

## Linear systems of equations

Linearly constrained least-squares problems

Linear dynamical systems

Eigenvalues and eigenvectors

The singular-value decomposition

# Linear systems of equations

system of linear equations:

$$\begin{array}{rclclclclcl} y_1 & = & A_{11} & x_1 & + & \cdots & + & A_{1n} & x_n \\ & & \vdots & & & & & & \\ y_m & = & A_{m1} & x_1 & + & \cdots & + & A_{mn} & x_n \end{array}$$

## Linear systems of equations

system of linear equations:

$$\begin{aligned}y_1 &= A_{11} x_1 + \cdots + A_{1n} x_n \\&\vdots \\y_m &= A_{m1} x_1 + \cdots + A_{mn} x_n\end{aligned}$$

matrix representation:

$$y = Ax$$

## Linear systems of equations

system of linear equations:

$$\begin{aligned} y_1 &= A_{11} x_1 + \cdots + A_{1n} x_n \\ &\vdots \\ y_m &= A_{m1} x_1 + \cdots + A_{mn} x_n \end{aligned}$$

matrix representation:

$$y = Ax$$

where

$$y = \begin{bmatrix} y_1 \\ \vdots \\ y_m \end{bmatrix}, \quad A = \begin{bmatrix} A_{11} & \cdots & A_{1n} \\ \vdots & \ddots & \vdots \\ A_{m1} & \cdots & A_{mn} \end{bmatrix}, \quad x = \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}$$

right side of system *defines* matrix-vector multiplication

Linear systems of equations

Linearly constrained least-squares problems

Linear dynamical systems

Eigenvalues and eigenvectors

The singular-value decomposition

## Linearly constrained least-squares problems

$$\begin{aligned} &\underset{x \in \mathbb{R}^n}{\text{minimize}} : \|Ax - b\| \\ &\text{subject to} : Cx = d \end{aligned}$$


## Linearly constrained least-squares problems

$$\begin{aligned} &\underset{x \in \mathbb{R}^n}{\text{minimize}} : \|Ax - b\| \\ &\text{subject to} : Cx = d \end{aligned}$$

normal equations:


$$\begin{bmatrix} A^T A & C^T \\ C & 0 \end{bmatrix} \begin{bmatrix} x \\ \lambda \end{bmatrix} = \begin{bmatrix} A^T b \\ d \end{bmatrix}$$

## The orthogonality principle






## The orthogonality principle



$$b - \hat{b} \perp A \text{ null}(C) \quad \Rightarrow \quad A^T(b - \hat{b}) \in \text{null}(C)^\perp = \text{range}(C^T)$$

## The orthogonality principle



$$b - \hat{b} \perp A \text{ null}(C) \quad \Rightarrow \quad A^T(b - \hat{b}) \in \text{null}(C)^\perp = \text{range}(C^T)$$

$$\begin{bmatrix} A^T A & C^T \\ C & 0 \end{bmatrix} \begin{bmatrix} x \\ \lambda \end{bmatrix} = \begin{bmatrix} A^T b \\ d \end{bmatrix} \quad \begin{array}{l} \text{orthogonality} \\ \text{feasibility} \end{array}$$

Linear systems of equations

Linearly constrained least-squares problems

Linear dynamical systems

Eigenvalues and eigenvectors

The singular-value decomposition

## Linear dynamical systems

discrete-time linear dynamical system:

$$\begin{aligned}x(t+1) &= Ax(t) + Bu(t), \\ y(t) &= Cx(t) + Du(t)\end{aligned}$$

## Linear dynamical systems

discrete-time linear dynamical system:

$$\begin{aligned}x(t+1) &= Ax(t) + Bu(t), \\ y(t) &= Cx(t) + Du(t)\end{aligned}$$

solutions of the state and measurement equations:

$$\begin{aligned}x(t) &= A^t x(0) + \sum_{\tau=0}^{t-1} A^{t-\tau-1} Bu(\tau), \\ y(t) &= CA^t x(0) + \sum_{\tau=0}^{t-1} CA^{t-\tau-1} Bu(\tau) + Du(t)\end{aligned}$$

Linear systems of equations

Linearly constrained least-squares problems

Linear dynamical systems

**Eigenvalues and eigenvectors**

The singular-value decomposition

## Eigenvalues and eigenvectors

suppose  $A \in \mathbb{R}^{n \times n}$

- ▶  $v \in \mathbb{C}^n$  is (right) eigenvector of  $A$  with eigenvalue  $\lambda \in \mathbb{C}$  if

$$Av = \lambda v, \quad v \neq 0$$

## Eigenvalues and eigenvectors

suppose  $A \in \mathbb{R}^{n \times n}$

- ▶  $v \in \mathbb{C}^n$  is (right) eigenvector of  $A$  with eigenvalue  $\lambda \in \mathbb{C}$  if

$$Av = \lambda v, \quad v \neq 0$$

- ▶ in matrix form:

$$A \begin{bmatrix} v_1 & \cdots & v_k \end{bmatrix} = \begin{bmatrix} v_1 & \cdots & v_k \end{bmatrix} \begin{bmatrix} \lambda_1 & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & \lambda_k \end{bmatrix}$$



## Eigenvalues and eigenvectors

- ▶  $w \in \mathbb{C}^n$  left eigenvector of  $A$  with eigenvalue  $\lambda \in \mathbb{C}$  if

$$w^T A = \lambda w^T$$

## Eigenvalues and eigenvectors

- ▶  $w \in \mathbb{C}^n$  left eigenvector of  $A$  with eigenvalue  $\lambda \in \mathbb{C}$  if

$$w^T A = \lambda w^T$$

- ▶ in matrix form:

$$\begin{bmatrix} w_1^T \\ \vdots \\ w_k^T \end{bmatrix} A = \begin{bmatrix} \lambda_1 & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & \lambda_k \end{bmatrix} \begin{bmatrix} w_1^T \\ \vdots \\ w_k^T \end{bmatrix}$$

## Diagonalizable matrices

- ▶ A diagonalizable if there is linearly independent set of  $n$  eigenvectors:

$$A = \begin{bmatrix} v_1 & \cdots & v_n \end{bmatrix} \begin{bmatrix} \lambda_1 & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & \lambda_n \end{bmatrix} \begin{bmatrix} w_1^T \\ \vdots \\ w_n^T \end{bmatrix}$$
$$= \sum_{i=1}^n \lambda_i v_i w_i^T$$

## Diagonalizable matrices

- ▶ A diagonalizable if there is linearly independent set of  $n$  eigenvectors:

$$\begin{aligned} A &= \begin{bmatrix} v_1 & \cdots & v_n \end{bmatrix} \begin{bmatrix} \lambda_1 & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & \lambda_n \end{bmatrix} \begin{bmatrix} w_1^T \\ \vdots \\ w_n^T \end{bmatrix} \\ &= \sum_{i=1}^n \lambda_i v_i w_i^T \end{aligned}$$

- ▶ where

$$\begin{bmatrix} w_1^T \\ \vdots \\ w_n^T \end{bmatrix} = \begin{bmatrix} v_1 & \cdots & v_n \end{bmatrix}^{-1}$$

## Dominant-eigenvalue analysis

- ▶ order eigenvalues such that  $|\lambda_1| \geq \dots \geq |\lambda_n|$

## Dominant-eigenvalue analysis

- ▶ order eigenvalues such that  $|\lambda_1| \geq \dots \geq |\lambda_n|$
- ▶  $\lambda_1$  is unique dominant eigenvalue if  $|\lambda_1| > |\lambda_2|$

## Dominant-eigenvalue analysis

- ▶ order eigenvalues such that  $|\lambda_1| \geq \dots \geq |\lambda_n|$
- ▶  $\lambda_1$  is unique dominant eigenvalue if  $|\lambda_1| > |\lambda_2|$
- ▶ then, for large  $t$ ,

$$A^t x \sim \lambda_1^t v_1 w_1^T x$$

assuming  $w_1^T x \neq 0$

## Dominant-eigenvalue analysis

- ▶ order eigenvalues such that  $|\lambda_1| \geq \dots \geq |\lambda_n|$
- ▶  $\lambda_1$  is unique dominant eigenvalue if  $|\lambda_1| > |\lambda_2|$
- ▶ then, for large  $t$ ,

$$A^t x \sim \lambda_1^t v_1 w_1^T x$$

assuming  $w_1^T x \neq 0$

- ▶ for example, if  $x(t+1) = Ax(t)$ , then

$$\lim_{t \rightarrow \infty} \frac{\|x(t+1)\|}{\|x(t)\|} = |\lambda_1|$$

assuming  $w_1^T x(0) \neq 0$



## Dominant-eigenvalue analysis

- ▶ order eigenvalues such that  $|\lambda_1| \geq \dots \geq |\lambda_n|$
- ▶  $\lambda_1$  is unique dominant eigenvalue if  $|\lambda_1| > |\lambda_2|$
- ▶ then, for large  $t$ ,

$$A^t x \sim \lambda_1^t v_1 w_1^T x$$

assuming  $w_1^T x \neq 0$

- ▶ for example, if  $x(t+1) = Ax(t)$ , then

$$\lim_{t \rightarrow \infty} \frac{\|x(t+1)\|}{\|x(t)\|} = |\lambda_1|$$

assuming  $w_1^T x(0) \neq 0$

- ▶ if  $|\lambda_1| > |\lambda_2| > |\lambda_3|$ , then  $\lambda_2$  is “vice” dominant eigenvalue

## Quadratic forms

- ▶ quadratic form is function of the form

$$x^T A x = \sum_{i=1}^n \sum_{j=1}^n A_{ij} x_i x_j$$

## Quadratic forms

- ▶ quadratic form is function of the form

$$x^T A x = \sum_{i=1}^n \sum_{j=1}^n A_{ij} x_i x_j$$

- ▶ unique representation using symmetric matrix  $A \in \mathbb{S}^n$

## Symmetric matrices

suppose  $A \in \mathbb{R}^{n \times n}$  is symmetric

## Symmetric matrices

suppose  $A \in \mathbb{R}^{n \times n}$  is symmetric

- ▶ there is an orthonormal set of  $n$  eigenvectors:

$$A = Q\Lambda Q^T = \sum_{i=1}^n \lambda_i q_i q_i^T$$

## Symmetric matrices

suppose  $A \in \mathbb{R}^{n \times n}$  is symmetric

- ▶ there is an orthonormal set of  $n$  eigenvectors:

$$A = Q\Lambda Q^T = \sum_{i=1}^n \lambda_i q_i q_i^T$$

- ▶  $A$  called positive definite if  $x^T A x > 0$  for all  $x \neq 0$

## Symmetric matrices

suppose  $A \in \mathbb{R}^{n \times n}$  is symmetric

- ▶ there is an orthonormal set of  $n$  eigenvectors:

$$A = Q\Lambda Q^T = \sum_{i=1}^n \lambda_i q_i q_i^T$$

- ▶  $A$  called positive definite if  $x^T A x > 0$  for all  $x \neq 0$ 
  - ▶ if and only if all  $\lambda_i > 0$

## Symmetric matrices

suppose  $A \in \mathbb{R}^{n \times n}$  is symmetric

- ▶ there is an orthonormal set of  $n$  eigenvectors:

$$A = Q\Lambda Q^T = \sum_{i=1}^n \lambda_i q_i q_i^T$$

- ▶  $A$  called positive definite if  $x^T A x > 0$  for all  $x \neq 0$ 
  - ▶ if and only if all  $\lambda_i > 0$
- ▶  $A$  called positive semidefinite if  $x^T A x \geq 0$  for all  $x$



## Symmetric matrices

suppose  $A \in \mathbb{R}^{n \times n}$  is symmetric

- ▶ there is an orthonormal set of  $n$  eigenvectors:

$$A = Q\Lambda Q^T = \sum_{i=1}^n \lambda_i q_i q_i^T$$

- ▶  $A$  called positive definite if  $x^T A x > 0$  for all  $x \neq 0$ 
  - ▶ if and only if all  $\lambda_i > 0$
- ▶  $A$  called positive semidefinite if  $x^T A x \geq 0$  for all  $x$ 
  - ▶ if and only if all  $\lambda_i \geq 0$

## Extremal-trace problems

$$\begin{aligned} &\underset{Q \in \mathbb{R}^{n \times k}}{\text{maximize}} : \text{tr}(Q^T A Q) \\ &\text{subject to} : Q^T Q = I \end{aligned}$$

## Extremal-trace problems

$$\begin{aligned} & \text{maximize} : \text{tr}(Q^T A Q) \\ & \quad Q \in \mathbb{R}^{n \times k} \\ & \text{subject to} : Q^T Q = I \end{aligned}$$

- equivalent formulation:

$$\begin{aligned} & \text{maximize} : \sum_{i=1}^k q_i^T A q_i \\ & \quad q_1, \dots, q_k \in \mathbb{R}^n \\ & \text{subject to} : q_i^T q_j = \delta_{ij} \end{aligned}$$

## Extremal-trace problems

$$\begin{aligned} & \underset{Q \in \mathbb{R}^{n \times k}}{\text{maximize}} : \text{tr}(Q^T A Q) \\ & \text{subject to} : Q^T Q = I \end{aligned}$$

- equivalent formulation:

$$\begin{aligned} & \underset{q_1, \dots, q_k \in \mathbb{R}^n}{\text{maximize}} : \sum_{i=1}^k q_i^T A q_i \\ & \text{subject to} : q_i^T q_j = \delta_{ij} \end{aligned}$$

- solution is matrix  $Q$  whose columns are eigenvectors of  $A$  corresponding to  $k$  largest eigenvalues

## Extremal-trace problems

$$\begin{aligned} & \text{maximize} : \text{tr}(Q^T A Q) \\ & \quad Q \in \mathbb{R}^{n \times k} \\ & \text{subject to} : Q^T Q = I \end{aligned}$$

- ▶ equivalent formulation:

$$\begin{aligned} & \text{maximize} : \sum_{i=1}^k q_i^T A q_i \\ & \quad q_1, \dots, q_k \in \mathbb{R}^n \\ & \text{subject to} : q_i^T q_j = \delta_{ij} \end{aligned}$$

- ▶ solution is matrix  $Q$  whose columns are eigenvectors of  $A$  corresponding to  $k$  largest eigenvalues
- ▶ to minimize, take matrix whose columns are eigenvectors corresponding to  $k$  smallest eigenvalues

Linear systems of equations

Linearly constrained least-squares problems

Linear dynamical systems

Eigenvalues and eigenvectors

The singular-value decomposition

## The singular-value decomposition

- ▶  $A^T A$  and  $A A^T$  are symmetric and positive semidefinite

## The singular-value decomposition

- ▶  $A^T A$  and  $AA^T$  are symmetric and positive semidefinite
- ▶ eigenvalue decompositions:

$$A^T A = V \Sigma^2 V^T, \quad AA^T = U \Sigma^2 U^T$$



## The singular-value decomposition

- ▶  $A^T A$  and  $AA^T$  are symmetric and positive semidefinite
- ▶ eigenvalue decompositions:

$$A^T A = V \Sigma^2 V^T, \quad AA^T = U \Sigma^2 U^T$$

- ▶ eigenvalues are nonnegative

## The singular-value decomposition

- ▶  $A^T A$  and  $AA^T$  are symmetric and positive semidefinite
- ▶ eigenvalue decompositions:

$$A^T A = V \Sigma^2 V^T, \quad AA^T = U \Sigma^2 U^T$$

- ▶ eigenvalues are nonnegative
- ▶ eigenvalues are the same

## The singular-value decomposition

- ▶  $A^T A$  and  $AA^T$  are symmetric and positive semidefinite
- ▶ eigenvalue decompositions:

$$A^T A = V \Sigma^2 V^T, \quad AA^T = U \Sigma^2 U^T$$

- ▶ eigenvalues are nonnegative
  - ▶ eigenvalues are the same
- ▶ singular-value decomposition:

$$A = U \Sigma V^T$$

## The singular-value decomposition (matrix form)

$$A = U\Sigma V^T = \sum_{i=1}^r \sigma_i u_i v_i^T$$

## The singular-value decomposition (matrix form)

$$A = U\Sigma V^T = \sum_{i=1}^r \sigma_i u_i v_i^T$$

- $U \in \mathbb{R}^{m \times r}$  has orthonormal columns

## The singular-value decomposition (matrix form)

$$A = U\Sigma V^T = \sum_{i=1}^r \sigma_i u_i v_i^T$$

- ▶  $U \in \mathbb{R}^{m \times r}$  has orthonormal columns
- ▶  $\Sigma \in \mathbb{R}^{r \times r}$  diagonal, nonsingular

## The singular-value decomposition (matrix form)

$$A = U\Sigma V^T = \sum_{i=1}^r \sigma_i u_i v_i^T$$

- ▶  $U \in \mathbb{R}^{m \times r}$  has orthonormal columns
- ▶  $\Sigma \in \mathbb{R}^{r \times r}$  diagonal, nonsingular
- ▶  $V \in \mathbb{R}^{n \times r}$  has orthonormal columns

# Ellipsoids

- ▶ generate members:

$$\{x_0 + Az : \|z\| \leq 1\}$$



## Ellipsoids

- ▶ generate members:

$$\{x_0 + Az : \|z\| \leq 1\}$$

- ▶ test membership:

$$\{x \in \mathbb{R}^n : (x - x_0)^T S (x - x_0) \leq 1\}$$

## Ellipsoids

- ▶ generate members:

$$\{x_0 + Az : \|z\| \leq 1\}$$

- ▶ test membership:

$$\{x \in \mathbb{R}^n : (x - x_0)^T S (x - x_0) \leq 1\}$$

- ▶ equivalent representations:  $S = (AA^T)^{-1}$

## Ellipsoids

- ▶ generate members:

$$\{x_0 + Az : \|z\| \leq 1\}$$

- ▶ test membership:

$$\{x \in \mathbb{R}^n : (x - x_0)^T S (x - x_0) \leq 1\}$$

- ▶ equivalent representations:  $S = (AA^T)^{-1}$
- ▶ image of unit ball under  $A$  is ellipsoid with principal axes  $\sigma_i u_i$

## The singular-value decomposition (dyadic expansion)

$$A = \sum_{i=1}^r \sigma_i u_i v_i^T$$

- ▶  $\sigma_1 \geq \dots \geq \sigma_r > 0$  are singular values
- ▶  $u_1, \dots, u_r \in \mathbb{R}^m$  are output singular vectors
- ▶  $v_1, \dots, v_r \in \mathbb{R}^n$  are input singular vectors
- ▶  $u_i v_i^T$  is unit atom
  - ▶  $\text{rank}(u_i v_i^T) = 1$
  - ▶  $\|u_i v_i^T\| = 1$
- ▶ SVD decomposes  $A$  into sum of unit atoms
  - ▶ singular values rank atoms in terms of importance

## The pseudoinverse

$$A^\dagger = V\Sigma^{-1}U^\top = \sum_{i=1}^r \frac{1}{\sigma_i} v_i u_i^\top$$

## The pseudoinverse

$$A^\dagger = V\Sigma^{-1}U^T = \sum_{i=1}^r \frac{1}{\sigma_i} v_i u_i^T$$

- ▶  $A^\dagger b$  is least-norm least-squares vector

$$\text{minimize : } \|x\|$$

$x \in \mathbb{R}^n$

$$\text{subject to : } \|Ax - b\| = \min_z \|Az - b\|$$

## Extremal-trace problems (input)

$$\begin{aligned} &\text{maximize : } \text{tr}(Q^T A^T A Q) \\ &\quad Q \in \mathbb{R}^{n \times k} \\ &\text{subject to : } Q^T Q = I \end{aligned}$$

## Extremal-trace problems (input)

$$\begin{aligned} &\text{maximize : } \text{tr}(Q^T A^T A Q) \\ &Q \in \mathbb{R}^{n \times k} \\ &\text{subject to : } Q^T Q = I \end{aligned}$$

- equivalent formulation:

$$\begin{aligned} &\text{maximize : } \sum_{i=1}^k \|Aq_i\|^2 \\ &q_1, \dots, q_k \in \mathbb{R}^n \\ &\text{subject to : } q_i^T q_j = \delta_{ij} \end{aligned}$$



## Extremal-trace problems (input)

$$\begin{aligned} &\text{maximize} : \text{tr}(Q^T A^T A Q) \\ &Q \in \mathbb{R}^{n \times k} \\ &\text{subject to} : Q^T Q = I \end{aligned}$$

- equivalent formulation:

$$\begin{aligned} &\text{maximize} : \sum_{i=1}^k \|Aq_i\|^2 \\ &q_1, \dots, q_k \in \mathbb{R}^n \\ &\text{subject to} : q_i^T q_j = \delta_{ij} \end{aligned}$$

- solution is matrix  $Q$  whose columns are eigenvectors of  $A^T A = V \Sigma^2 V^T$  corresponding to  $k$  largest eigenvalues

## Extremal-trace problems (input)

$$\begin{aligned} & \text{maximize : } \text{tr}(Q^T A^T A Q) \\ & \quad Q \in \mathbb{R}^{n \times k} \\ & \text{subject to : } Q^T Q = I \end{aligned}$$

- ▶ equivalent formulation:

$$\begin{aligned} & \text{maximize : } \sum_{i=1}^k \|Aq_i\|^2 \\ & \quad q_1, \dots, q_k \in \mathbb{R}^n \\ & \text{subject to : } q_i^T q_j = \delta_{ij} \end{aligned}$$

- ▶ solution is matrix  $Q$  whose columns are eigenvectors of  $A^T A = V \Sigma^2 V^T$  corresponding to  $k$  largest eigenvalues
- ▶ equivalently, solution is matrix  $Q$  whose columns are the first  $k$  input singular vectors of  $A$

## Extremal-trace problems (output)

$$\begin{aligned} &\text{maximize : } \text{tr}(Q^T A A^T Q) \\ &Q \in \mathbb{R}^{m \times k} \\ &\text{subject to : } Q^T Q = I \end{aligned}$$

## Extremal-trace problems (output)

$$\begin{aligned} &\text{maximize : } \text{tr}(Q^T A A^T Q) \\ &Q \in \mathbb{R}^{m \times k} \\ &\text{subject to : } Q^T Q = I \end{aligned}$$

► equivalent formulation:

$$\begin{aligned} &\text{maximize}_{q_1, \dots, q_k \in \mathbb{R}^m} : \sum_{i=1}^k \sum_{j=1}^n \|q_i q_i^T A_{*j}\|^2 \\ &\text{subject to} : q_i^T q_j = \delta_{ij} \end{aligned}$$

## Extremal-trace problems (output)

$$\begin{aligned} & \text{maximize : } \text{tr}(Q^T A A^T Q) \\ & \quad Q \in \mathbb{R}^{m \times k} \\ & \text{subject to : } Q^T Q = I \end{aligned}$$

- ▶ equivalent formulation:

$$\begin{aligned} & \text{maximize}_{q_1, \dots, q_k \in \mathbb{R}^m} : \sum_{i=1}^k \sum_{j=1}^n \|q_i q_i^T A_{*j}\|^2 \\ & \text{subject to} : q_i^T q_j = \delta_{ij} \end{aligned}$$

- ▶ solution is matrix  $Q$  whose columns are eigenvectors of  $AA^T = U\Sigma^2U^T$  corresponding to  $k$  largest eigenvalues

## Extremal-trace problems (output)

$$\begin{aligned} & \text{maximize : } \text{tr}(Q^T A A^T Q) \\ & \quad Q \in \mathbb{R}^{m \times k} \\ & \text{subject to : } Q^T Q = I \end{aligned}$$

- ▶ equivalent formulation:

$$\begin{aligned} & \text{maximize}_{q_1, \dots, q_k \in \mathbb{R}^m} : \sum_{i=1}^k \sum_{j=1}^n \|q_i q_i^T A_{*j}\|^2 \\ & \text{subject to} : q_i^T q_j = \delta_{ij} \end{aligned}$$

- ▶ solution is matrix  $Q$  whose columns are eigenvectors of  $A A^T = U \Sigma^2 U^T$  corresponding to  $k$  largest eigenvalues
- ▶ equivalently, solution is matrix  $Q$  whose columns are the first  $k$  output singular vectors of  $A$

## Low-rank approximation

singular-value decomposition of  $A \in \mathbb{R}^{m \times n}$ :

$$A = \sum_{i=1}^r \sigma_i u_i v_i^T$$

## Low-rank approximation

singular-value decomposition of  $A \in \mathbb{R}^{m \times n}$ :

$$A = \sum_{i=1}^r \sigma_i u_i v_i^T$$

best rank- $p$  approximation of  $A$ :

$$\hat{A}_p = \sum_{i=1}^p \sigma_i u_i v_i^T$$