




Spectral Classification of White Dwarfs by Dimensionality Reduction

Xander Byrne


Amy Bonsor – Laura Rogers – Christopher Manser

Outline

- Spectroscopic surveys
- White dwarfs
- Dimensionality reduction (DR)
- Variation: focus on spectral lines




Spectroscopic Surveys




Spectroscopic Surveys

DESI




Spectroscopic Surveys

DESI




GALAH

White Dwarfs 101




White Dwarfs 101

- 97% of stars will be WDs ($M \sim M_{\odot}$, $R \sim R_E$)





White Dwarfs 101

- 97% of stars will be WDs ($M \sim M_{\odot}$, $R \sim R_E$)
- Spectral types:





White Dwarfs 101

- 97% of stars will be WDs ($M \sim M_{\odot}$, $R \sim R_E$)
- Spectral types:
 - **DA** (hydrogen)





White Dwarfs 101

- 97% of stars will be WDs ($M \sim M_{\odot}$, $R \sim R_E$)
- Spectral types:
 - **DA** (hydrogen)
 - **DB** (helium)





White Dwarfs 101

- 97% of stars will be WDs ($M \sim M_{\odot}$, $R \sim R_E$)
- Spectral types:
 - **DA** (hydrogen)
 - **DB** (helium)
 - **DZ** (Ca, Mg, ...)





White Dwarfs 101

- 97% of stars will be WDs ($M \sim M_{\odot}$, $R \sim R_E$)
- Spectral types:
 - **DA** (hydrogen)
 - **DB** (helium)
 - **DZ** (Ca, Mg, ...)
 - ...



White Dwarfs 101

- 97% of stars will be WDs ($M \sim M_{\odot}$, $R \sim R_E$)
- Spectral types:
 - **DA** (hydrogen)
 - **DB** (helium)
 - **DZ** (Ca, Mg, ...)
 - ...
 - **CV** (double-peaked)




~{ Interlude: Astrocolonialism }~

- DESI is on the Mayall 4m telescope at **Iolkam Du'ag / Kitt Peak**
- This site is leased (dubiously?) from the **Tohono O'odham Nation**
- astrobites:
'A tale of two observatories'





DESI EDR – 4000 WD candidates

DESI EDR – 4000 WD candidates



DESI EDR – 4000 WD candidates



Dimensionality Reduction (DR)


- An unsupervised method of **simplifying a high-D dataset**

Dimensionality Reduction (DR)

- An unsupervised method of **simplifying a high-D dataset**
- Deprojects dataset into **2D map**, preserving distances


Dimensionality Reduction (DR)

- An unsupervised method of **simplifying a high-D dataset**
- Deprojects dataset into **2D map**, preserving distances




Dimensionality Reduction (DR)

- An unsupervised method of **simplifying a high-D dataset**
- Deprojects dataset into **2D map**, preserving distances





Dimensionality Reduction (DR)

- An unsupervised method of **simplifying a high-D dataset**
- Deprojects dataset into **2D map**, preserving distances



DR on a spectroscopic survey


- Each data point (e.g. spectrum) **converted to a vector**:



(1.93, 2.01, ..., -0.32)

DR on a spectroscopic survey


- Each data point (e.g. spectrum) **converted to a vector**:



- => a dataset of ~8000D vectors


DR on a spectroscopic survey


- Each data point (e.g. spectrum) **converted to a vector**:



(1.93, 2.01, ..., -0.32)

- => a dataset of ~8000D vectors, arranged by type





Advantages over visual classif'n

- **V. fast** (5.5s for $N \approx 4000$) and scales as $O(N \log N)$



Advantages over visual classif'n

- **V. fast** (5.5s for $N \approx 4000$) and scales as $O(N \log N)$
- => Much more **reproducible / modifiable**

Advantages over visual classif'n


- **V. fast** (5.5s for $N \approx 4000$) and scales as $O(N \log N)$
- => Much more **reproducible / modifiable**
- **Objective** – doesn't rely on human interpretation

Focusing on spectral lines




(1.93, 2.01, ..., -0.32)


Focusing on spectral lines



(1.63, 1.61, ..., 0.89)




Focusing on spectral lines




- Vectors now clustered based on a spectral feature



Focusing on spectral lines



(1.63, 1.61, ..., 0.89)

- Vectors now clustered based on a spectral feature
- Removes 'distraction' of rest of the spectrum





Byrne+, in rev.


Conclusions

- Spectroscopic surveys return valuable data for WD science
- Dimensionality reduction is a useful classification tool for spectroscopic surveys
- Focusing on spectral lines can improve classification power




Extra slides


Preprocessing





- Can **spot mistakes**



- Can **spot mistakes**



- Can **spot mistakes**




Automated WD Classification

Automated WD Classification

- **Supervised** methods rely on a training set
 - Have shown promise in WD classification (Vincent+23, 24)



Automated WD Classification

- **Supervised** methods rely on a training set
 - Have shown promise in WD classification (Vincent+23, 24)
- But can suffer from **class imbalance**
 - ...which always exists for WDs



Automated WD Classification

- **Supervised** methods rely on a training set
 - Have shown promise in WD classification (Vincent+23, 24)
- But can suffer from **class imbalance**
 - ...which always exists for WDs
- Also, sometimes **weirdly confident**



Jonathan Ramkissoon


Unsupervised Methods

- Don't rely on a training set!



Unsupervised Methods

- Don't rely on a training set!
- Looks for **inherent structure** in a dataset (trends, clusters, ...)



Unsupervised Methods

- **Don't rely on a training set!**
- Looks for **inherent structure** in a dataset (trends, clusters, ...)
- Usually involves **comparing data points to each other**

