

WEEK 5 HOMEWORK

INSTRUCTIONS

- Every learner should submit his/her own homework solutions. However, you are allowed to discuss the homework with each other (in fact, I encourage you to form groups and/or use the forums) – but everyone must submit his/her own solution; you may not copy someone else's solution.
- The homework will be peer-graded.
- The homework grading scale reflects the fact that the primary purpose of homework is learning:

| Rating | Meaning | Point value (out of 100) |
|--------|--|-----------------------------|
| 4 | All correct (perhaps except a few details) <u>with</u> a deeper solution than expected | 100 |
| 3 | Most or all correct | 90 |
| 2 | Not correct, but a reasonable attempt | 75 |
| 1 | Not correct, insufficient effort | 50 |
| 0 | Not submitted | 0 |

Question 1

Using the crime data set from Homework 3, build a regression model using:

1. Stepwise regression
2. Lasso
3. Elastic net

For Parts 2 and 3, remember to scale the data first – otherwise, the regression coefficients will be on different scales and the constraint won't have the desired effect.

For Parts 2 and 3, use the `glmnet` function in R.

Notes on R:

- For the elastic net model, what we called λ in the videos, `glmnet` calls "alpha"; you can get a range of results by varying alpha from 1 (lasso) to 0 (ridge regression) [and, of course, other values of alpha in between].
- In a function call like `glmnet(x,y,family="mgaussian",alpha=1)` the predictors `x` need to be in R's matrix format, rather than data frame format. You can convert a data frame to a matrix using `as.matrix` – for example, `x <- as.matrix(data[,1:n-1])`
- Rather than specifying a value of `T`, `glmnet` returns models for a variety of values of `T`.

Question 2

Describe a situation or problem from your job, everyday life, current events, etc., for which a design of experiments approach would be appropriate.

Question 3

To determine the value of 10 different yes/no features to the market value of a house (large yard, solar roof, etc.), a real estate agent plans to survey 50 potential buyers, showing a fictitious house with different combinations of features. To reduce the survey size, the agent wants to show just 16 fictitious houses. Use R's FrF2 function (in the FrF2 package) to find a fractional factorial design for this experiment: what set of features should each of the 16 fictitious houses? Note: the output of FrF2 is "1" (include) or "-1" (don't include) for each feature.

Question 4

For each of the following distributions, give an example of data that you would expect to follow this distribution (besides the examples already discussed in class).

- a. Binomial
- b. Geometric
- c. Poisson
- d. Exponential
- e. Weibull