




Calculus Solution



Contents

I	Antiderivatives
1	The Indefinite Integral 4
2	Chain Rule in Reverse 10
3	The Method of Substitution 14



Antiderivatives

1	The Indefinite Integral	4
1.1	The Indefinite Integral	
1.2	Basic Antiderivative Rules	
2	Chain Rule in Reverse	10
2.1	Chain Rule in Reverse	
3	The Method of Substitution	14
3.1	The Method of Substitution	



1. The Indefinite Integral

1.1 The Indefinite Integral

How do we undo a derivative? If we were given the derivative of a function $f'(x)$, how could we find the original function $f(x)$? The answer is called the antiderivative of $f(x)$, which we will denote by the associated capital letter $F(x)$.

Another way to think about this question is: "What function do I have to take the derivative of in order to get the answer?" The antiderivative of $f(x) = 2x$ is x^2 .

But what about $F(x) = x^2 + 1$? This works too! In fact, since when we take the derivative of a constant, we get zero. We could have chosen any constant. As a result, we report our antiderivative in its most general form $x^2 + C$. The constant C is an important part of the antiderivative.

Notationally, we denote the operation of take the antiderivative (known as integration) as:

$$\int 2x \, dx = x^2 + C$$

This is also called the indefinite integral of the function $f(x)$, or sometimes just the integral of $f(x)$, where $f(x)$ is called the integrand.

That was a pretty simple example, so how do we find antiderivatives of more complicated expressions? In much the same way as we did with derivatives, we can generate a set of rules for finding antiderivatives, derived simply by thinking of our familiar derivative rules in reverse.

1.2 Basic Antiderivative Rules

The power rule for derivatives multiplies by the power and then subtracts one from the power. Reversing these operations means that we add one to the power and divide by the new power.

1.2.1 Reverse Power Rule

Theorem 1.2.1 — Reverse Power Rule.

$$\int x^n dx = \frac{x^{n+1}}{n+1} + C, \text{ where } n \neq -1 \quad (1.1)$$

Exercise 1.1 Find the antiderivatives.

(a)

$$\begin{aligned} \int x^6 dx \\ = \frac{x^7}{7} + C \end{aligned}$$

(b)

$$\begin{aligned} \int \sqrt[4]{t} dt \\ = \int t^{\frac{1}{4}} dt \\ = \frac{t^{5/4}}{5/4} + C \\ = \frac{4t^{5/4}}{5} + C \end{aligned}$$

(c)

$$\begin{aligned} \int \frac{1}{x^{5/3}} dx \\ = \int x^{-5/3} dx \\ = \frac{x^{-2/3}}{-2/3} + C \\ = -\frac{3}{2x^{2/3}} + C \end{aligned}$$

1.2.2 Antiderivative of Zero

Theorem 1.2.2 — Antiderivative of Zero.

$$\int 0 \, dx = C \quad (1.2)$$

1.2.3 Antiderivative of a Constant

Theorem 1.2.3 — Antiderivative of a Constant.

$$\int k \, dx = kx + C, \text{ where } k \text{ is any constant} \quad (1.3)$$

Exercise 1.2 Find the antiderivatives.

$$\begin{aligned} \int \pi \, dx \\ = \pi x + C \end{aligned}$$

1.2.4 Multiplicative Constants

Theorem 1.2.4 — Multiplicative Constantst.

$$\int k f(x) \, dx = k \int f(x) \, dx \quad (1.4)$$

Exercise 1.3 Find the antiderivatives.

(a)

$$\begin{aligned} \int 4x^7 \, dx \\ = 4 \int x^7 \, dx \\ = 4 \left(\frac{x^8}{8} + C \right) \\ = \frac{x^8}{2} + C \end{aligned}$$

(b)

$$\begin{aligned}\int \frac{\pi}{\sqrt{t}} dt \\&= \pi \int t^{-\frac{1}{2}} dt \\&= \pi \cdot \frac{t^{1/2}}{1/2} + C \\&= 2\pi t^{1/2} + C\end{aligned}$$

1.2.5 Sum / Difference

Theorem 1.2.5 — Sum / Difference.

$$\int (f(x) \pm g(x)) dx = \int f(x) dx \pm \int g(x) dx \quad (1.5)$$

Exercise 1.4 Find the antiderivatives.

(a)

$$\begin{aligned}\int (3x^2 + 5) dx \\&= \frac{3x^3}{3} + 5x + C \\&= x^3 + 5x + C\end{aligned}$$

(b)

$$\begin{aligned}\int \left(\frac{1}{x^3} - \frac{2}{x^2} \right) dx \\&= \int (x^{-3} - 2x^{-2}) dx \\&= \frac{x^{-2}}{-2} - 2 \left(\frac{x^{-1}}{-1} \right) + C \\&= -\frac{1}{2x^2} + \frac{2}{x} + C\end{aligned}$$

1.2.6 Trigonometric Functions

Theorem 1.2.6 — Trigonometric Functions.

$$\int \sin(x) \, dx = -\cos(x) + C \quad (1.6)$$

$$\int \cos(x) \, dx = \sin(x) + C \quad (1.7)$$

$$\int \csc^2(x) \, dx = -\cot(x) + C \quad (1.8)$$

$$\int \sec^2(x) \, dx = \tan(x) + C \quad (1.9)$$

$$\int \sec(x)\tan(x) \, dx = \sec(x) + C \quad (1.10)$$

$$\int \csc(x)\cot(x) \, dx = -\csc(x) + C \quad (1.11)$$

1.2.7 Exponential / Logarithmic**Theorem 1.2.7 — Exponential / Logarithmic.**

$$\int a^x \ln(a) \, dx = a^x + C \quad (1.12)$$

$$\int e^x \ln(e) \, dx = e^x + C \quad (1.13)$$

$$\int \frac{1}{x \ln(a)} \, dx = \log_a |x| + C \quad (1.14)$$

$$\int \frac{1}{x} \, dx = \ln|x| + C \quad (1.15)$$

Exercise 1.5 Find the antiderivatives.

(a)

$$\begin{aligned} \int 4^x \ln(4) + 5e^x - \frac{6}{x} \, dx \\ = 4^x + 5e^x - 6\ln|x| + C \end{aligned}$$

(b)

$$\begin{aligned} \int 3^z \, dz \\ = \frac{1}{\ln(3)} \int 3^z \ln(3) \, dz \\ = \frac{1}{\ln(3)} \cdot 3^z + C \end{aligned}$$


Sometimes we need to manipulate the integral a little bit before we can apply the rules. ■

Exercise 1.6 Find the antiderivatives.

(a)

$$\begin{aligned} & \int \frac{2s^3 - 5s^4}{3s^2} ds \\ &= \int \frac{2}{3}s - \frac{5}{3}s^2 ds \\ &= \frac{2}{3} \cdot \frac{s^2}{2} - \frac{5}{3} \cdot \frac{s^3}{3} + C \\ &= \frac{s^2}{3} - \frac{5s^3}{9} + C \end{aligned}$$

(b)

$$\begin{aligned} & \int \left(\frac{1}{x} + \frac{1}{x^2} \right) (3 + 2x^2) dx \\ &= \int \frac{3}{x} + 2x + \frac{3}{x^2} + 2 dx \\ &= 3\ln|x| + x^2 + 3 \cdot \frac{x^{-1}}{-1} + 2x + C \\ &= 3\ln|x| + x^2 + \frac{3}{x} + 2x + C \end{aligned}$$


2. Chain Rule in Reverse

2.1 Chain Rule in Reverse

The derivative of $f(u(x))$ is $f'(u(x))u'(x)$, so

Theorem 2.1.1 — Chain Rule in Reverse.

$$\int f'(u(x))u'(x) dx = f(u(x)) + C \quad (2.1)$$

Notice that in the integration, the $u'(x)$ piece disappears, being absorbed back into $f(x)$. The steps for finding the antiderivative of composition functions are as follows:

1. Identify the core layer $u(x)$.
2. Identify the derivative of the core layer $u'(x)$.
3. Identify the outer layer f' , and integrate f' leaving $u(x)$ inside.

Exercise 2.1 Find the antiderivatives.

(a)

$$\int (6x^2 + 1) \sin(2x^3 + x) \, dx$$

$$u = 2x^3 + x$$

$$u' = 6x^2 + 1$$

$$\begin{aligned} \int (6x^2 + 1) \sin(2x^3 + x) \, dx \\ = -\cos(2x^3 + x) + C \end{aligned}$$

(b)

$$\int \sec^2(4t) \, dt$$

$$u = 4t$$

$$u' = 4$$

$$\begin{aligned} \int \sec^2(4t) \, dt \\ = \frac{1}{4} \int 4 \sec^2(4t) \, dt \\ = \frac{1}{4} \tan(4t) + C \end{aligned}$$

(c)

$$\int 4x^3(3x^4 - 1)^{14} dx$$

$$u = 3x^4 - 1$$

$$u' = 12x^3$$

$$\begin{aligned} & \int 4x^3(3x^4 - 1)^{14} dx \\ &= \frac{1}{3} \int 12x^3(3x^4 - 1)^{14} dx \\ &= \frac{1}{3} \cdot \frac{(3x^4 - 1)^{15}}{15} + C \\ &= \frac{(3x^4 - 1)^{15}}{45} + C \end{aligned}$$

(d)

$$\int \frac{e^{\frac{1}{x}}}{4x^2} dx$$

$$u = \frac{1}{x}$$

$$u' = -\frac{1}{x^2}$$

$$\begin{aligned} & \int \frac{e^{\frac{1}{x}}}{4x^2} dx \\ &= \frac{1}{4} \int \frac{1}{x^2} e^{\frac{1}{x}} dx \\ &= -\frac{1}{4} \int -\frac{1}{x^2} e^{\frac{1}{x}} dx \\ &= -\frac{1}{4} e^{\frac{1}{x}} + C \end{aligned}$$

■

Exercise 2.2 Integrate in one step.

$$\begin{aligned} & \int e^{-2t} + \sin(3t) + \cos\left(\frac{1}{4}t\right) dt \\ &= -\frac{1}{2}e^{-2t} - \frac{1}{3}\cos(3t) + 4\sin\left(\frac{1}{4}t\right) + C \end{aligned}$$



3. The Method of Substitution

3.1 The Method of Substitution

The idea behind the method of substitution is to change a difficult integral in terms of one variable into an easier integral in terms of some other variable using a substitution.

Theorem 3.1.1 — The Method of Substitution.

$$\int f'(u(x)) \frac{du}{dx} = \int f'(u) du \quad (3.1)$$

Exercise 3.1 The method of substitution.

$$\int (6x + 4)(3x^2 + 4x)^5 dx$$

1. Identify the core layer $u(x) = 3x^2 + 4x$.
2. Find the derivative of the core $\frac{du}{dx} = 6x + 4$.
3. Transform from an integral in x to an integral in the new variable u using the change of variable theorem.

$$\begin{aligned}
 & \int (6x+4)(3x^2+4x)^5 dx \\
 &= \int \frac{du}{dx} u^5 dx \\
 &= \int u^5 du \\
 &= \frac{u^6}{6} + C
 \end{aligned}$$

4. Convert back to the original variable by substituting $u(x)$ back in.

$$\begin{aligned}
 & \frac{u^6}{6} + C \\
 &= \frac{(3x^2+4x)^6}{6} + C
 \end{aligned}$$

Exercise 3.2 Calculate using the method of substitution.

(a)