


NUCLEO-OLIVARY INHIBITION EXPLAINS EXTINCTION ON A COMPUTATIONAL MODEL OF THE VESTIBULO-OCULAR REFLEX

Xavier Duran, supervised by Ivan Herreros and Paul Verschure

July 7, 2015

Master in Cognitive Systems and Interactive Media

VESTIBULO-OCULAR REFLEX (VOR)




1

This reflex functions to **stabilize images** on the retinas during **head movement** by producing **eye movements** in the direction opposite to head movement, thus preserving the image on the center of the visual field.

¹<http://bit.ly/10x3Qd6>

VOR ADAPTATION

VOR adaptation is **trained with light** and **measured in the dark**
[Boyden et al., 2004]



- **head movements** in the absence of **visual stimulation** cause a loss of the learned **eye movement response**
- changes in the amplitude, or gain of the VOR
- is mediated by an **active, extinction-like process** (not by passive forgetting)

[Cohen et al., 2004]

State of the art computational models of the vestibulo-ocular reflex don't define a physiological mechanism for extinction

NUCLEO-OLIVARY INHIBITION (NOI): A CANDIDATE SIGNAL

Inhibition of climbing fibres (retinal slip) serves as a teaching signal for extinction [Medina et al., 2002]




Figure 1: Neuroanatomical circuitry involved in eyeblink conditioning [Schonewille et al., 2010]

Would nucleo-olivary inhibition explain extinction in the vestibulo-ocular reflex computational models?

In direct evidence

- There is extinction of the adaptive response in the absence of peripheral error
- NOI has a role in the eye-blink reflex. The gain of the NOI is what determines the amplitude of the response on adaptive reflexes [Emken et al., 2007, Herreros and Verschure, 2013]

Adding nucleo-olivary inhibition on a vestibulo-ocular reflex computational model would explain extinction as in the experimental behavior of the reflex

METHODS

This computational model is made bottom-up from physiological and behavioral observations

- Plasticity on the cerebellar cortex
 - quick
 - short-term
 - error-based learning
- Plasticity on the brainstem
 - slow
 - long-term

[Clopath et al., 2014]

LEARNING BALANCE

- learned adaptation at the cerebellar cortex is slowly transferred to the brainstem
- cortical plasticity remains flexible to further adaptations
- savings help faster response on reacquisition




Figure 2: Short and long-term memory on eyeblink reflex adaptation [Yamazaki et al., 2015]


Extinction on Clopath's model

- Weakly modulated by head movement (vestibular signal)
- Weights on cortical plasticity decay linearly to their initial value

Extinction on NOI model

- Extinction is defined as proportional to cerebellar output
- More simple mechanism


VOR PHASE REVERSAL TRAINING PROTOCOL SIMULATION



[Gao et al., 2012]

- Day 1: VOR cancellation
- Day 2: VOR reversal with gain -0.5
- Day 3 and 4: Phase reversal with gain -1

VOR PHASE REVERSAL TRAINING PROTOCOL SIMULATION



[Gao et al., 2012]

- Day 1: VOR cancellation
- Day 2: VOR reversal with gain -0.5
- Day 3 and 4: Phase reversal with gain -1
- One week of light deprivation with vestibular stimulation

RESULTS

REPRODUCING CLOPATH'S RESULTS






Figure 3: Experimental results (left) and simulations (right)

1. Training session of gain 0.000000 (50 min)




Figure 4: Day 1: VOR cancellation training




Figure 5: Night 1: partial extinction

3. Training session of gain -0,500000 (50 min)




Figure 6: Day 2: VOR phase reversal training




Figure 7: Night 2: partial extinction

WHAT HAPPENS IN THE DARK?

- Cortical
 - Extinguishes progressively to a baseline
- Nuclear
 - Continues transference from cortical memory until it arrives at a nuclear balance
- Total
 - Extinction and transference go in opposite directions
 - On the dark adaptation continues consolidating until an inflexion point where extinction overtakes transference
 - After a long period on the dark, all cortical memory is consolidated on the brainstem and cortical contribution is at its baseline

CONCLUSIONS

- NOI **explains extinction** on VOR adaptation
- Extinction is triggered when vestibular information is available
- Teaching or error signal is modulated by **cerebellar output**
- Savings

CONCLUSIONS




Figure 8: GAD+IO neuron populations receive signals from cerebellar output [Uusisaari and Knöpfel, 2011]

More detailed bottom-up models

- transgenic mouse lines
- better electro-physiological recordings
- models with distributed plasticity

THANK YOU



Boyden, E. S., Katoh, A., and Raymond, J. L. (2004).

Cerebellum-dependent learning: the role of multiple plasticity mechanisms.

Annu. Rev. Neurosci., 27:581–609.



Clopath, C., Badura, A., De Zeeuw, C. I., and Brunel, N. (2014).

A cerebellar learning model of vestibulo-ocular reflex adaptation in wild-type and mutant mice.

J. Neurosci., 34(21):7203–15.



Cohen, M. R., Meissner, G. W., Schafer, R. J., and Raymond, J. L. (2004).

Reversal of motor learning in the vestibulo-ocular reflex in the absence of visual input.

Learn. Mem., 11(5):559–565.



Emken, J. L., Benitez, R., Sideris, A., Bobrow, J. E., and Reinkensmeyer, D. J. (2007).

Motor adaptation as a greedy optimization of error and effort.

J. Neurophysiol., 97(6):3997–4006.



Gao, Z., van Beugen, B. J., and De Zeeuw, C. I. (2012).

Distributed synergistic plasticity and cerebellar learning.

Nat. Rev. Neurosci., 13(9):619–35.



Herreros, I. and Verschure, P. F. M. J. (2013).

Nucleo-olivary inhibition balances the interaction between the reactive and adaptive layers in motor control.

Neural Netw., 47:64–71.



Medina, J. F., Nores, W. L., and Mauk, M. D. (2002).

Inhibition of climbing fibres is a signal for the extinction of conditioned eyelid responses.

Nature, 416(6878):330–333.



Schonewille, M., Belmeguenai, a., Koekkoek, S. K., Houtman, S. H., Boele, H. J., van Beugen, B. J., Gao, Z., Badura, a., Ohtsuki, G., Amerika, W. E., Hosy, E., Hoebeek, F. E., Elgersma, Y., Hansel, C., and De Zeeuw, C. I. (2010).
Purkinje cell-specific knockout of the protein phosphatase PP2B impairs potentiation and cerebellar motor learning.
Neuron, 67(4):618–28.



Uusisaari, M. and Knöpfel, T. (2011).
Functional classification of neurons in the mouse lateral Cerebellar Nuclei.
Cerebellum, 10(4):637–646.



Yamazaki, T., Nagao, S., Lennon, W., and Tanaka, S. (2015).
Modeling memory consolidation during posttraining periods in cerebellovestibular learning.
Proc. Natl. Acad. Sci., 112(11):201413798.