

UNIVERSITY OF LONDON

EC2020 ZA

BSc degrees and Diplomas for Graduates in Economics, Management, Finance and the Social Sciences, the Diplomas in Economics and Social Sciences

Elements of Econometrics

Thursday, 18 May 2017 : 14:30 to 17:30

Candidates should answer **EIGHT** of the following **TEN** questions: **ALL** of the questions in Section A (8 marks each) and **THREE** questions from Section B (20 marks each).

Candidates are strongly advised to divide their time accordingly.

Extracts from statistical tables are given after the final question on this paper.

Graph paper is provided at the end of this question paper. If used, it must be detached and fastened securely inside the answer book.

A calculator may be used when answering questions on this paper and it must comply in all respects with the specification given with your Admission Notice. The make and type of machine must be clearly stated on the front cover of the answer book.

If more questions are answered than requested, only the first answers attempted will be counted.

PLEASE TURN OVER

SECTION A

Answer all questions from this section

1. Discuss the drawbacks and advantages of using the Linear Probability Model when trying to explain a binary decision. In your answer clearly indicate what the Linear Probability Model is. **(8 marks)**
2. Discuss the consequences of measurement error.

In your answer consider the following empirical study attempting to estimate the relationship between advertising and magazine circulation rates. A simple linear relationship is postulated between A_t , the advertising rate in magazine t , and C_t , the circulation figure for the magazine in question:

$$A_t = \alpha + \beta C_t + \varepsilon_t, \quad t = 1, \dots, T,$$

The relation is estimated by least squares on data for $T = 75$ magazines. Unfortunately, there are considerable errors in measurement in the reported circulation figures. Critically discuss the following statement: 'The estimated coefficient $\hat{\beta}$ will be too small.' Rigour of your answer will be rewarded. **(8 marks)**

3. Consider the following regression model:

$$Y_i = \beta_0 + \beta_1 \frac{1}{X_i} + u_i, \quad i = 1, \dots, n.$$

We assume that the errors $\{u_i\}_{i=1}^n$ are independent normal random variables with zero mean and variance σ^2/X_i^2 . The regressor, $1/X_i$, is nonstochastic with positive sample variability and $X_i \neq 0$ for all i . You are interested in testing the hypothesis $H_0 : \beta_0 = 0$ against $H_1 : \beta_0 \neq 0$. You are advised to use the BLUE estimator of β_0 for this purpose.

Discuss how you would obtain the BLUE estimator of β_0 (note, you are not asked to derive this estimator).

Give two reasons why you would prefer using the BLUE estimator for β_0 instead of the OLS estimator $\hat{\beta}_{0,OLS} = \bar{Y} - \hat{\beta}_{1,OLS} \overline{(1/X)}$ when testing this hypothesis, where $\bar{Y} = \frac{1}{n} \sum Y_i$ and $\overline{(1/X)} = \frac{1}{n} \sum \frac{1}{X_i}$. **(8 marks)**

4. Consider the following non-stationary process:

$$y_t = \gamma_0 + \gamma_1 t + u_t, \text{ with } u_t = \varepsilon_t + \theta_1 \varepsilon_{t-1} + \theta_2 \varepsilon_{t-2}$$

and ε_t i.i.d. $(0, \sigma^2)$. Indicate (with explanation) the source(s) of non-stationarity of y_t . Discuss how you would test whether y_t indeed is non-stationary? Clearly indicate the null and the alternative hypothesis, the test statistic and the rejection rule. What name do we give such a non-stationary process? **(8 marks)**

5. Suppose you are given a random sample X_1, \dots, X_n from the exponential distribution:

$$f(x) = \lambda \exp(-\lambda x), \quad x > 0, \lambda > 0.$$

According to this distribution $E(X_i) = 1/\lambda$ and $Var(X_i) = 1/\lambda^2$ for $i = 1, \dots, n$.

Show that the maximum likelihood estimator for λ is $1/\bar{X}$ where $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i$. Is the estimator unbiased and/or consistent? Prove your claims. **(8 marks)**

SECTION B

Answer three questions from this section.

6. The following equation was estimated by Ordinary Least Squares based on second semester students in the fall term

$$\begin{aligned} \widehat{trmgpa} = & -2.12 + .900 crsgpa + .193 cumgpa + .0014 tothrs \\ & \begin{matrix} (.55) & (.175) & (.064) & (.0012) \\ [.55] & [.166] & [.074] & [.0012] \end{matrix} \\ & + .0018 sat - .0039 hspers + .351 female - .157 season \\ & \begin{matrix} (.0002) & (.0018) & (.085) & (.098) \\ [.0002] & [.0019] & [.079] & [.080] \end{matrix} \\ n = & 269; R^2 = .465, \end{aligned} \quad (6.1)$$

where $trmgpa_i$ is the term GPA (grade point average) of individual i . $crsgpa_i$ is a measure of difficulty of courses taken by individual i (weighted overall average of GPA in selected courses), $cumgpa_i$ is the GPA of individual i prior to the current semester, $tothrs_i$ is the total credit hours of individual i prior to the semester, sat_i is his/her SAT score (test taken for college admission in the USA), $hsperc_i$ is the graduating percentile of individual i in high school class, $female_i$ is a gender dummy, and $season_i$ is a dummy variable equal to unity if the student's sport is in season during the fall term. The usual standard errors are in parentheses and the White's heteroskedasticity-robust standard errors are in squared brackets.

- Explain the concept of heteroskedasticity and discuss the properties of the OLS estimator in the presence of heteroskedasticity. **(5 marks)**
- Do the variables $crsgpa$, $cumgpa$ and $tothrs$ have the expected estimated effects? Which of these variables are statistically significant at the 5% level? Does it matter which standard errors are used? **(5 marks)**
- In view of your concern about the presence of heteroskedasticity, provide the 95% confidence interval for β_{crsgpa} and use it to test the hypothesis $H_0 : \beta_{crsgpa} = 1$ against $H_A : \beta_{crsgpa} \neq 1$. Describe your conclusions. In view of your answer, indicate whether the p-value of the test is bigger or smaller than 5%. **(5 marks)**
- Discuss how you would conduct a test for heteroskedasticity in this setting. Clearly indicate the assumptions that underlie the test you suggest. Detail of your answer will be rewarded. **(5 marks)**

7. Consider the model:

$$y_t = \alpha_1 y_{t-1} + u_t, t = 1, \dots, T$$

where $y_0 = 0$. $E(u_t) = 0$, $E(u_t^2) = \sigma^2$ and $E(u_t u_s) = 0$ when $s \neq t$, for all $s, t = 1, \dots, T$

- (a) Discuss what we mean by the concept of stationarity (more precisely "covariance stationarity") and indicate under what condition $\{y_t\}_{t=1}^T$ will be stationary. **(3 marks)**
- (b) Discuss the Dickey-Fuller procedure used to test for the presence of a unit root in the above model. Clearly indicate the null and alternative hypothesis, test statistic and rejection rule. **(5 marks)**

(c) Consider a slight variation of the above model:

$$y_t = \alpha_1 y_{t-1} + \alpha_2 y_{t-2} + v_t, t = 1, \dots, T,$$

where $y_0 = 0$. $E(v_t) = 0$, $E(v_t^2) = \sigma^2$ and $E(v_t v_s) = 0$ when $s \neq t$, for all $s, t = 1, \dots, T$
What do we call such a process? Discuss the problem you will have when conducting your test as in (b). **(3 marks)**

- (d) Instead of conducting the Dickey-Fuller procedure, you are told to apply the Augmented DF test. Indicate how you would conduct the test for the presence of a unit root here. Derivation of the test equation will be required for full marks. **(5 marks)**
- (e) What are the potential problems associated with performing a regression with $I(1)$ variables? In your answer explain what it means for a variable to be $I(1)$. **(4 marks)**

8. For the US economy, let $gprice$ denote the monthly growth in the overall price level and let $gwage$ be the monthly growth in hourly wages. We have estimated the following distributed lag model (ADL(0,12)):

$$\begin{aligned}\widehat{gprice}_t = & -.00093 + .119gwage_t + .097gwage_{t-1} + .040gwage_{t-2} \\ & (.00057) \quad (.052) \quad (.039) \quad (.039) \\ & +.038gwage_{t-3} + .081gwage_{t-4} + .107gwage_{t-5} + .095gwage_{t-6} \\ & (.039) \quad (.039) \quad (.039) \quad (.039) \\ & +.104gwage_{t-7} + .103gwage_{t-8} + .159gwage_{t-9} + .110gwage_{t-10} \\ & (.039) \quad (.039) \quad (.039) \quad (.039) \\ & +.103gwage_{t-11} + .016gwage_{t-12}, \\ & (.039) \quad (.052) \\ n = & 273, R^2 = .317, \bar{R}^2 = .283, DW = .99.\end{aligned}$$

The usual standard errors are in parentheses.

- (a) What is the estimated long-run effect (LRP)? Is it very different from one? Explain what the LRP tells us in this example. How does this differ from the short-run effect? **(4 marks)**
- (b) We want to test whether the LRP is significantly smaller than one. Clearly indicating the null and the alternative hypothesis, give the test statistic and the rejection rule. What regression would you run to obtain the standard error of the LRP directly? **(6 marks)**
- (c) Your result in (b) may be affected by the presence of autocorrelation. Briefly explain this and discuss how you could use the Durbin-Watson test to detect whether this is indeed the case. Indicate clearly the assumptions underlying the Durbin Watson test, the test statistic and the rejection rule. **(5 marks)**
- Hint:* For the critical values of the DW test, use the largest sample size and number of regressors available in the table.
- (d) Assuming the Durbin-Watson test finds evidence of autocorrelation, discuss how you could resolve the problem by means of the iterated Cochrane-Orcutt procedure. For notational purposes, you may simplify your model to an ADL(0,1). **(5 marks)**

9. Let us consider the demand for fish. Using 97 daily price (*avgprc*) and quantity (*totqty*) observations on fish prices at the Fulton Fish Market in Manhattan, the following results were obtained by OLS:

$$\begin{aligned}\widehat{\log(\text{totqty}_t)} = & 8.244 - .425 \log(\text{avgprc}_t) - .311 \text{mon}_t - .683 \text{tues}_t \\ & \quad \quad \quad (.163) \quad (.176) \quad \quad \quad (.226) \quad \quad \quad (.223) \\ & - .533 \text{wed}_t + .067 \text{thurs}_t. \\ & \quad \quad \quad (.220) \quad \quad \quad (.220)\end{aligned}\quad (9.1)$$

The equation allows demand to differ across the days of the week, and Friday is the excluded dummy variable. The standard errors are in parentheses.

- (a) Interpret the coefficient of $\log(\text{avgprc})$ and discuss whether it is significant. **(3 marks)**
- (b) It is commonly thought that prices are jointly determined with quantity in equilibrium where demand equals supply. What are the consequences of this simultaneity for the properties of the OLS estimator? **(3 marks)**
- (c) The variables *wave2_t* and *wave3_t* are measures of ocean wave heights over the past several days. In view of your answer in (b), what two assumptions do we need to make in order to use *wave2_t* and *wave3_t* as instruments for $\log(\text{avgprc}_t)$ in estimating the demand equation? Discuss whether these assumptions are reasonable. **(4 marks)**
- (d) Below we report two sets of regression results, where the dependent variable is $\log(\text{avgprc}_t)$. Are *wave2_t* and *wave3_t* jointly significant? State the test statistic and rejection rule. How is your finding related to your answer in (c)? **(4 marks)**

Dependent Variable	Regressors				<i>R</i> ²	<i>RSS</i>	<i>n</i>
	<i>constant</i>	<i>wave2</i>	<i>wave3</i>	<i>day-of-the-week dummies</i>			
$\log(\text{avgprc}_t)$							
Regression (9.2)	−1.022 (.144)	.094 (.021)	.053 (.020)	<i>yes</i>	.3041	10.934	97
Regression (9.3)	−.276 (.092)	—	—	<i>yes</i>	.0088	15.576	97

- (e) The following IV results were obtained in Stata:

$$\begin{aligned}\widehat{\log(\text{totqty}_t)} = & 8.164 - .815 \log(\text{avgprc}_t) - .307 \text{mon}_t - .685 \text{tues}_t \\ & \quad \quad \quad (.182) \quad (.327) \quad \quad \quad (.229) \quad \quad \quad (.226) \\ & - .521 \text{wed}_t + .095 \text{thurs}_t. \\ & \quad \quad \quad (.223) \quad \quad \quad (.225)\end{aligned}\quad (9.4)$$

Discuss how these results can be obtained using Two Stage Least Squares (2SLS) and briefly discuss how you would test whether the results in (9.1) and (9.4) are significantly different from each other. **(6 marks)**

10. Let us consider an analysis on recidivism (probability of re-arrest) among a group of young men in California who have at least one arrest prior to 1986. The dependent variable, *arr86*, is equal to unity if the man was arrested at least once during 1986, and zero otherwise.

	OLS A	OLS B	Logit A	Logit B	Logit B Marginal Effect
<i>pcnv</i>	-.152 (.021)	-.162 (.021)	-.880 (.122)	-.901 (.120)	-.176 (.023)
<i>avgsen</i>	.005 (.006)	.006 (.006)	.027 (.035)	.031 (.034)	.006 (.007)
<i>totttime</i>	-.003 (.005)	-.002 (.005)	-.014 (.028)	-.010 (.027)	-.002 (.005)
<i>ptime86</i>	-.023 (.005)	-.022 (.005)	-.140 (.031)	-.127 (.031)	-.025 (.006)
<i>qemp86</i>	-.038 (.005)	-.043 (.005)	-.199 (.028)	-.216 (.028)	-.042 (.005)
<i>black</i>	.170 (.024)	—	.823 (.117)	—	—
<i>hispan</i>	.096 (.021)	—	.522 (.109)	—	—
<i>constant</i>	.380 (.019)	.380 (.019)	-.464 (.095)	-.169 (.084)	
R^2	.068	.047			
$\log L$			-1512.35	-1541.24	

pcnv is the proportion of prior arrests that led to a conviction, *avgsen* is the average sentence served from prior convictions, *totttime* is the months spent in prison since age 18 prior to 1986, *ptime86* is months spent in prison in 1986, *qemp86* is the number of quarters the man was legally employed in 1986, while *black* and *hispan* are two race dummies (*white* the excluded dummy). The standard errors are reported in parentheses.

- (a) When estimating the parameters by OLS, we are using the Linear Probability Model. Why might you then report heteroskedasticity-robust standard errors? **(2 marks)**
- (b) Using the OLS B results, what is the estimated effect on the probability of arrest if *pcnv* goes from 0.25 to 0.75 holding everything else constant? **(4 marks)**

(question continues on next page)

- (c) It is argued that the linear probability model is not appropriate for explaining the binary variable *arr86* and a logit regression model has been estimated. Explain how the Logit estimates are obtained. **(5 marks)**

Hint: You may recall, that for the Logit model A, we will specify

$$\Pr(arr86_i = 1) = \Lambda(\beta_0 + \beta_1 pcnv + \beta_2 avg\textit{sen} + \dots + \beta_6 black + \beta_7 hispan),$$

where $\Lambda(z) = \frac{1}{1 + \exp(-z)}$.


- (d) Using the Logit model results, discuss whether *black* and *hispan* are jointly significant. Clearly indicate the null and alternative hypothesis, the test statistic and the rejection rule. **(3 marks)**
- (e) An important distinction between the two approaches is that the marginal effect of *pcnv* on the probability of re-arrest is constant for the LPM unlike the marginal effect using the logit analysis. What this means for instance is that the estimated effect on the probability of arrest if *pcnv* goes from 0.25 to 0.75 will depend on the other characteristics.
- Explain how the marginal effects evaluated at the mean values of the explanatory variables (reported in the last column) were obtained. Give a brief comment as to how they compare to the marginal effect of the associated LPM. **(3 marks)**
 - Using the Logit B results, how would you obtain the estimated effect on the probability of arrest if *pcnv* goes from 0.25 to 0.75 for a *white* man, with characteristics *avg\textit{sen}*=1, *tot\textit{time}*=1, *p\textit{time}*86=0 and *qemp*86=2. A clear explanation of what calculations are required is sufficient. **(3 marks)**

END OF PAPER

TABLE 4. THE NORMAL DISTRIBUTION FUNCTION

The function tabulated is $\Phi(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^x e^{-\frac{1}{2}t^2} dt$. $\Phi(x)$ is

the probability that a random variable, normally distributed with zero mean and unit variance, will be less than or equal to x . When $x < 0$ use $\Phi(x) = 1 - \Phi(-x)$, as the normal distribution with zero mean and unit variance is symmetric about zero.



x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$
0.00	0.5000	0.40	0.6554	0.80	0.7881	1.20	0.8849	1.60	0.9452	2.00	0.97725
0.01	.5040	0.41	.6591	0.81	.7910	1.21	.8869	1.61	.9463	2.01	.97778
0.02	.5080	0.42	.6628	0.82	.7939	1.22	.8888	1.62	.9474	2.02	.97831
0.03	.5120	0.43	.6664	0.83	.7967	1.23	.8907	1.63	.9484	2.03	.97882
0.04	.5160	0.44	.6700	0.84	.7995	1.24	.8925	1.64	.9495	2.04	.97932
0.05	.5199	0.45	.6736	0.85	.8023	1.25	.8944	1.65	.9505	2.05	.97982
0.06	.5239	0.46	.6772	0.86	.8051	1.26	.8962	1.66	.9515	2.06	.98030
0.07	.5279	0.47	.6808	0.87	.8078	1.27	.8980	1.67	.9525	2.07	.98077
0.08	.5319	0.48	.6844	0.88	.8106	1.28	.8997	1.68	.9535	2.08	.98124
0.09	.5359	0.49	.6879	0.89	.8133	1.29	.9015	1.69	.9545	2.09	.98169
0.10	.5398	0.50	.6915	0.90	.8159	1.30	.9032	1.70	.9554	2.10	.98214
0.11	.5438	0.51	.6950	0.91	.8186	1.31	.9049	1.71	.9564	2.11	.98257
0.12	.5478	0.52	.6985	0.92	.8212	1.32	.9066	1.72	.9573	2.12	.98300
0.13	.5517	0.53	.7019	0.93	.8238	1.33	.9082	1.73	.9582	2.13	.98341
0.14	.5557	0.54	.7054	0.94	.8264	1.34	.9099	1.74	.9591	2.14	.98382
0.15	.5596	0.55	.7088	0.95	.8289	1.35	.9115	1.75	.9599	2.15	.98422
0.16	.5636	0.56	.7123	0.96	.8315	1.36	.9131	1.76	.9608	2.16	.98461
0.17	.5675	0.57	.7157	0.97	.8340	1.37	.9147	1.77	.9616	2.17	.98500
0.18	.5714	0.58	.7190	0.98	.8365	1.38	.9162	1.78	.9625	2.18	.98537
0.19	.5753	0.59	.7224	0.99	.8389	1.39	.9177	1.79	.9633	2.19	.98574
0.20	.5793	0.60	.7257	1.00	.8413	1.40	.9192	1.80	.9641	2.20	.98610
0.21	.5832	0.61	.7291	0.01	.8438	1.41	.9207	1.81	.9649	2.21	.98645
0.22	.5871	0.62	.7324	0.02	.8461	1.42	.9222	1.82	.9656	2.22	.98679
0.23	.5910	0.63	.7357	0.03	.8485	1.43	.9236	1.83	.9664	2.23	.98713
0.24	.5948	0.64	.7389	0.04	.8508	1.44	.9251	1.84	.9671	2.24	.98745
0.25	.5987	0.65	.7422	1.05	.8531	1.45	.9265	1.85	.9678	2.25	.98778
0.26	.6026	0.66	.7454	0.06	.8554	1.46	.9279	1.86	.9686	2.26	.98809
0.27	.6064	0.67	.7486	0.07	.8577	1.47	.9292	1.87	.9693	2.27	.98840
0.28	.6103	0.68	.7517	0.08	.8599	1.48	.9306	1.88	.9699	2.28	.98870
0.29	.6141	0.69	.7549	0.09	.8621	1.49	.9319	1.89	.9706	2.29	.98899
0.30	.6179	0.70	.7580	1.10	.8643	1.50	.9332	1.90	.9713	2.30	.98928
0.31	.6217	0.71	.7611	0.11	.8665	1.51	.9345	1.91	.9719	2.31	.98956
0.32	.6255	0.72	.7642	0.12	.8686	1.52	.9357	1.92	.9726	2.32	.98983
0.33	.6293	0.73	.7673	0.13	.8708	1.53	.9370	1.93	.9732	2.33	.99010
0.34	.6331	0.74	.7704	0.14	.8729	1.54	.9382	1.94	.9738	2.34	.99036
0.35	.6368	0.75	.7734	1.15	.8749	1.55	.9394	1.95	.9744	2.35	.99061
0.36	.6406	0.76	.7764	0.16	.8770	1.56	.9406	1.96	.9750	2.36	.99086
0.37	.6443	0.77	.7794	0.17	.8790	1.57	.9418	1.97	.9756	2.37	.99111
0.38	.6480	0.78	.7823	0.18	.8810	1.58	.9429	1.98	.9761	2.38	.99134
0.39	.6517	0.79	.7852	0.19	.8830	1.59	.9441	1.99	.9767	2.39	.99158
0.40	.6554	0.80	.7881	1.20	.8849	1.60	.9452	2.00	.9772	2.40	.99180

TABLE 4. THE NORMAL DISTRIBUTION FUNCTION

x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$
2.40	0.99180	2.55	0.99461	2.70	0.99653	2.85	0.99781	3.00	0.99865	3.15	0.99918
41	.99202	56	.99477	71	.99664	86	.99788	01	.99869	16	.99921
42	.99224	57	.99492	72	.99674	87	.99795	02	.99874	17	.99924
43	.99245	58	.99506	73	.99683	88	.99801	03	.99878	18	.99926
44	.99266	59	.99520	74	.99693	89	.99807	04	.99882	19	.99929
2.45	0.99286	2.60	0.99534	2.75	0.99702	2.90	0.99813	3.05	0.99886	3.20	0.99931
46	.99305	61	.99547	76	.99711	91	.99819	06	.99889	21	.99934
47	.99324	62	.99560	77	.99720	92	.99825	07	.99893	22	.99936
48	.99343	63	.99573	78	.99728	93	.99831	08	.99896	23	.99938
49	.99361	64	.99585	79	.99736	94	.99836	09	.99900	24	.99940
2.50	0.99379	2.65	0.99598	2.80	0.99744	2.95	0.99841	3.10	0.99903	3.25	0.99942
51	.99396	66	.99609	81	.99752	96	.99846	11	.99906	26	.99944
52	.99413	67	.99621	82	.99760	97	.99851	12	.99910	27	.99946
53	.99430	68	.99632	83	.99767	98	.99856	13	.99913	28	.99948
54	.99446	69	.99643	84	.99774	99	.99861	14	.99916	29	.99950
2.55	0.99461	2.70	0.99653	2.85	0.99781	3.00	0.99865	3.15	0.99918	3.30	0.99952

The critical table below gives on the left the range of values of x for which $\Phi(x)$ takes the value on the right, correct to the last figure given; in critical cases, take the upper of the two values of $\Phi(x)$ indicated.

3.075	0.9990	3.263	0.9994	3.731	0.99990	3.916	0.99995
3.105	0.9990	3.320	0.9995	3.759	0.99991	3.976	0.99996
3.138	0.9991	3.389	0.9996	3.791	0.99992	4.055	0.99997
3.174	0.9992	3.480	0.9997	3.826	0.99993	4.173	0.99998
3.215	0.9993	3.615	0.9998	3.867	0.99994	4.417	0.99999
	0.9994		0.9999		0.99995		1.00000


When $x > 3.3$ the formula $1 - \Phi(x) \doteq \frac{e^{-x^2}}{x\sqrt{2\pi}} \left[1 - \frac{1}{x^2} + \frac{3}{x^4} - \frac{15}{x^6} + \frac{105}{x^8} \right]$ is very accurate, with relative error less than $945/x^{10}$.

TABLE 5. PERCENTAGE POINTS OF THE NORMAL DISTRIBUTION

This table gives percentage points $x(P)$ defined by the equation

$$\frac{P}{100} = \frac{1}{\sqrt{2\pi}} \int_{x(P)}^{\infty} e^{-\frac{1}{2}t^2} dt.$$

If X is a variable, normally distributed with zero mean and unit variance, $P/100$ is the probability that $X \geq x(P)$. The lower P per cent points are given by symmetry as $-x(P)$, and the probability that $|X| \geq x(P)$ is $2P/100$.



P	$x(P)$	P	$x(P)$	P	$x(P)$	P	$x(P)$	P	$x(P)$	P	$x(P)$
50	0.0000	5.0	1.6449	3.0	1.8808	2.0	2.0537	1.0	2.3263	0.10	3.0902
45	0.1257	4.8	1.6646	2.9	1.8957	1.9	2.0749	0.9	2.3656	0.09	3.1214
40	0.2533	4.6	1.6849	2.8	1.9110	1.8	2.0969	0.8	2.4089	0.08	3.1559
35	0.3853	4.4	1.7060	2.7	1.9268	1.7	2.1201	0.7	2.4573	0.07	3.1947
30	0.5244	4.2	1.7279	2.6	1.9431	1.6	2.1444	0.6	2.5121	0.06	3.2389
25	0.6745	4.0	1.7507	2.5	1.9600	1.5	2.1701	0.5	2.5758	0.05	3.2905
20	0.8416	3.8	1.7744	2.4	1.9774	1.4	2.1973	0.4	2.6521	0.01	3.7190
15	1.0364	3.6	1.7991	2.3	1.9954	1.3	2.2262	0.3	2.7478	0.005	3.8906
10	1.2816	3.4	1.8250	2.2	2.0141	1.2	2.2571	0.2	2.8782	0.001	4.2649
5	1.6449	3.2	1.8522	2.1	2.0335	1.1	2.2904	0.1	3.0902	0.0005	4.4172


TABLE 8. PERCENTAGE POINTS OF THE χ^2 -DISTRIBUTION

This table gives percentage points $\chi^2_\nu(P)$ defined by the equation

$$\frac{P}{100} = \frac{1}{2^{\nu/2} \Gamma(\frac{\nu}{2})} \int_{\chi^2_\nu(P)}^{\infty} x^{\frac{\nu}{2}-1} e^{-\frac{x}{2}} dx.$$

If X is a variable distributed as χ^2 with ν degrees of freedom, $P/100$ is the probability that $X \geq \chi^2_\nu(P)$.

For $\nu > 100$, $\sqrt{2X}$ is approximately normally distributed with mean $\sqrt{2\nu-1}$ and unit variance.



(The above shape applies for $\nu \geq 3$ only. When $\nu < 3$ the mode is at the origin.)

P	99.95	99.9	99.5	99	97.5	95	90	80	70	60
$\nu = 1$	0.003927	0.001571	0.003927	0.001571	0.003927	0.003927	0.01579	0.06418	0.1485	0.2750
2	0.001000	0.002001	0.01003	0.02010	0.05064	0.1026	0.2107	0.4463	0.7133	1.022
3	0.01528	0.02430	0.07172	0.1148	0.2158	0.3518	0.5844	1.005	1.424	1.869
4	0.06392	0.09080	0.2070	0.2971	0.4844	0.7107	1.064	1.649	2.195	2.753
5	0.1581	0.2102	0.4117	0.5543	0.8312	1.145	1.610	2.343	3.000	3.655
6	0.2994	0.3811	0.6757	0.8721	1.237	1.635	2.204	3.070	3.828	4.570
7	0.4849	0.5985	0.9893	1.239	1.690	2.167	2.833	3.822	4.671	5.493
8	0.7104	0.8571	1.344	1.646	2.180	2.733	3.490	4.594	5.527	6.423
9	0.9717	1.152	1.735	2.088	2.700	3.325	4.168	5.380	6.393	7.357
10	1.265	1.479	2.156	2.558	3.247	3.940	4.865	6.179	7.267	8.295
11	1.587	1.834	2.603	3.053	3.816	4.575	5.578	6.989	8.148	9.237
12	1.934	2.214	3.074	3.571	4.404	5.226	6.304	7.807	9.034	10.18
13	2.305	2.617	3.565	4.107	5.009	5.892	7.042	8.634	9.926	11.13
14	2.697	3.041	4.075	4.660	5.629	6.571	7.790	9.467	10.82	12.08
15	3.108	3.483	4.601	5.229	6.262	7.261	8.547	10.31	11.72	13.03
16	3.536	3.942	5.142	5.812	6.908	7.962	9.312	11.15	12.62	13.98
17	3.980	4.416	5.697	6.408	7.564	8.672	10.09	12.00	13.53	14.94
18	4.439	4.905	6.265	7.015	8.231	9.390	10.86	12.86	14.44	15.89
19	4.912	5.407	6.844	7.633	8.907	10.12	11.65	13.72	15.35	16.85
20	5.398	5.921	7.434	8.260	9.591	10.85	12.44	14.58	16.27	17.81
21	5.896	6.447	8.034	8.897	10.28	11.59	13.24	15.44	17.18	18.77
22	6.404	6.983	8.643	9.542	10.98	12.34	14.04	16.31	18.10	19.73
23	6.924	7.529	9.260	10.20	11.69	13.09	14.85	17.19	19.02	20.69
24	7.453	8.085	9.886	10.86	12.40	13.85	15.66	18.06	19.94	21.65
25	7.991	8.649	10.52	11.52	13.12	14.61	16.47	18.94	20.87	22.62
26	8.538	9.222	11.16	12.20	13.84	15.38	17.29	19.82	21.79	23.58
27	9.093	9.803	11.81	12.88	14.57	16.15	18.11	20.70	22.72	24.54
28	9.656	10.39	12.46	13.56	15.31	16.93	18.94	21.59	23.65	25.51
29	10.23	10.99	13.12	14.26	16.05	17.71	19.77	22.48	24.58	26.48
30	10.80	11.59	13.79	14.95	16.79	18.49	20.60	23.36	25.51	27.44
32	11.98	12.81	15.13	16.36	18.29	20.07	22.27	25.15	27.37	29.38
34	13.18	14.06	16.50	17.79	19.81	21.66	23.95	26.94	29.24	31.31
36	14.40	15.32	17.89	19.23	21.34	23.27	25.64	28.73	31.12	33.25
38	15.64	16.61	19.29	20.69	22.88	24.88	27.34	30.54	32.99	35.19
40	16.91	17.92	20.71	22.16	24.43	26.51	29.05	32.34	34.87	37.13
50	23.46	24.67	27.99	29.71	32.36	34.76	37.69	41.45	44.31	46.86
60	30.34	31.74	35.53	37.48	40.48	43.19	46.46	50.64	53.81	56.62
70	37.47	39.04	43.28	45.44	48.76	51.74	55.33	59.90	63.35	66.40
80	44.79	46.52	51.17	53.54	57.15	60.39	64.28	69.21	72.92	76.19
90	52.28	54.16	59.20	61.75	65.65	69.13	73.29	78.56	82.51	85.99
100	59.90	61.92	67.33	70.06	74.22	77.93	82.36	87.95	92.13	95.81


TABLE 8. PERCENTAGE POINTS OF THE χ^2 -DISTRIBUTION

This table gives percentage points $\chi^2_\nu(P)$ defined by the equation

$$\frac{P}{100} = \frac{1}{2^{\nu/2} \Gamma(\frac{\nu}{2})} \int_{\chi^2_\nu(P)}^{\infty} x^{\frac{\nu}{2}-1} e^{-\frac{x}{2}} dx.$$

If X is a variable distributed as χ^2 with ν degrees of freedom, $P/100$ is the probability that $X \geq \chi^2_\nu(P)$.

For $\nu > 100$, $\sqrt{2X}$ is approximately normally distributed with mean $\sqrt{2\nu-1}$ and unit variance.



(The above shape applies for $\nu \geq 3$ only. When $\nu < 3$ the mode is at the origin.)


P	50	40	30	20	10	5	2.5	1	0.5	0.1	0.05
$\nu = 1$	0.4549	0.7083	1.074	1.642	2.706	3.841	5.024	6.635	7.879	10.83	12.12
2	1.386	1.833	2.408	3.219	4.605	5.991	7.378	9.210	10.60	13.82	15.20
3	2.366	2.946	3.665	4.642	6.251	7.815	9.348	11.34	12.84	16.27	17.73
4	3.357	4.045	4.878	5.989	7.779	9.488	11.14	13.28	14.86	18.47	20.00
5	4.351	5.132	6.064	7.289	9.236	11.07	12.83	15.09	16.75	20.52	22.11
6	5.348	6.211	7.231	8.558	10.64	12.59	14.45	16.81	18.55	22.46	24.10
7	6.346	7.283	8.383	9.803	12.02	14.07	16.01	18.48	20.28	24.32	26.02
8	7.344	8.351	9.524	11.03	13.36	15.51	17.53	20.09	21.95	26.12	27.87
9	8.343	9.414	10.66	12.24	14.68	16.92	19.02	21.67	23.59	27.88	29.67
10	9.342	10.47	11.78	13.44	15.99	18.31	20.48	23.21	25.19	29.59	31.42
11	10.34	11.53	12.90	14.63	17.28	19.68	21.92	24.72	26.76	31.26	33.14
12	11.34	12.58	14.01	15.81	18.55	21.03	23.34	26.22	28.30	32.91	34.82
13	12.34	13.64	15.12	16.98	19.81	22.36	24.74	27.69	29.82	34.53	36.48
14	13.34	14.69	16.22	18.15	21.06	23.68	26.12	29.14	31.32	36.12	38.11
15	14.34	15.73	17.32	19.31	22.31	25.00	27.49	30.58	32.80	37.70	39.72
16	15.34	16.78	18.42	20.47	23.54	26.30	28.85	32.00	34.27	39.25	41.31
17	16.34	17.82	19.51	21.61	24.77	27.59	30.19	33.41	35.72	40.79	42.88
18	17.34	18.87	20.60	22.76	25.99	28.87	31.53	34.81	37.16	42.31	44.43
19	18.34	19.91	21.69	23.90	27.20	30.14	32.85	36.19	38.58	43.82	45.97
20	19.34	20.95	22.77	25.04	28.41	31.41	34.17	37.57	40.00	45.31	47.50
21	20.34	21.99	23.86	26.17	29.62	32.67	35.48	38.93	41.40	46.80	49.01
22	21.34	23.03	24.94	27.30	30.81	33.92	36.78	40.29	42.80	48.27	50.51
23	22.34	24.07	26.02	28.43	32.01	35.17	38.08	41.64	44.18	49.73	52.00
24	23.34	25.11	27.10	29.55	33.20	36.42	39.36	42.98	45.56	51.18	53.48
25	24.34	26.14	28.17	30.68	34.38	37.65	40.65	44.31	46.93	52.62	54.95
26	25.34	27.18	29.25	31.79	35.56	38.89	41.92	45.64	48.29	54.05	56.41
27	26.34	28.21	30.32	32.91	36.74	40.11	43.19	46.96	49.64	55.48	57.86
28	27.34	29.25	31.39	34.03	37.92	41.34	44.46	48.28	50.99	56.89	59.30
29	28.34	30.28	32.46	35.14	39.09	42.56	45.72	49.59	52.34	58.30	60.73
30	29.34	31.32	33.53	36.25	40.26	43.77	46.98	50.89	53.67	59.70	62.16
32	31.34	33.38	35.66	38.47	42.58	46.19	49.48	53.49	56.33	62.49	65.00
34	33.34	35.44	37.80	40.68	44.90	48.60	51.97	56.06	58.96	65.25	67.80
36	35.34	37.50	39.92	42.88	47.21	51.00	54.44	58.62	61.58	67.99	70.59
38	37.34	39.56	42.05	45.08	49.51	53.38	56.90	61.16	64.18	70.70	73.35
40	39.34	41.62	44.16	47.27	51.81	55.76	59.34	63.69	66.77	73.40	76.09
50	49.33	51.89	54.72	58.16	63.17	67.50	71.42	76.15	79.49	86.66	89.56
60	59.33	62.13	65.23	68.97	74.40	79.08	83.30	88.38	91.95	99.61	102.7
70	69.33	72.36	75.69	79.71	85.53	90.53	95.02	100.4	104.2	112.3	115.6
80	79.33	82.57	86.12	90.41	96.58	101.9	106.6	112.3	116.3	124.8	128.3
90	89.33	92.76	96.52	101.1	107.6	113.1	118.1	124.1	128.3	137.2	140.8
100	99.33	102.9	106.9	111.7	118.5	124.3	129.6	135.8	140.2	149.4	153.2

TABLE 10. PERCENTAGE POINTS OF THE t -DISTRIBUTION

This table gives percentage points $t_\nu(P)$ defined by the equation

$$\frac{P}{100} = \frac{1}{\sqrt{\nu\pi}} \frac{\Gamma(\frac{1}{2}\nu + \frac{1}{2})}{\Gamma(\frac{1}{2}\nu)} \int_{t_\nu(P)}^{\infty} \frac{dt}{(1+t^2/\nu)^{\frac{1}{2}(\nu+1)}}.$$

Let X_1 and X_2 be independent random variables having a normal distribution with zero mean and unit variance and a χ^2 -distribution with ν degrees of freedom respectively; then $t = X_1/\sqrt{X_2/\nu}$ has Student's t -distribution with ν degrees of freedom, and the probability that $t \geq t_\nu(P)$ is $P/100$. The lower percentage points are given by symmetry as $-t_\nu(P)$, and the probability that $|t| \geq t_\nu(P)$ is $2P/100$.



The limiting distribution of t as ν tends to infinity is the normal distribution with zero mean and unit variance. When ν is large interpolation in ν should be harmonic.

P	40	30	25	20	15	10	5	2.5	1	0.5	0.1	0.05
$\nu = 1$	0.3249	0.7265	1.0000	1.3764	1.963	3.078	6.314	12.71	31.82	63.66	318.3	636.6
2	0.2887	0.6172	0.8165	1.0607	1.386	1.886	2.920	4.303	6.965	9.925	22.33	31.60
3	0.2767	0.5844	0.7649	0.9785	1.250	1.638	2.353	3.182	4.541	5.841	10.21	12.92
4	0.2707	0.5686	0.7407	0.9410	1.190	1.533	2.132	2.776	3.747	4.604	7.173	8.610
5	0.2672	0.5594	0.7267	0.9195	1.156	1.476	2.015	2.571	3.365	4.032	5.893	6.869
6	0.2648	0.5534	0.7176	0.9057	1.134	1.440	1.943	2.447	3.143	3.707	5.208	5.959
7	0.2632	0.5491	0.7111	0.8960	1.119	1.415	1.895	2.365	2.998	3.499	4.785	5.408
8	0.2619	0.5459	0.7064	0.8889	1.108	1.397	1.860	2.306	2.896	3.355	4.501	5.041
9	0.2610	0.5435	0.7027	0.8834	1.100	1.383	1.833	2.262	2.821	3.250	4.297	4.781
10	0.2602	0.5415	0.6998	0.8791	1.093	1.372	1.812	2.228	2.764	3.169	4.144	4.587
11	0.2596	0.5399	0.6974	0.8755	1.088	1.363	1.796	2.201	2.718	3.106	4.025	4.437
12	0.2590	0.5386	0.6955	0.8726	1.083	1.356	1.782	2.179	2.681	3.055	3.930	4.318
13	0.2586	0.5375	0.6938	0.8702	1.079	1.350	1.771	2.160	2.650	3.012	3.852	4.221
14	0.2582	0.5366	0.6924	0.8681	1.076	1.345	1.761	2.145	2.624	2.977	3.787	4.140
15	0.2579	0.5357	0.6912	0.8662	1.074	1.341	1.753	2.131	2.602	2.947	3.733	4.073
16	0.2576	0.5350	0.6901	0.8647	1.071	1.337	1.746	2.120	2.583	2.921	3.686	4.015
17	0.2573	0.5344	0.6892	0.8633	1.069	1.333	1.740	2.110	2.567	2.898	3.646	3.965
18	0.2571	0.5338	0.6884	0.8620	1.067	1.330	1.734	2.101	2.552	2.878	3.610	3.922
19	0.2569	0.5333	0.6876	0.8610	1.066	1.328	1.729	2.093	2.539	2.861	3.579	3.883
20	0.2567	0.5329	0.6870	0.8600	1.064	1.325	1.725	2.086	2.528	2.845	3.552	3.850
21	0.2566	0.5325	0.6864	0.8591	1.063	1.323	1.721	2.080	2.518	2.831	3.527	3.819
22	0.2564	0.5321	0.6858	0.8583	1.061	1.321	1.717	2.074	2.508	2.819	3.505	3.792
23	0.2563	0.5317	0.6853	0.8575	1.060	1.319	1.714	2.069	2.500	2.807	3.485	3.768
24	0.2562	0.5314	0.6848	0.8569	1.059	1.318	1.711	2.064	2.492	2.797	3.467	3.745
25	0.2561	0.5312	0.6844	0.8562	1.058	1.316	1.708	2.060	2.485	2.787	3.450	3.725
26	0.2560	0.5309	0.6840	0.8557	1.058	1.315	1.706	2.056	2.479	2.779	3.435	3.707
27	0.2559	0.5306	0.6837	0.8551	1.057	1.314	1.703	2.052	2.473	2.771	3.421	3.690
28	0.2558	0.5304	0.6834	0.8546	1.056	1.313	1.701	2.048	2.467	2.763	3.408	3.674
29	0.2557	0.5302	0.6830	0.8542	1.055	1.311	1.699	2.045	2.462	2.756	3.396	3.659
30	0.2556	0.5300	0.6828	0.8538	1.055	1.310	1.697	2.042	2.457	2.750	3.385	3.646
32	0.2555	0.5297	0.6822	0.8530	1.054	1.309	1.694	2.037	2.449	2.738	3.365	3.622
34	0.2553	0.5294	0.6818	0.8523	1.052	1.307	1.691	2.032	2.441	2.728	3.348	3.601
36	0.2552	0.5291	0.6814	0.8517	1.052	1.306	1.688	2.028	2.434	2.719	3.333	3.582
38	0.2551	0.5288	0.6810	0.8512	1.051	1.304	1.686	2.024	2.429	2.712	3.319	3.566
40	0.2550	0.5286	0.6807	0.8507	1.050	1.303	1.684	2.021	2.423	2.704	3.307	3.551
50	0.2547	0.5278	0.6794	0.8489	1.047	1.299	1.676	2.009	2.403	2.678	3.261	3.496
60	0.2545	0.5272	0.6786	0.8477	1.045	1.296	1.671	2.000	2.390	2.660	3.232	3.460
120	0.2539	0.5258	0.6765	0.8446	1.041	1.289	1.658	1.980	2.358	2.617	3.160	3.373
∞	0.2533	0.5244	0.6745	0.8416	1.036	1.282	1.645	1.960	2.326	2.576	3.090	3.291


TABLE 12(a). 10 PER CENT POINTS OF THE F-DISTRIBUTION

The function tabulated is $F(P) = F(P|\nu_1, \nu_2)$ defined by the equation

$$\frac{P}{100} = \frac{\Gamma(\frac{1}{2}\nu_1 + \frac{1}{2}\nu_2)}{\Gamma(\frac{1}{2}\nu_1) \Gamma(\frac{1}{2}\nu_2)} \nu_1^{\frac{1}{2}\nu_1} \nu_2^{\frac{1}{2}\nu_2} \int_{F(P)}^{\infty} \frac{F^{\frac{1}{2}\nu_1-1}}{(\nu_2 + \nu_1 F)^{\frac{1}{2}(\nu_1+\nu_2)}} dF,$$

for $P = 10, 5, 2.5, 1, 0.5$ and 0.1 . The lower percentage points, that is the values $F'(P) = F'(P|\nu_1, \nu_2)$ such that the probability that $F \leq F'(P)$ is equal to $P/100$, may be found by the formula

$$F'(P|\nu_1, \nu_2) = 1/F(P|\nu_2, \nu_1).$$




(This shape applies only when $\nu_1 \geq 3$. When $\nu_1 < 3$ the mode is at the origin.)

$\nu_1 =$	1	2	3	4	5	6	7	8	10	12	24	∞
$\nu_2 = 1$	39.86	49.50	53.59	55.83	57.24	58.20	58.91	59.44	60.19	60.71	62.00	63.33
2	8.526	9.000	9.162	9.243	9.293	9.326	9.349	9.367	9.392	9.408	9.450	9.491
3	5.538	5.462	5.391	5.343	5.309	5.285	5.266	5.252	5.230	5.216	5.176	5.134
4	4.545	4.325	4.191	4.107	4.051	4.010	3.979	3.955	3.920	3.896	3.831	3.761
5	4.060	3.780	3.619	3.520	3.453	3.405	3.368	3.339	3.297	3.268	3.191	3.105
6	3.776	3.463	3.289	3.181	3.108	3.055	3.014	2.983	2.937	2.905	2.818	2.722
7	3.589	3.257	3.074	2.961	2.883	2.827	2.785	2.752	2.703	2.668	2.575	2.471
8	3.458	3.113	2.924	2.806	2.726	2.668	2.624	2.589	2.538	2.502	2.404	2.293
9	3.360	3.006	2.813	2.693	2.611	2.551	2.505	2.469	2.416	2.379	2.277	2.159
10	3.285	2.924	2.728	2.605	2.522	2.461	2.414	2.377	2.323	2.284	2.178	2.055
11	3.225	2.860	2.660	2.536	2.451	2.389	2.342	2.304	2.248	2.209	2.100	1.972
12	3.177	2.807	2.606	2.480	2.394	2.331	2.283	2.245	2.188	2.147	2.036	1.904
13	3.136	2.763	2.560	2.434	2.347	2.283	2.234	2.195	2.138	2.097	1.983	1.846
14	3.102	2.726	2.522	2.395	2.307	2.243	2.193	2.154	2.095	2.054	1.938	1.797
15	3.073	2.695	2.490	2.361	2.273	2.208	2.158	2.119	2.059	2.017	1.899	1.755
16	3.048	2.668	2.462	2.333	2.244	2.178	2.128	2.088	2.028	1.985	1.866	1.718
17	3.026	2.645	2.437	2.308	2.218	2.152	2.102	2.061	2.001	1.958	1.836	1.686
18	3.007	2.624	2.416	2.286	2.196	2.130	2.079	2.038	1.977	1.933	1.810	1.657
19	2.990	2.606	2.397	2.266	2.176	2.109	2.058	2.017	1.956	1.912	1.787	1.631
20	2.975	2.589	2.380	2.249	2.158	2.091	2.040	1.999	1.937	1.892	1.767	1.607
21	2.961	2.575	2.365	2.233	2.142	2.075	2.023	1.982	1.920	1.875	1.748	1.586
22	2.949	2.561	2.351	2.219	2.128	2.060	2.008	1.967	1.904	1.859	1.731	1.567
23	2.937	2.549	2.339	2.207	2.115	2.047	1.995	1.953	1.890	1.845	1.716	1.549
24	2.927	2.538	2.327	2.195	2.103	2.035	1.983	1.941	1.877	1.832	1.702	1.533
25	2.918	2.528	2.317	2.184	2.092	2.024	1.971	1.929	1.866	1.820	1.689	1.518
26	2.909	2.519	2.307	2.174	2.082	2.014	1.961	1.919	1.855	1.809	1.677	1.504
27	2.901	2.511	2.299	2.165	2.073	2.005	1.952	1.909	1.845	1.799	1.666	1.491
28	2.894	2.503	2.291	2.157	2.064	1.996	1.943	1.900	1.836	1.790	1.656	1.478
29	2.887	2.495	2.283	2.149	2.057	1.988	1.935	1.892	1.827	1.781	1.647	1.467
30	2.881	2.489	2.276	2.142	2.049	1.980	1.927	1.884	1.819	1.773	1.638	1.456
32	2.869	2.477	2.263	2.129	2.036	1.967	1.913	1.870	1.805	1.758	1.622	1.437
34	2.859	2.466	2.252	2.118	2.024	1.955	1.901	1.858	1.793	1.745	1.608	1.419
36	2.850	2.456	2.243	2.108	2.014	1.945	1.891	1.847	1.781	1.734	1.595	1.404
38	2.842	2.448	2.234	2.099	2.005	1.935	1.881	1.838	1.772	1.724	1.584	1.390
40	2.835	2.440	2.226	2.091	1.997	1.927	1.873	1.829	1.763	1.715	1.574	1.377
60	2.791	2.393	2.177	2.041	1.946	1.875	1.819	1.775	1.707	1.657	1.511	1.291
120	2.748	2.347	2.130	1.992	1.896	1.824	1.767	1.722	1.652	1.601	1.447	1.193
∞	2.706	2.303	2.084	1.945	1.847	1.774	1.717	1.670	1.599	1.546	1.383	1.000

TABLE 12(b). 5 PER CENT POINTS OF THE F-DISTRIBUTION

If $F = \frac{X_1/\nu_1}{X_2/\nu_2}$, where X_1 and X_2 are independent random variables distributed as χ^2 with ν_1 and ν_2 degrees of freedom respectively, then the probabilities that $F \geq F(P)$ and that $F \leq F(P)$ are both equal to $P/100$. Linear interpolation in ν_1 and ν_2 will generally be sufficiently accurate except when either $\nu_1 > 12$ or $\nu_2 > 40$, when harmonic interpolation should be used.




(This shape applies only when $\nu_1 \geq 3$. When $\nu_1 < 3$ the mode is at the origin.)

$\nu_1 =$	1	2	3	4	5	6	7	8	10	12	24	∞
$\nu_2 = 1$	161.4	199.5	215.7	224.6	230.2	234.0	236.8	238.9	241.9	243.9	249.1	254.3
2	18.51	19.00	19.16	19.25	19.30	19.33	19.35	19.37	19.40	19.41	19.45	19.50
3	10.13	9.552	9.277	9.117	9.013	8.941	8.887	8.845	8.786	8.745	8.639	8.526
4	7.709	6.944	6.591	6.388	6.256	6.163	6.094	6.041	5.964	5.912	5.774	5.628
5	6.608	5.786	5.409	5.192	5.050	4.950	4.876	4.818	4.735	4.678	4.527	4.365
6	5.987	5.143	4.757	4.534	4.387	4.284	4.207	4.147	4.060	4.000	3.841	3.669
7	5.591	4.737	4.347	4.120	3.972	3.866	3.787	3.726	3.637	3.575	3.410	3.230
8	5.318	4.459	4.066	3.838	3.687	3.581	3.500	3.438	3.347	3.284	3.115	2.928
9	5.117	4.256	3.863	3.633	3.482	3.374	3.293	3.230	3.137	3.073	2.900	2.707
10	4.965	4.103	3.708	3.478	3.326	3.217	3.135	3.072	2.978	2.913	2.737	2.538
11	4.844	3.982	3.587	3.357	3.204	3.095	3.012	2.948	2.854	2.788	2.609	2.404
12	4.747	3.885	3.490	3.259	3.106	2.996	2.913	2.849	2.753	2.687	2.505	2.296
13	4.667	3.806	3.411	3.179	3.025	2.915	2.832	2.767	2.671	2.604	2.420	2.206
14	4.600	3.739	3.344	3.112	2.958	2.848	2.764	2.699	2.602	2.534	2.349	2.131
15	4.543	3.682	3.287	3.056	2.901	2.790	2.707	2.641	2.544	2.475	2.288	2.066
16	4.494	3.634	3.239	3.007	2.852	2.741	2.657	2.591	2.494	2.425	2.235	2.010
17	4.451	3.592	3.197	2.965	2.810	2.699	2.614	2.548	2.450	2.381	2.190	1.960
18	4.414	3.555	3.160	2.928	2.773	2.661	2.577	2.510	2.412	2.342	2.150	1.917
19	4.381	3.522	3.127	2.895	2.740	2.628	2.544	2.477	2.378	2.308	2.114	1.878
20	4.351	3.493	3.098	2.866	2.711	2.599	2.514	2.447	2.348	2.278	2.082	1.843
21	4.325	3.467	3.072	2.840	2.685	2.573	2.488	2.420	2.321	2.250	2.054	1.812
22	4.301	3.443	3.049	2.817	2.661	2.549	2.464	2.397	2.297	2.226	2.028	1.783
23	4.279	3.422	3.028	2.796	2.640	2.528	2.442	2.375	2.275	2.204	2.005	1.757
24	4.260	3.403	3.009	2.776	2.621	2.508	2.423	2.355	2.255	2.183	1.984	1.733
25	4.242	3.385	2.991	2.759	2.603	2.490	2.405	2.337	2.236	2.165	1.964	1.711
26	4.225	3.369	2.975	2.743	2.587	2.474	2.388	2.321	2.220	2.148	1.946	1.691
27	4.210	3.354	2.960	2.728	2.572	2.459	2.373	2.305	2.204	2.132	1.930	1.672
28	4.196	3.340	2.947	2.714	2.558	2.445	2.359	2.291	2.190	2.118	1.915	1.654
29	4.183	3.328	2.934	2.701	2.545	2.432	2.346	2.278	2.177	2.104	1.901	1.638
30	4.171	3.316	2.922	2.690	2.534	2.421	2.334	2.266	2.165	2.092	1.887	1.622
32	4.149	3.295	2.901	2.668	2.512	2.399	2.313	2.244	2.142	2.070	1.864	1.594
34	4.130	3.276	2.883	2.650	2.494	2.380	2.294	2.225	2.123	2.050	1.843	1.569
36	4.113	3.259	2.866	2.634	2.477	2.364	2.277	2.209	2.106	2.033	1.824	1.547
38	4.098	3.245	2.852	2.619	2.463	2.349	2.262	2.194	2.091	2.017	1.808	1.527
40	4.085	3.232	2.839	2.606	2.449	2.336	2.249	2.180	2.077	2.003	1.793	1.509
60	4.001	3.150	2.758	2.525	2.368	2.254	2.167	2.097	1.993	1.917	1.700	1.389
120	3.920	3.072	2.680	2.447	2.290	2.175	2.087	2.016	1.910	1.834	1.608	1.254
∞	3.841	2.996	2.605	2.372	2.214	2.099	2.010	1.938	1.831	1.752	1.517	1.000

TABLE 12(d). 1 PER CENT POINTS OF THE *F*-DISTRIBUTION

If $F = \frac{X_1}{\nu_1} / \frac{X_2}{\nu_2}$, where X_1 and X_2 are independent random variables distributed as χ^2 with ν_1 and ν_2 degrees of freedom respectively, then the probabilities that $F \geq F(P)$ and that $F \leq F(P)$ are both equal to $P/100$. Linear interpolation in ν_1 or ν_2 will generally be sufficiently accurate except when either $\nu_1 > 12$ or $\nu_2 > 40$, when harmonic interpolation should be used.



(This shape applies only when $\nu_1 \geq 3$. When $\nu_1 < 3$ the mode is at the origin.)

$\nu_1 =$	1	2	3	4	5	6	7	8	10	12	24	∞
$\nu_2 = 1$	4052	4999	5403	5625	5764	5859	5928	5981	6056	6106	6235	6366
2	98.50	99.00	99.17	99.25	99.30	99.33	99.36	99.37	99.40	99.42	99.46	99.50
3	34.12	30.82	29.46	28.71	28.24	27.91	27.67	27.49	27.23	27.05	26.60	26.13
4	21.20	18.00	16.69	15.98	15.52	15.21	14.98	14.80	14.55	14.37	13.93	13.46
5	16.26	13.27	12.06	11.39	10.97	10.67	10.46	10.29	10.05	9.888	9.466	9.020
6	13.75	10.92	9.780	9.148	8.746	8.466	8.260	8.102	7.874	7.718	7.313	6.880
7	12.25	9.547	8.451	7.847	7.460	7.191	6.993	6.840	6.620	6.469	6.074	5.650
8	11.26	8.649	7.591	7.006	6.632	6.371	6.178	6.029	5.814	5.667	5.279	4.859
9	10.56	8.022	6.992	6.422	6.057	5.802	5.613	5.467	5.257	5.111	4.729	4.311
10	10.04	7.559	6.552	5.994	5.636	5.386	5.200	5.057	4.849	4.706	4.327	3.909
11	9.646	7.206	6.217	5.668	5.316	5.069	4.886	4.744	4.539	4.397	4.021	3.602
12	9.330	6.927	5.953	5.412	5.064	4.821	4.640	4.499	4.296	4.155	3.780	3.361
13	9.074	6.701	5.739	5.205	4.862	4.620	4.441	4.302	4.100	3.960	3.587	3.165
14	8.862	6.515	5.564	5.035	4.695	4.456	4.278	4.140	3.939	3.800	3.427	3.004
15	8.683	6.359	5.417	4.893	4.556	4.318	4.142	4.004	3.805	3.666	3.294	2.868
16	8.531	6.226	5.292	4.773	4.437	4.202	4.026	3.890	3.691	3.553	3.181	2.753
17	8.400	6.112	5.185	4.669	4.336	4.102	3.927	3.791	3.593	3.455	3.084	2.653
18	8.285	6.013	5.092	4.579	4.248	4.015	3.841	3.705	3.508	3.371	2.999	2.566
19	8.185	5.926	5.010	4.500	4.171	3.939	3.765	3.631	3.434	3.297	2.925	2.489
20	8.096	5.849	4.938	4.431	4.103	3.871	3.699	3.564	3.368	3.231	2.859	2.421
21	8.017	5.780	4.874	4.369	4.042	3.812	3.640	3.506	3.310	3.173	2.801	2.360
22	7.945	5.719	4.817	4.313	3.988	3.758	3.587	3.453	3.258	3.121	2.749	2.305
23	7.881	5.664	4.765	4.264	3.939	3.710	3.539	3.406	3.211	3.074	2.702	2.256
24	7.823	5.614	4.718	4.218	3.895	3.667	3.496	3.363	3.168	3.032	2.659	2.211
25	7.770	5.568	4.675	4.177	3.855	3.627	3.457	3.324	3.129	2.993	2.620	2.169
26	7.721	5.526	4.637	4.140	3.818	3.591	3.421	3.288	3.094	2.958	2.585	2.131
27	7.677	5.488	4.601	4.106	3.785	3.558	3.388	3.256	3.062	2.926	2.552	2.097
28	7.636	5.453	4.568	4.074	3.754	3.528	3.358	3.226	3.032	2.896	2.522	2.064
29	7.598	5.420	4.538	4.045	3.725	3.499	3.330	3.198	3.005	2.868	2.495	2.034
30	7.562	5.390	4.510	4.018	3.699	3.473	3.304	3.173	2.979	2.843	2.469	2.006
32	7.499	5.336	4.459	3.969	3.652	3.427	3.258	3.127	2.934	2.798	2.423	1.956
34	7.444	5.289	4.416	3.927	3.611	3.386	3.218	3.087	2.894	2.758	2.383	1.911
36	7.396	5.248	4.377	3.890	3.574	3.351	3.183	3.052	2.859	2.723	2.347	1.872
38	7.353	5.211	4.343	3.858	3.542	3.319	3.152	3.021	2.828	2.692	2.316	1.837
40	7.314	5.179	4.313	3.828	3.514	3.291	3.124	2.993	2.801	2.665	2.288	1.805
60	7.077	4.977	4.126	3.649	3.339	3.119	2.953	2.823	2.632	2.496	2.115	1.601
120	6.851	4.787	3.949	3.480	3.174	2.956	2.792	2.663	2.472	2.336	1.950	1.381
∞	6.635	4.605	3.782	3.319	3.017	2.802	2.639	2.511	2.321	2.185	1.791	1.000

Durbin-Watson test statistic d : 1% significance points of d_L and d_U .

n	$k'=1$		$k'=2$		$k'=3$		$k'=4$		$k'=5$	
	d_L	d_U	d_L	d_U	d_L	d_U	d_L	d_U	d_L	d_U
15	0.81	1.07	0.70	1.25	0.59	1.46	0.49	1.70	0.39	1.96
16	0.84	1.09	0.74	1.25	0.63	1.44	0.53	1.66	0.44	1.90
17	0.87	1.10	0.77	1.25	0.67	1.43	0.57	1.63	0.48	1.85
18	0.90	1.12	0.80	1.26	0.71	1.42	0.61	1.60	0.52	1.80
19	0.93	1.13	0.83	1.26	0.74	1.41	0.65	1.58	0.56	1.77
20	0.95	1.15	0.86	1.27	0.77	1.41	0.68	1.57	0.60	1.74
21	0.97	1.16	0.89	1.27	0.80	1.41	0.72	1.55	0.63	1.71
22	1.00	1.17	0.91	1.28	0.83	1.40	0.75	1.54	0.66	1.69
23	1.02	1.19	0.94	1.29	0.86	1.40	0.77	1.53	0.70	1.67
24	1.04	1.20	0.96	1.30	0.88	1.41	0.80	1.53	0.72	1.66
25	1.05	1.21	0.98	1.30	0.90	1.41	0.83	1.52	0.75	1.65
26	1.07	1.22	1.00	1.31	0.93	1.41	0.85	1.52	0.78	1.64
27	1.09	1.23	1.02	1.32	0.95	1.41	0.88	1.51	0.81	1.63
28	1.10	1.24	1.04	1.32	0.97	1.41	0.90	1.51	0.83	1.62
29	1.12	1.25	1.05	1.33	0.99	1.42	0.92	1.51	0.85	1.61
30	1.13	1.26	1.07	1.34	1.01	1.42	0.94	1.51	0.88	1.61
31	1.15	1.27	1.08	1.34	1.02	1.42	0.96	1.51	0.90	1.60
32	1.16	1.28	1.10	1.35	1.04	1.43	0.98	1.51	0.92	1.60
33	1.17	1.29	1.11	1.36	1.05	1.43	1.00	1.51	0.94	1.59
34	1.18	1.30	1.13	1.36	1.07	1.43	1.01	1.51	0.95	1.59
35	1.19	1.31	1.14	1.37	1.08	1.44	1.03	1.51	0.97	1.59
36	1.21	1.32	1.15	1.38	1.10	1.44	1.04	1.51	0.99	1.59
37	1.22	1.32	1.16	1.38	1.11	1.45	1.06	1.51	1.00	1.59
38	1.23	1.33	1.18	1.39	1.12	1.45	1.07	1.52	1.02	1.58
39	1.24	1.34	1.19	1.39	1.14	1.45	1.09	1.52	1.03	1.58
40	1.25	1.34	1.20	1.40	1.15	1.46	1.10	1.52	1.05	1.58
45	1.29	1.38	1.24	1.42	1.20	1.48	1.16	1.53	1.11	1.58
50	1.32	1.40	1.28	1.45	1.24	1.49	1.20	1.54	1.16	1.59
55	1.36	1.43	1.32	1.47	1.28	1.51	1.25	1.55	1.21	1.59
60	1.38	1.45	1.35	1.48	1.32	1.52	1.28	1.56	1.25	1.60
65	1.41	1.47	1.38	1.50	1.35	1.53	1.31	1.57	1.28	1.61
70	1.43	1.49	1.40	1.52	1.37	1.55	1.34	1.58	1.31	1.61
75	1.45	1.50	1.42	1.53	1.39	1.56	1.37	1.59	1.34	1.62
80	1.47	1.52	1.44	1.54	1.42	1.57	1.39	1.60	1.36	1.62
85	1.48	1.53	1.46	1.55	1.43	1.58	1.41	1.60	1.39	1.63
90	1.50	1.54	1.47	1.56	1.45	1.59	1.43	1.61	1.41	1.64
95	1.51	1.55	1.49	1.57	1.47	1.60	1.45	1.62	1.42	1.64
100	1.52	1.56	1.50	1.58	1.48	1.60	1.46	1.63	1.44	1.65

n = number of observations

k' = number of explanatory variables

Durbin-Watson test statistic d : 5% significance points of d_L and d_U .

n	$k'=1$		$k'=2$		$k'=3$		$k'=4$		$k'=5$	
	d_L	d_U	d_L	d_U	d_L	d_U	d_L	d_U	d_L	d_U
15	1.08	1.36	0.95	1.54	0.82	1.75	0.69	1.97	0.56	2.21
16	1.10	1.37	0.98	1.54	0.86	1.73	0.74	1.93	0.62	2.15
17	1.13	1.38	1.02	1.54	0.90	1.71	0.78	1.90	0.67	2.10
18	1.16	1.39	1.05	1.53	0.93	1.69	0.82	1.87	0.71	2.06
19	1.18	1.40	1.08	1.53	0.97	1.68	0.86	1.85	0.75	2.02
20	1.20	1.41	1.10	1.54	1.00	1.68	0.90	1.83	0.79	1.99
21	1.22	1.42	1.13	1.54	1.03	1.67	0.93	1.81	0.83	1.96
22	1.24	1.43	1.15	1.54	1.05	1.66	0.96	1.80	0.86	1.94
23	1.26	1.44	1.17	1.54	1.08	1.66	0.99	1.79	0.90	1.92
24	1.27	1.45	1.19	1.55	1.10	1.66	1.01	1.78	0.93	1.90
25	1.29	1.45	1.21	1.55	1.12	1.66	1.04	1.77	0.95	1.89
26	1.30	1.46	1.22	1.55	1.14	1.65	1.06	1.76	0.98	1.88
27	1.32	1.47	1.24	1.56	1.16	1.65	1.08	1.76	1.01	1.86
28	1.33	1.48	1.26	1.56	1.18	1.65	1.10	1.75	1.03	1.85
29	1.34	1.48	1.27	1.56	1.20	1.65	1.12	1.74	1.05	1.84
30	1.35	1.49	1.28	1.57	1.21	1.65	1.14	1.74	1.07	1.83
31	1.36	1.50	1.30	1.57	1.23	1.65	1.16	1.74	1.09	1.83
32	1.37	1.50	1.31	1.57	1.24	1.65	1.18	1.73	1.11	1.82
33	1.38	1.51	1.32	1.58	1.26	1.65	1.19	1.73	1.13	1.81
34	1.39	1.51	1.33	1.58	1.27	1.65	1.21	1.73	1.15	1.81
35	1.40	1.52	1.34	1.58	1.28	1.65	1.22	1.73	1.16	1.80
36	1.41	1.52	1.35	1.59	1.29	1.65	1.24	1.73	1.18	1.80
37	1.42	1.53	1.36	1.59	1.31	1.66	1.25	1.72	1.19	1.80
38	1.43	1.54	1.37	1.59	1.32	1.66	1.26	1.72	1.21	1.79
39	1.43	1.54	1.38	1.60	1.33	1.66	1.27	1.72	1.22	1.79
40	1.44	1.54	1.39	1.60	1.34	1.66	1.29	1.72	1.23	1.79
45	1.48	1.57	1.43	1.62	1.38	1.67	1.34	1.72	1.29	1.78
50	1.50	1.59	1.46	1.63	1.42	1.67	1.38	1.72	1.34	1.77
55	1.53	1.60	1.49	1.64	1.45	1.68	1.41	1.72	1.38	1.77
60	1.55	1.62	1.51	1.65	1.48	1.69	1.44	1.73	1.41	1.77
65	1.57	1.63	1.54	1.66	1.50	1.70	1.47	1.73	1.44	1.77
70	1.58	1.64	1.55	1.67	1.52	1.70	1.49	1.74	1.46	1.77
75	1.60	1.65	1.57	1.68	1.54	1.71	1.51	1.74	1.49	1.77
80	1.61	1.66	1.59	1.69	1.56	1.72	1.53	1.74	1.51	1.77
85	1.62	1.67	1.60	1.70	1.57	1.72	1.55	1.75	1.52	1.77
90	1.63	1.68	1.61	1.70	1.59	1.73	1.57	1.75	1.54	1.78
95	1.64	1.69	1.62	1.71	1.60	1.73	1.58	1.75	1.56	1.78
100	1.65	1.69	1.63	1.72	1.61	1.74	1.59	1.76	1.57	1.78

n = number of observations

k' = number of explanatory variables