

# Ornstein - Uhlenbeck processes. Novikov martingale method.

Kasianova, Shulyak

Higher School of Economics

May 18, 2020

# Some results on AR(1)

OU  
processes

We shall assume that we have some underlying probability space  $(\Omega, \mathcal{F}, P)$  in our discussions below and  $\mathbf{E}$  will be expectations with respect to this probability.

Some  
results on  
AR(1)

Ornstein -  
Uhlenbeck  
processes

We shall study some aspects of the simplest autoregressive time series : AR(1). It is given as a solution of

Stationary  
Solutions

$$X_t = \alpha X_{t-1} + \varepsilon_t$$

Remarks

Crossing-  
time  
problem

for  $t \in T$  where  $T = \{0, 1, 2, 3, \dots\}$  or  $T = \{0, \pm 1, \pm 2, \pm 3, \dots\}$ .

Integral  
Equation  
Methods

We are mainly interested in the following problem: given that  $X_0 = x > 0$  and  $\tau(x) = \min\{n \geq 0 : X_n \leq 0\}$ , what is the distribution of  $\tau$ ? In other words we see values for  $P\{\tau(x) = n\}$  for each integer  $n \geq 0$ . The difficulty with this problem is the fact that we do not have  $X_{\tau(x)} = 0$ .

Martingales  
and first  
passage  
times of  
AR(1)  
sequences

We shall assume that the process  $\{X_t\}$  is real valued, but there are generalizations of this in the literature.

Vervaat's  
Theorem

Remark

Contribu-  
tions by  
Novikov

## Definition 1

Let  $\{X_t : t \in \mathbb{T}\}$  be a process with  $V(X_t) < \infty$ . Then the **autocorrelation**  $\gamma$  of it is defined by

$$\gamma(r, s) = \text{cov}(X_r, X_s) = \mathbf{E}[(X_r - \mathbf{E}[X_r])(X_s - \mathbf{E}[X_s])]$$

for  $r, s \in \mathbb{T}$ .

# Some results on AR(1)

OU  
processes

## Definition 2

Let  $T = \{0, \pm 1, \pm 2, \pm 3, \dots\}$ . We say that  $\{X_t\}$  is stationary if

- (i)  $\mathbf{E}[|X_t|^2] < \infty$  for all  $t \in T$
- (ii)  $\mathbf{E}[X_t] = m$  for all  $t \in T$  and some real  $m$
- (iii)  $\gamma(r, s) = \gamma(r + t, s + t)$  for all  $s, t, r \in T$

We note that (iii) implies  $\gamma(r, s) = \gamma(r - s, 0)$  if we put  $t = -s$ .

We then write  $\gamma(h)$  for  $\gamma(h, 0)$ .

Alternatively, such processes are also called weak stationary, covariance stationary, stationary in wide sense, and second order stationary.

## Definition 3

We say that  $\{X_t : t \in T\}$  is strictly stationary if

$$\{X_{t_1}, X_{t_2}, \dots, X_{t_n}\} \stackrel{d}{=} \{X_{t_1+h}, X_{t_2+h}, \dots, X_{t_n+h}\}$$

for all positive integers  $n$  and  $t_1, t_2, \dots, t_n, h \in T$ .

Some  
results on  
AR(1)

Ornstein -  
Uhlenbeck  
processes

Stationary  
Solutions

Remarks

Crossing-  
time  
problem

Integral  
Equation  
Methods

Martingales  
and first  
passage  
times of  
AR(1)  
sequences

Vervaat's  
Theorem

Remark

Contribu-  
tions by  
Novikov

# Some results on AR(1)

OU  
processes

In the case of AR(1) we have

$$X_t = \alpha X_{t-1} + \varepsilon_t \quad (1)$$

with  $\mathbf{E}[\varepsilon_t] = 0$  and  $\mathbf{E}[\varepsilon_t^2] = \sigma^2$  for all  $t$ , and the correlation

$$\gamma(h) = \text{cov}(\varepsilon_t, \varepsilon_{t+h}) = \begin{cases} 0 & \text{if } h \neq 0 \\ \sigma^2 & \text{if } h = 0 \end{cases}$$

If this is the case, the noises are uncorrelated and  $\{\varepsilon_t\}$  is said to be  $\text{WN}(0, \sigma^2)$ , a white noise process.

If  $\{\varepsilon_t\}$  are iid, then  $\{\varepsilon_t\}$  is said to be an  $\text{IID}(0, \sigma^2)$  process.

In the classic case we have  $\{\varepsilon_t\} \sim \mathcal{N}(0, \sigma^2)$  and uncorrelated or we have  $\{\varepsilon_t\} \sim \mathcal{N}(0, \sigma^2)$  and independent.

In the case that  $\{\varepsilon_t\}$  are iid and  $\mathbf{E}[\varepsilon_t^2] < \infty$ , then it turns out that the two types of stationarity are the same.

Time series people are interested in **stationary solutions** of equation (1), and estimation of such time series are based on this assumption.

Some  
results on  
AR(1)

Ornstein -  
Uhlenbeck  
processes

Stationary  
Solutions

Remarks

Crossing-  
time  
problem

Integral  
Equation  
Methods

Martingales  
and first  
passage  
times of  
AR(1)  
sequences

Vervaat's  
Theorem

Remark

Contributions  
by  
Novikov

# Ornstein - Uhlenbeck processes

## Recall

- Arithmetic BM:  $dX_t = \mu dt + \sigma dB_t$ .
- Geometric BM:  $dX_t = \mu X_t dt + \sigma X_t dB_t$ .
- OU process:  $dX_t = \theta(\mu - X_t)dt + \sigma dB_t$ .

OU process is a mean-reverting process, i.e. over time, the process tends to drift towards its mean function.

Direction and the magnitude of the drift is not constant but changes depending on the difference between the value of the process and its long term mean.

When  $X_t$  is below  $\mu$  the positive drift pulls it back up towards  $\mu$ , whereas the opposite occurs when  $X_t$  is above  $\mu$ . Therefore,  $\mu$  may be interpreted as the “long run mean” of the process ( $E[X_t] \rightarrow \mu$  as  $t \rightarrow \infty$ ) and the mean-reversion coefficient  $\theta$  is the speed of adjustment of  $X_t$  towards its long run. Stochastic term  $\sigma dB_t$  captures the instantaneous volatility, i.e. constant instantaneous variance  $\sigma^2$ , which ensures that the process erratically and continuously fluctuates around the level  $\mu$ .

# Ornstein - Uhlenbeck processes

OU  
processes

$$dX_t = \theta(\mu - X_t)dt + \sigma dB_t \quad (1)$$

Figure 1 shows three sample paths of equation (1) for  $\sigma = 0.05$ ,  $\mu = 1$ , and for increasing values of  $\theta$ . It can be seen that the smaller  $\theta$ , the more  $X_t$  drifts away from  $\mu$ . For  $\theta = 2$ ,  $X_t$  displays only small and short-lived deviations from  $\mu$ .




Figure 1: Three sample paths of the mean-reverting Ornstein-Uhlenbeck process.

To solve SDE

$$dX_t = \theta(\mu - X_t)dt + \sigma dB_t \quad (2)$$

let's first consider the deterministic version of this equation

$$dX_t = \theta(\mu - X_t)dt \quad (3)$$

$$dX_t + \theta X_t dt = \theta \mu dt \quad (4)$$

$$\frac{dX_t}{dt} + \theta X_t = \theta \mu \quad (5)$$

Can be solved via the integrating factor method.

The integrating factor is  $e^{\int_0^t \theta ds} = e^{\theta t}$

$$e^{\theta t} \frac{dX_t}{dt} + e^{\theta t} \theta X_t = e^{\theta t} \theta \mu \quad (6)$$

The left hand side is an exact differential of the product.

$$\frac{d}{dt}(e^{\theta t} X_t) = e^{\theta t} \theta \mu \quad (7)$$



Recall, that for OU process by Ito's lemma  $dX_t^2 = \sigma^2 dt$

$$dX_t = \theta(\mu - X_t)dt + \sigma dB_t \quad (8)$$

We are looking for solution in a form

$$d(e^{\theta t} X_t) = e^{\theta t} \theta \mu dt \quad (9)$$

$$dX_t + \theta X_t dt = \theta \mu dt + \sigma dB_t \quad (10)$$

Using the same integrating factor

$$e^{\theta t} dX_t + \theta e^{\theta t} X_t dt = \theta \mu e^{\theta t} dt + \sigma e^{\theta t} dB_t \quad (11)$$

By Ito's product rule  $d(e^{\theta t} X_t) = d(e^{\theta t})X_t + e^{\theta t} dX_t + d(e^{\theta t})dX_t$ ,  $d(e^{\theta t})dX_t = 0$  as a covariance between a deterministic and stochastic function is zero

$$d(e^{\theta t} X_t) = \theta \mu e^{\theta t} dt + \sigma e^{\theta t} dB_t \quad (12)$$

$$\int_0^T d(e^{\theta t} X_t) = \int_0^T \theta \mu e^{\theta t} dt + \int_0^T \sigma e^{\theta t} dB_t \quad (13)$$

$$e^{\theta T} X_T - e^0 X_0 = \theta \mu \frac{e^{\theta T} - e^0}{\theta} + \int_0^T \sigma e^{\theta t} dB_t \quad (14)$$

$$X_T = e^{-\theta T} X_0 + \mu(1 - e^{-\theta T}) + \sigma e^{-\theta T} \int_0^T e^{\theta t} dB_t \quad (15)$$

SDE:

$$dX_t = \theta(\mu - X_t)dt + \sigma dB_t \quad (16)$$

Solution:

$$X_T = e^{-\theta T}X_0 + \mu(1 - e^{-\theta T}) + \sigma \int_0^T e^{-\theta(T-t)}dB_t \quad (17)$$

We now characterize its probability distribution. It is easy to see that  $X$  is normally distributed, as the integral of a deterministic function w.r.t. Brownian is Gaussian.

The expected value of a deterministic function w.r.t. Brownian is zero. Therefore,

$$E[X_T] = E[e^{-\theta T}X_0 + \mu(1 - e^{-\theta T}) + \sigma \int_0^T e^{-\theta(T-t)}dB_t] = e^{-\theta T}X_0 + \mu(1 - e^{-\theta T}) \quad (18)$$

Using Ito's isometry rule

$$E[(\int_0^T X_t dW_t)^2] = E[\int_0^T X_t^2 dt] \quad (19)$$

gives us a deterministic integral that we can easily evaluate

$$Var[X_T] = E[(X_T - E[X_T])^2] = E[(\sigma \int_0^T e^{-\theta(T-t)}dB_t)^2] = \sigma^2 E[\int_0^T e^{-2\theta(T-t)}dt] = \sigma^2 \frac{1 - e^{-2\theta T}}{2\theta} \quad (20)$$

Up to this point we have assumed the start point to be 0 and the end point to be time T. The distribution of the value of the process at time T given the information at time 0.

$$X_T \sim N(e^{-\theta T} X_0 + \mu(1 - e^{-\theta T}), \sigma^2 \frac{1 - e^{-2\theta T}}{2\theta}) \quad (21)$$

We can generalize this to an arbitrary starting point  $X_t$

$$X_{t+\Delta t} = e^{-\theta \Delta t} X_t + \mu(1 - e^{-\theta \Delta t}) + \sigma \sqrt{\frac{1 - e^{-2\theta \Delta t}}{2\theta}} N(0, 1) \quad (22)$$

It is easy to see that the process is similar to AR(1)

$$y_{t+1} = \beta y_t + \alpha + e_{t+1} \quad (23)$$

# Ornstein - Uhlenbeck processes

OU  
processes

In this section  $\{X_t\}$  is an Ornstein-Uhlenbeck process.

Some  
results on  
AR(1)

Suppose that  $X_0 = x$  and

$$\tau(x) = \inf \{t \geq 0 : X_t \leq 0\}$$

Ornstein -  
Uhlenbeck  
processes

then we wish to find  $P(\tau(x) \leq t)$ .

Stationary  
Solutions

Clearly we need only study the case where  $x > 0$ .

Remarks

Crossing-  
time  
problem

Integral  
Equation  
Methods

Martingales  
and first  
passage  
times of  
AR(1)  
sequences

Vervaat's  
Theorem

Remark

Contribu-  
tions by  
Novikov

# Ornstein - Uhlenbeck processes

Yi, Chuang. "On the first passage time distribution of an Ornstein–Uhlenbeck process." Quantitative Finance 10, no. 9 (2010): 957-960.

Let

$$dX_t = -\alpha X_t dt + \sigma \sqrt{2\alpha} dB_t \quad (24)$$

and let  $X_0 = x > 0$ .

Then for each  $t > 0$  the solution  $X_t$  is Gaussian and

$$X_t \sim \mathcal{N}(x e^{-\alpha t}, \sigma^2(1 - e^{-2\alpha t}))$$

In fact

$$d(e^{\alpha t} X_t) = e^{\alpha t} (\alpha X_t dt + dX_t) = e^{\alpha t} \sigma \sqrt{2\alpha} dB_t$$

implies that

$$e^{\alpha t} X_t = X_0 + \int_0^t e^{\alpha s} \sigma \sqrt{2\alpha} dB_s$$

and hence

$$X_t = e^{-\alpha t} x + \sigma \sqrt{2\alpha} \int_0^t e^{-\alpha(t-s)} dB_s$$

Thus

$$X_t \sim \mathcal{N}(m(t), n(t)^2)$$

where

$$m(t) = \mathbf{E}[X(t)] = e^{-\alpha t} x$$

and

$$n(t)^2 = \mathbf{E} \left[ \left( \sigma \sqrt{2\alpha} \int_0^t e^{-\alpha(t-s)} dB_s \right)^2 \right] = \sigma^2 2\alpha \int_0^t e^{-2\alpha(t-s)} ds = \sigma^2 (1 - e^{-2\alpha t})$$

## Theorem 1

We have

$$\mathbf{P}(\tau(x) \leq t) = 2 \mathbf{P}(X_t \leq 0)$$

## Corollary

From the theorem

$$\begin{aligned} P(\tau(x) \leq t) &= 2 P\left(\frac{X_t - m(t)}{n(t)} \leq -\frac{m(t)}{n(t)}\right) \\ &= 2 \Phi\left(-\frac{m(t)}{n(t)}\right) \\ &= 2 \Phi\left(\frac{-xe^{-\alpha t}}{\sigma\sqrt{1 - e^{-2\alpha t}}}\right) \\ &= \int_0^t f(u) du \end{aligned}$$

where  $\Phi$  stands for cumulative distribution function on  $\mathcal{N}(0, 1)$  and

$$f(u) = \frac{x}{2\sigma\sqrt{\pi\alpha}} \exp\left\{-\frac{x^2 e^{-\alpha u}}{4\sigma^2 \sinh(\alpha u)} + \frac{\alpha u}{2}\right\} \left(\frac{\alpha}{\sinh(\alpha u)}\right)^{\frac{3}{2}}$$

for  $u > 0$  will be a consequence.

## Proof of the Theorem

We note at

$$\begin{aligned}d(-X_t) &= -\lambda(-X_t) dt - \sigma dB_t \\ &= -\lambda(-X_t) dt + \sigma d\tilde{B}_t\end{aligned}$$

where  $\tilde{B}_t = -B_t$ .

Define a new process by

$$\tilde{X}_t = \begin{cases} X_t & \text{if } t \leq \tau \\ -X_t & \text{if } t > \tau \end{cases}$$

then  $\tilde{X}_0 = X_0 = x$ .

We now claim that  $\tilde{X}_t \stackrel{d}{=} X_t$ .



# Ornstein - Uhlenbeck processes

OU  
processes

To show this, let  $g$  be an arbitrary bounded Borel function. Then

$$\begin{aligned}\mathbf{E}[g(\tilde{X}_t)] &= \mathbf{E}[g(\tilde{X}_t)\mathbf{I}(t \leq \tau)] + \mathbf{E}[g(\tilde{X}_t)\mathbf{I}(t > \tau)] \\ &= \mathbf{E}[g(X_t)\mathbf{I}(t \leq \tau)] + \mathbf{E}[g(-X_t)\mathbf{I}(t > \tau)] \\ &= \mathbf{E}[g(X_t)] + \mathbf{E}[g(-X_t)\mathbf{I}(t > \tau)] - \mathbf{E}[g(X_t)\mathbf{I}(t > \tau)] \\ &= \mathbf{E}[g(X_t)]\end{aligned}$$

using the fact that  $X_\tau = -X_\tau = 0$ .

To be very precise, the cancellation follows from the strong Markov property for Itô processes. See Øksendal<sup>2</sup> Theorem 7.2.4.

Some  
results on  
AR(1)

Ornstein -  
Uhlenbeck  
processes

Stationary  
Solutions

Remarks

Crossing-  
time  
problem

Integral  
Equation  
Methods

Martingales  
and first  
passage  
times of  
AR(1)  
sequences

Vervaat's  
Theorem

Remark

Contribu-  
tions by  
Novikov

# Ornstein - Uhlenbeck processes

OU  
processes

$$P(\tau(x) \leq t) = P(\{\tau(x) \leq t\} \cap \{X_t \leq 0\}) + P(\{\tau(x) \leq t\} \cap \{X_t < 0\})$$

$$= P(X_t \leq 0) + P(\tilde{X}_t \leq 0)$$

Some  
results on  
AR(1)

Ornstein -  
Uhlenbeck  
processes

To see this, we note that  $X_t \leq 0$  implies that  $\tau \leq t$  and hence

$$\{t \leq \tau\} \cap \{X_t \leq 0\} = \{X_t \leq 0\}$$

Stationary  
Solutions

Also

Remarks

Crossing-  
time  
problem

$$\begin{aligned} \{\tau \leq t\} \cap \{X_t > 0\} &= \{\tau \leq t\} \cap \{-X_t < 0\} \\ &= \{\tau < t\} \cap \{-X_t < 0\} \\ &= \{\tau < t\} \cap \{\tilde{X}_t < 0\} \\ &= \{\tau \leq t\} \cap \{\tilde{X}_t < 0\} \\ &\stackrel{P}{=} \{\tau \leq t\} \cap \{\tilde{X}_t \leq 0\} \\ &= \{\tilde{X}_t \leq 0\} \end{aligned}$$

Integral  
Equation  
Methods

Martingales  
and first  
passage  
times of  
AR(1)  
sequences

where we wrote  $\stackrel{P}{=}$  to indicate equality up to a set of probability zero.

Vervaat's  
Theorem

Therefore

Remark

Contribu-  
tions by  
Novikov

$$P(\tau(x) \leq t) = P(X_t \leq 0) + P(\tilde{X}_t \leq 0) = 2P(X_t \leq 0)$$

# Stationary Solutions

OU  
processes

$$X_t = \alpha X_{t-1} + \varepsilon_t \quad (1)$$

We can iterate the AR(1) equation to obtain

$$X_t = \varepsilon_t + \alpha \varepsilon_{t-1} + \alpha^2 \varepsilon_{t-2} + \dots + \alpha^{k+1} X_{t-k-1} \quad (2)$$

If  $|\alpha| < 1$  we shall show that

$$X_t = \sum_{j=0}^{\infty} \alpha^j \varepsilon_{t-j} \quad (3)$$

converges and provides the unique stationary solution to equation (1).

Let us first observe that stationarity of  $\{X_t\}$  implies that

$$\mathbf{E}[X_t^2] = \gamma(0) + \mathbf{E}[X_t]^2 = \gamma(0) + m^2 = \mathbf{E}[X_0^2]$$

for all  $t \in \mathbb{T}$ .

The sum in (3) converges in mean square as (using (2))

$$\mathbf{E} \left[ X_t - \sum_{j=0}^k \alpha^j \varepsilon_{t-j} \right]^2 = \mathbf{E} [\alpha^{2(k+1)} X_{t-k-1}^2] = \alpha^{2(k+1)} \mathbf{E}[X_0^2] \rightarrow 0$$

as  $k \rightarrow \infty$ .

Some  
results on  
AR(1)

Ornstein -  
Uhlenbeck  
processes

Stationary  
Solutions

Remarks

Crossing-  
time  
problem

Integral  
Equation  
Methods

Martingales  
and first  
passage  
times of  
AR(1)  
sequences

Vervaat's  
Theorem

Remark

Contribu-  
tions by  
Novikov

# Stationary Solutions

OU  
processes

From Chebyshev inequality

This implies convergence in probability, because

$$\mathbf{P}\left(\left|X_t - \sum_{j=0}^k \alpha^j \varepsilon_{t-j}\right| \geq \varepsilon\right) \leq \frac{1}{\varepsilon^2} \alpha^{2(k+1)} \mathbf{E}[X_0^2] \rightarrow 0$$

as  $k \rightarrow \infty$ .

In fact (3) converges almost surely. In the case when  $\{\varepsilon_t\}$  are independent, then we can use Lecture 55, Lemma 4, page 816 as

$$\sum_{j=0}^{\infty} \mathbf{E}[\alpha^{2j} \varepsilon_{t-j}^2] = \sigma^2 \sum_{j=0}^{\infty} \alpha^{2j} = \frac{\sigma^2}{1 - \alpha^2} < \infty,$$

Some  
results on  
AR(1)

Ornstein -  
Uhlenbeck  
processes

Stationary  
Solutions

Remarks

Crossing-  
time  
problem

Integral  
Equation  
Methods

Martingales  
and first  
passage  
times of  
AR(1)  
sequences

Vervaat's  
Theorem

Remark

Contribu-  
tions by  
Novikov

# Stationary Solutions

OU  
processes

## Lemma 1

If  $X$  is a random variable with  $\mathbf{E}[|X|] < \infty$ , then  $X$  is finite almost surely.

## Proof

Let

$$A = \{\omega \in \Omega : |X(\omega)| = \infty\}$$

It suffices to show that  $A \in \mathcal{F}$  and  $P(A) = 0$ .

In fact, for positive integers  $n$ , let

$$A_n = \{\omega \in \Omega : |X(\omega)| > n\}$$

Then  $A_n \in \mathcal{F}$  and so

$$A = \bigcap_{n=1}^{\infty} A_n \in \mathcal{F}$$

and for each  $n$

$$\infty > \mathbf{E}[|X|] \geq \mathbf{E}[|X| \mathbf{I}(A)] \geq n \mathbf{E}[\mathbf{I}(A)] = n P(A)$$

implies

$$P(A) \leq \frac{\mathbf{E}[|X|]}{n} \rightarrow 0$$

as  $n \rightarrow \infty$ . ■

Some  
results on  
AR(1)

Ornstein -  
Uhlenbeck  
processes

Stationary  
Solutions

Remarks

Crossing-  
time  
problem

Integral  
Equation  
Methods

Martingales  
and first  
passage  
times of  
AR(1)  
sequences

Vervaat's  
Theorem

Remark

Contribu-  
tions by  
Novikov

# Stationary Solutions

OU  
processes

We note, using monotone convergence, that

$$\begin{aligned}\mathbf{E} \left[ \sum_{j=0}^{\infty} |\alpha|^j |\varepsilon_{t-j}| \right] &= \lim_{n \rightarrow \infty} \mathbf{E} \left[ \sum_{j=0}^n |\alpha|^j |\varepsilon_{t-j}| \right] \\ &= \lim_{n \rightarrow \infty} \sum_{j=0}^n |\alpha|^j \mathbf{E} [|\varepsilon_{t-j}|] \\ &\leq \lim_{n \rightarrow \infty} \sum_{j=0}^n |\alpha|^j \sqrt{\mathbf{E} [\varepsilon_{t-j}^2]} = \frac{\sigma}{1 - |\alpha|} < \infty\end{aligned}$$

Some  
results on  
AR(1)

Ornstein -  
Uhlenbeck  
processes

Stationary  
Solutions

Remarks

Crossing-  
time  
problem

Integral  
Equation  
Methods

Martingales  
and first  
passage  
times of  
AR(1)  
sequences

Vervaat's  
Theorem

Remark

Contribu-  
tions by  
Novikov

So there exists  $\Omega_0 \subset \Omega$  with  $P(\Omega_0) = 0$  and when  $\omega \notin \Omega_0$  we have

$$\sum_{j=0}^{\infty} |\alpha|^j |\varepsilon_{t-j}(\omega)| < \infty$$

For such  $\omega \notin \Omega_0$  the expression in (3) converges because if we set

$$S_n(\omega) = \sum_{j=0}^n \alpha^j \varepsilon_{t-j}(\omega)$$

then

$$|S_n(\omega) - S_m(\omega)| \equiv \left| \sum_{j=m+1}^n \alpha^j \varepsilon_{t-j}(\omega) \right| \leq \sum_{j=m+1}^n |\alpha|^j |\varepsilon_{t-j}(\omega)| \rightarrow 0$$

as  $m, n \rightarrow \infty$ .

# Remarks

OU  
processes

1. If  $|\alpha| > 1$ , there is also a stationary solution

$$X_t = - \sum_{j=1}^{\infty} \alpha^{-j} \varepsilon_{t+j}$$

as can easily be checked. However it is not what is called **causal** as  $X_t$  depend on the future values  $\{\varepsilon_s : s > t\}$ . Such a solution is regarded as unnatural.

2. When  $|\alpha| = 1$  there is no stationary solution. For example, let  $\alpha = 1$ . If there were a stationary solution, then

$$X_t - X_0 = \varepsilon_1 + \varepsilon_2 + \cdots + \varepsilon_t$$

implies

$$\mathbf{E}[(X_t - X_0)^2] = \mathbf{E}[(\varepsilon_1 + \varepsilon_2 + \cdots + \varepsilon_t)^2]$$

But now the right hand side is  $t\sigma^2 \rightarrow \infty$  as  $t \rightarrow \infty$  while the left hand side remains bounded as

$$\mathbf{E}[(X_t - X_0)^2] \leq \mathbf{E}[2X_t^2 + 2X_0^2] = 4\mathbf{E}[X_0^2] < \infty$$

for all  $t \in \mathbf{T}$ .

When  $\alpha = -1$  we use

$$\mathbf{E}[(X_{2t} - X_0)^2] = \mathbf{E}[(-\varepsilon_1 + \varepsilon_2 - \varepsilon_3 + \cdots + \varepsilon_{2t})^2]$$

and come to a similar contradiction.

Some  
results on  
AR(1)

Ornstein -  
Uhlenbeck  
processes

Stationary  
Solutions

Remarks

Crossing-  
time  
problem

Integral  
Equation  
Methods

Martingales  
and first  
passage  
times of  
AR(1)  
sequences

Vervaat's  
Theorem

Remark

Contribu-  
tions by  
Novikov

# Remarks

3. When  $|\alpha| < 1$ , the formula for  $\gamma$  is obtained from (3). In fact, (3) implies that  $\mathbf{E}[X_t] = 0$  for all  $t \in T$  and for  $r \geq s$ ,

$$\gamma(r, s) = \mathbf{E}[X_r X_s] = \sum_{j=0}^{\infty} \sum_{k=0}^{\infty} \alpha^{j+k} \mathbf{E}[\varepsilon_{r-j} \varepsilon_{s-k}] = \sigma^2 \sum_{k=0}^{\infty} \alpha^{2k+r-s} = \frac{\alpha^{r-s}}{1-\alpha^2} \sigma^2$$

and if  $r < s$


$$\gamma(r, s) = \sum_{j=0}^{\infty} \sum_{k=0}^{\infty} \alpha^{j+k} \mathbf{E}[\varepsilon_{r-j} \varepsilon_{s-k}] = \sigma^2 \sum_{j=0}^{\infty} \alpha^{2j+s-r} = \frac{\alpha^{s-r}}{1-\alpha^2} \sigma^2$$

so


$$\gamma(h) = \frac{\alpha^{|h|}}{1-\alpha^2} \sigma^2$$

for  $h \in T$ .

ACF, AR(1),  $y_t = 0.7y_{t-1} + \varepsilon_t$



PACF, AR(1),  $y_t = 0.7y_{t-1} + \varepsilon_t$





# Crossing-time problem

OU  
processes

Let  $x \in \mathbb{R}$  and define

$$\tau(x) = \min\{n : X_n \leq 0, X_0 = x\}^3$$

and

$$\psi_k(x) = P(\tau(x) = k) \quad \text{for } k = 0, 1, 2, \dots$$

We would like to compute the values of  $\psi_k(x)$ , but this is an unsolved problem. So we need to find a method to find its values or develop approximate methods.

There are two methods to study this question.

- (i) Use martingales.
- (ii) Use integral equations.

---

<sup>3</sup>We use the convention : if  $X_n > 0$  for all  $n$ , then  $\tau(x) = \infty$

Contributions by  
Novikov

# Integral Equation Methods

OU  
processes

We suppose that the  $\{\varepsilon_t\}$  is iid with common cumulative distribution function  $F$  and density  $f$ .

We have

$$\psi_0(x) = \begin{cases} 0 & \text{if } x > 0 \\ 1 & \text{if } x \leq 0 \end{cases}$$

and for  $x > 0$

$$\psi_1(x) = P(X_1 \leq 0) = P(\varepsilon_1 \leq -\alpha x) = F(-\alpha x) .$$

We obtain the recurrence. For  $x > 0$

$$\begin{aligned} \psi_{k+1}(x) &= \mathbf{E}[\psi_k(X_1) \mathbf{I}(X_1 > 0)] \\ &= \mathbf{E}[\psi_k(\alpha x + \varepsilon_1) \mathbf{I}(\alpha x + \varepsilon_1 > 0)] \\ &= \int_{-\alpha x}^{\infty} \psi_k(\alpha x + u) f(u) du \\ &= \int_0^{\infty} \psi_k(v) f(v - \alpha x) dv \quad (\text{putting } v = \alpha x + u) \end{aligned}$$

Hence, the integral equation recurrence is

$$\psi_{k+1}(x) = \int_0^{\infty} \psi_k(v) f(v - \alpha x) dv$$

Some  
results on  
AR(1)

Ornstein -  
Uhlenbeck  
processes

Stationary  
Solutions

Remarks

Crossing-  
time  
problem

Integral  
Equation  
Methods

Martingales  
and first  
passage  
times of  
AR(1)  
sequences

Vervaat's  
Theorem

Remark

Contribu-  
tions by  
Novikov

# Integral Equation Methods

OU  
processes

We can introduce a generating function

$$\psi(x, z) = \sum_{j=1}^{\infty} \psi_k(x) z^k$$

for  $|z| \leq 1$ . Then using  $\psi_0(v) = 0$  when  $v > 0$ ,

$$\sum_{k=0}^{\infty} z^{k+1} \psi_{k+1}(x) = \int_0^{\infty} \sum_{k=0}^{\infty} z^{k+1} \psi_k(v) f(v - \alpha x) dv$$

$$= z \int_0^{\infty} \sum_{k=1}^{\infty} z^k \psi_k(v) f(v - \alpha x) dv$$

and so

$$\psi(x, z) = z \int_0^{\infty} \psi(v, z) f(v - \alpha x) dv$$

The general solution of this equation is unknown, but perhaps for some good choices of  $f$  an explicit solution may be possible. This is clearly a problem for exploration. The example in Exercise 1 could be explored.

# Martingales and first passage times of AR(1) sequences

OU  
processes

Some  
results on  
AR(1)

Ornstein -  
Uhlenbeck  
processes

Stationary  
Solutions

Remarks

Crossing-  
time  
problem

Integral  
Equation  
Methods

Martingales  
and first  
passage  
times of  
AR(1)  
sequences

Vervaat's  
Theorem

Remark

Contributions by  
Novikov

Alexander Novikov (University of Technology, Sydney) has several studies using martingale methods in the study of AR(1) processes. We will present some of these ideas, and provide some extra details not provided in these papers. While these methods will not lead to formulas for the characteristic function or the moment generating function of the crossing time, it can lead to some estimates and does lead under some technical conditions to the result that

$$P(\tau(x) < \infty) = 1$$

Applied to our study, Novikov looks at a reflection of what we are interested in. In order to be able to read the Novikov papers we revert to his setup.

## *Abstract*

Using the martingale approach we find sufficient conditions for exponential boundedness of first passage times over a level for ergodic first order autoregressive sequences. Further, we prove a martingale identity to be used in obtaining explicit bounds for the expectation of first passage times.

In applications, the distribution and expectation of such passage times are usually approximated via Monte-Carlo simulation or using Markov chain approximations (e.g. [16]). However, analytical bounds are also of interest (e.g. to control an accuracy of simulation algorithms).

# Martingales and first passage times of AR(1) sequences

OU  
processes

Let  $\varepsilon, \varepsilon_1, \varepsilon_2, \dots$  be iid random variables, and for  $0 < \alpha < 1$  and  $n = 1, 2, 3, \dots$

$$X_n = \alpha X_{n-1} + \varepsilon_n$$

and  $X_0 = x < 0$ .

We let

$$\tau = \min\{n > 0 : X_n \geq 0\}$$

The study of this crossing time is equivalent to the one we defined earlier.

As martingales will be presented, these will be with respect to the natural filtration defined by

$$\mathcal{F}_n = \sigma\{X_0, X_1, \dots, X_n\}$$

We shall assume that

$$M(u) = \mathbf{E}[\exp(u\varepsilon)] < \infty \tag{5}$$

for all  $u \geq 0$ .

Some  
results on  
AR(1)

Ornstein -  
Uhlenbeck  
processes

Stationary  
Solutions

Remarks

Crossing-  
time  
problem

Integral  
Equation  
Methods

Martingales  
and first  
passage  
times of  
AR(1)  
sequences

Vervaat's  
Theorem

Remark

Contribu-  
tions by  
Novikov

# Martingales and first passage times of AR(1) sequences

OU  
processes

Let us note that this implies that

$$\mathbf{E}[\varepsilon^+] = \mathbf{E}[\max(0, \varepsilon)] < \infty$$

This is because

$$\begin{aligned}\mathbf{E}[\varepsilon^+] &\leq \mathbf{E}[\exp(\varepsilon^+)] \\ &= \mathbf{E}[\mathbf{I}(\varepsilon \geq 0) \exp(\varepsilon)] \\ &\leq \mathbf{E}[\mathbf{I}(\varepsilon \geq 0) \exp(\varepsilon)] + \mathbf{E}[\mathbf{I}(\varepsilon < 0) \exp(\varepsilon)] = M(1) < \infty\end{aligned}$$

But (5) does not imply

$$\mathbf{E}[\varepsilon^-] = \mathbf{E}[\max(0, -\varepsilon)] < \infty$$

as the following example shows.

By the way, we recall that  $\varepsilon = \varepsilon^+ - \varepsilon^-$  and  $|\varepsilon| = \varepsilon^+ + \varepsilon^-$ .

Some  
results on  
AR(1)

Ornstein -  
Uhlenbeck  
processes

Stationary  
Solutions

Remarks

Crossing-  
time  
problem

Integral  
Equation  
Methods

Martingales  
and first  
passage  
times of  
AR(1)  
sequences

Vervaat's  
Theorem

Remark

Contribu-  
tions by  
Novikov

# Martingales and first passage times of AR(1) sequences

First, consider a martingale  $M_n$  of the form

$$M_n = \lambda^v q_v(X_n), \quad (5)$$

where a deterministic function  $q_v(y)$  depends on a parameter  $v$ , the variable  $y$  takes values from the domain  $D$  of  $X_n$ . Note that, typically,  $D = (-\infty, \infty)$  but if, for example, condition (3) holds and  $H(1 - \lambda) \geq x$  then  $D \subset (-\infty, (H/(1 - \lambda)))$ .

Under the assumption that  $M_n$  has a finite expectation, by definition of martingales

$$E[M_n | \mathcal{F}_{n-1}] = M_{n-1} \quad \text{a.s.}$$

which with (1) is equivalent to the equation

$$\lambda^v E[q_v(\lambda X_{n-1} + \eta_n) | \mathcal{F}_{n-1}] = \lambda^{v(n-1)} q_v(X_{n-1}) \quad \text{a.s.}$$

Here,  $\eta_n$  is independent of  $\mathcal{F}_{n-1}$  and  $X_{n-1}$  may take any value from the domain  $D$  of  $X_n$ . Therefore, if the function  $q_v(y)$  is a solution of the equation

$$Eq_v(\lambda y + \eta_1) = \lambda^{-v} q_v(y), \quad y \in D, \quad (6)$$

and the expectation  $Eq_v(\lambda y + \eta_n)$  is finite then  $\lambda^v q_v(X_n)$  is a martingale.

# Vervaat's Theorem

We present a result for AR(1) series which was presented in greater generality by Vervaat.<sup>8</sup>

**Theorem** (Vervaat)

Let  $\varepsilon, \varepsilon_1, \varepsilon_2, \dots$  be iid random variables and suppose that

$$\mathbf{E}[\log^+ |\varepsilon|] < \infty \quad (1)$$

Let us suppose that  $0 < |\alpha| < 1$  (for  $\alpha = 0$ , the result is trivial) and that

$$X_k = \alpha X_{k-1} + \varepsilon_k$$

for  $k = 1, 2, \dots$  and suppose  $X_0 = x$ .

Then there is a random variable  $\eta$  so that

$$X_n \xrightarrow{d} \eta \quad (2)$$

as  $n \rightarrow \infty$ . Further  $\eta$  and  $\varepsilon$  are independent and

$$\eta \stackrel{d}{=} \varepsilon + \alpha \eta. \quad (3)$$



# Vervaat's Theorem

OU  
processes

We note that

$$\log^+ |\varepsilon| \equiv \max\{0, \log |\varepsilon|\} \leq \max\{0, \log(1 + |\varepsilon|)\} = \log(1 + |\varepsilon|) \leq |\varepsilon|$$

This says that condition (1) will hold if  $\mathbf{E}[\log(1 + |\varepsilon|)] < \infty$  or  $\mathbf{E}[|\varepsilon|] < \infty$ .

We also recall that (2) means that

$$\mathbf{E}[g(X_n)] \rightarrow \mathbf{E}[g(\eta)]$$

as  $n \rightarrow \infty$  for any continuous bounded function  $g$ .

## Lemma 1

Assuming the set up of the previous theorem, condition (2) implies (3).

## Lemma 2

Assume that (1) holds and  $a > 1$ . There exists  $\Omega_0 \subset \Omega$  with  $P(\Omega_0) = 0$  so that for any  $\omega \notin \Omega_0$  there is a positive integer  $n_0 = n_0(\omega)$  so that for all  $k \geq n_0(\omega)$ ,

$$|\varepsilon_k(\omega)| \leq a^k \tag{4}$$

Some  
results on  
AR(1)

Ornstein -  
Uhlenbeck  
processes

Stationary  
Solutions

Remarks

Crossing-  
time  
problem

Integral  
Equation  
Methods

Martingales  
and first  
passage  
times of  
AR(1)  
sequences

Vervaat's  
Theorem

Remark

Contributions by  
Novikov

# Remark

OU  
processes

In the next section we will see something about the moment generating function of  $\eta$ .

Some  
results on  
AR(1)

The condition  $\mathbf{E}[\log^+ |\varepsilon|] < \infty$  often arises in the study of ARMA time series and is often required for the existence of stationary solutions. Brockwell refers to an early paper by Yohai and Maronna (1977) which indicated that for stationarity it is hard to relax this condition further.<sup>9</sup>

Ornstein -  
Uhlenbeck  
processes

We now give some further observations that follow from Vervaat's Theorem.

Stationary  
Solutions

Let us note that since

$$\mathbf{E}[\log^+ |\varepsilon|] < \infty$$

Remarks

we also have

$$\mathbf{E}[\log^+ |\varepsilon^\pm|] < \infty \quad (1')$$

Crossing-  
time  
problem

where  $\varepsilon^+ = \max(0, \varepsilon)$  and  $\varepsilon^- = \max(0, -\varepsilon)$  so that  $\varepsilon = \varepsilon^+ - \varepsilon^-$  and  $|\varepsilon| = \varepsilon^+ + \varepsilon^-$  and so  $0 \leq \varepsilon^+, \varepsilon^- \leq |\varepsilon|$  from which (1') follows.

Integral  
Equation  
Methods

We can then define two sequences

Martingales  
and first  
passage  
times of  
AR(1)  
sequences

$$\begin{aligned}\eta_n^\lambda &= \varepsilon_1^+ + \alpha \varepsilon_2^+ + \alpha^2 \varepsilon_3^+ + \cdots + \alpha^{n-1} \varepsilon_n^+ \\ \eta_n^\gamma &= \varepsilon_1^- + \alpha \varepsilon_2^- + \alpha^2 \varepsilon_3^- + \cdots + \alpha^{n-1} \varepsilon_n^-\end{aligned}$$

which each converge almost surely as  $n \rightarrow \infty$  by Vervaat's Theorem since each sequence  $\{\varepsilon_n^+\}$  and  $\{\varepsilon_n^-\}$  are iid. Let the limits be  $\eta^\lambda$  and  $\eta^\gamma$ .

Vervaat's  
Theorem

Let us note that  $\eta_n^\lambda \uparrow \eta^\lambda$  and  $\eta_n^\gamma \downarrow \eta^\gamma$  as  $n \rightarrow \infty$ , that  $\eta^\gamma \leq \eta_n \leq \eta^\lambda$  for each  $n$  and  $\eta = \eta^\lambda - \eta^\gamma$ .

Remark

We will use these constructions in some of the arguments that follow in the next section.

Contributions  
by  
Novikov

## Lemma A

If (5) holds for  $u \geq 0$  and  $\mathbf{E}[|\varepsilon|] < \infty$ , then the right hand derivative of  $M(u)$  at  $u = 0$  exists and is given by

$$M'(0+) = \mathbf{E}[\varepsilon]$$

## Remark

We note that  $\mathbf{E}[\varepsilon^+] \leq M_+(u) \leq M(u) < \infty$ . We also note that  $\mathbf{E}[\varepsilon^-] = \infty$  if and only if  $M'(0+) = -\infty$ .

# Contributions by Novikov

OU  
processes

The **cumulant function** of  $\varepsilon$  is defined by

$$\psi(u) = \log \mathbf{E} [\exp (u \varepsilon)]$$

## Remarks

1. The cumulant function is convex. Suppose that  $\psi(s)$  and  $\psi(t)$  are finite and  $0 < \lambda < 1$ , then

$$\lambda \psi(s) + (1 - \lambda) \psi(t) = \log [\mathbf{E}[\exp(s \varepsilon)]^\lambda \mathbf{E}[\exp(t \varepsilon)]^{1-\lambda}]$$

$$\geq \log [\mathbf{E}[\exp((\lambda s + (1 - \lambda) t) \varepsilon)]]$$

$$= \psi(\lambda s + (1 - \lambda) t)$$

In the process we used Hölder's inequality. In fact

$$\mathbf{E}[\exp((\lambda s + (1 - \lambda) t) \varepsilon)] = \mathbf{E}[\exp(\lambda s \varepsilon) \exp((1 - \lambda) t \varepsilon)]$$

$$\leq \mathbf{E}[\exp(p \lambda s \varepsilon)]^{\frac{1}{p}} \mathbf{E}[\exp(q (1 - \lambda) t \varepsilon)]^{\frac{1}{q}}$$

Where  $p, q > 1$  and  $\frac{1}{p} + \frac{1}{q} = 1$ . Choose  $p = 1/\lambda$  and  $q = 1/(1 - \lambda)$ .

2. It then follows that for  $0 < u < 1$  that

$$\psi(u) = \psi(u \cdot 1 + (1 - u) \cdot 0) \leq u \psi(1)$$

as  $\psi(0) = 0$ .

3. If  $\mathbf{E}[|\varepsilon|] < \infty$ , then from Lemma A,

$$\psi'(0+) = \mathbf{E}[\varepsilon]$$

Some  
results on  
AR(1)

Ornstein -  
Uhlenbeck  
processes

Stationary  
Solutions

Remarks

Crossing-  
time  
problem

Integral  
Equation  
Methods

Martingales  
and first  
passage  
times of  
AR(1)  
sequences

Vervaat's  
Theorem

Remark

Contribu-  
tions by  
Novikov

## Lemma B

If  $\mathbf{E}[|\varepsilon|] < \infty$ , then

$$\phi(u) = \sum_{k=0}^{\infty} \psi(\alpha^k u) \quad (8)$$

is convergent. It is differentiable for  $u > 0$ . It satisfies

$$\phi(u) = \log \mathbf{E}[\exp(u \eta)] \quad (9)$$

where  $\eta$  is the same random variable as in the previous section, and

$$\phi(u) = \phi(\alpha u) + \psi(u) \quad (10)$$

## Remark

Novikov<sup>10</sup> shows that (8) converges under the weaker condition

$$\mathbf{E}[\log^+ |\varepsilon|] < \infty \quad (11)$$

# Remark

Under the weaker condition

$$\mathbf{E}[\log^+ |\varepsilon|] < \infty$$

there are two cases (1)  $\mathbf{E}[\varepsilon^-] < \infty$  and (2)  $\mathbf{E}[\varepsilon^-] = \infty$ . In the first case we have  $\mathbf{E}[|\varepsilon|] < \infty$  and we have the assumptions of Lemma B. In the second case we have  $\psi'(0+) = -\infty$ . This implies that for some  $u_0 > 0$ ,  $\psi(u) < 0$  for  $0 < u < u_0$ . This implies that

$$\lim_{n \rightarrow \infty} \sum_{k=0}^{n-1} \psi(\alpha^k u)$$

is finite or  $-\infty$ . But the latter is not possible as

$$\begin{aligned} \liminf_{n \rightarrow \infty} \sum_{k=0}^{n-1} \psi(\alpha^k u) &= \liminf_{n \rightarrow \infty} \log \mathbf{E}[\exp(u \eta_n)] \\ &\geq \liminf_{n \rightarrow \infty} \log \mathbf{E}[\exp(-u \eta_n^\gamma)] \\ &= \log \mathbf{E}[\exp(-u \eta^\gamma)] > -\infty \end{aligned}$$

# Remark

This follows because  $\mathbf{E}[\log^+ |\varepsilon^-|] < \infty$  implies that  $0 \leq \eta^\gamma < \infty$  with probability 1, and if  $\log \mathbf{E}[\exp(-u \eta^\gamma)] = -\infty$  for  $u > 0$  we would have a contradiction. This implies that under the condition (11) that  $\phi(u)$  defined in (8) exists.

The proof of the Lemma B under (11) now follows as we have already described.

It is a further corollary to these discussions that

$$\phi(u) \geq \log \mathbf{E}[\exp(-u \eta^\gamma)] \geq -K = \log \mathbf{E}[\exp(-\eta^\gamma)] > -\infty$$

for all  $0 \leq u \leq 1$ .