

Home Assignment 2.

1. Set random seed to $M * 100 + D$, where M and D is your Month and Date of birth, respectively.
Generate two samples X of size 20 and Y of size 200 s.t. $E(X) \sim Unif[0, 1]$, $E(Y) \sim Unif[0, 1]$ from
 - two normal distributions with the same variance $\sigma_X^2 = \sigma_Y^2 = 0.1$;
 - two normal distributions with if M is odd $\sigma_X^2 = 0.1$ and $\sigma_Y^2 = 1$, else $\sigma_X^2 = 1$ and $\sigma_Y^2 = 0.1$
 - two normal distributions with 3 outliers added afterwards $Out_i \sim Exp(10) * (-1)^{U\{1,2\}}$;
 - two exponential distributions.
 - (a) Conduct t -test, Welch test, Mann-Whitney test 10000 times.
 - (b) Calculate nominal type I error rate and power (to calculate type I error fix mean for Y the same as for X).
 - (c) For each case described above plot how nominal power changes for different values of parameters of the distribution.
 - (d) Based on the plots, which test is the most appropriate in each of the cases. Will your answer change if both sample size are increased by 10.
2. Use dataset `SIC33.csv` with the following data:
 - *output* – Value added.
 - *labor* – Labor input.
 - *capital* – Capital stock.

Consider the following model $\ln Y = \beta_0 + \beta_1 \ln(K) + \beta_2 \ln(L) + u$. Suppose we are interested in estimating the quantity $\gamma = \beta_1 + \beta_2$ from the data. Test $H_0 : \gamma = 1$ against $H_a : \gamma \neq 1$ using:

- (a) Jackknife estimator;
- (b) paired bootstrap;
- (c) wild bootstrap (bootstrapping the residuals).

How the procedure should be changed in case u is heteroskedastic?

3. Consider K and L from the same dataset `SIC33.csv`.
 - (a) Illustrate principal components on a scatter plot.
 - (b) Calculate the explained variance by each component, the weight of K in the 1st component and the weight of 2nd component in the reconstruction of L .
 - (c) Repeat the analysis for $\ln(K)$ and $\ln(L)$.

Deadline: 2022-12-18, 21:00.