


## Week 2

July 17, 2014


# plotter script

- ▶ 324 sloc
- ▶ reads RotationShield and FieldMapping VI input
- ▶ uses new normalization method
  - ▶ average of data points near  $(0, 0, 0)$  vs. polynomial fit
  - ▶ calculates desired normalization level - average  $B_x$  of measured maps
- ▶ handles custom field slices
- ▶ to-do: field gradients, interpolation, smooth plots

# 10 cm offset between map & simulation



# 10 cm offset between map & simulation



modifications to account for 10 cm offset

## modifications to account for 10 cm offset

- ▶ varied metglas thickness from 5 cm to 1 mm (closest to actual)

## modifications to account for 10 cm offset


- ▶ varied metglas thickness from 5 cm to 1 mm (closest to actual)
- ▶ extended metglas slightly (2 cm) above  $B_0$  coil

## modifications to account for 10 cm offset


- ▶ varied metglas thickness from 5 cm to 1 mm (closest to actual)
- ▶ extended metglas slightly (2 cm) above  $B_0$  coil
- ▶ extended metglas far (10 cm) above  $B_0$  coil to highlight effects




varying thickness: small change in  $B$  magnitude, no shift




varying thickness: small change in  $B$  magnitude, no shift




## 2 cm longer on top: small magnitude change




## 2 cm longer on top: small magnitude change



# 10 cm longer on top: confirms magnitude change



# 10 cm longer on top: confirms magnitude change



## other things to try

- ▶ metglas shearing - hard to model

## other things to try

- ▶ metglas shearing - hard to model
  - ▶ RotationShield handles arbitrary fields (e.g. you can put a line current anywhere)




## other things to try

- ▶ metglas shearing - hard to model
  - ▶ RotationShield handles arbitrary fields (e.g. you can put a line current anywhere)
  - ▶ but before applying the field it builds an interaction matrix - requires azimuthal symmetry


## other things to try

- ▶ metglas shearing - hard to model
  - ▶ RotationShield handles arbitrary fields (e.g. you can put a line current anywhere)
  - ▶ but before applying the field it builds an interaction matrix - requires azimuthal symmetry
  - ▶ would probably affect azimuthal symmetry in measured map


# azimuthal symmetry: $0^\circ$




azimuthal symmetry:  $180^\circ$



azimuthal symmetry:  $0^\circ$ ,  $B_z$  axis flipped



azimuthal symmetry:  $180^\circ$



## other things to try

- ▶ metglas shearing - hard to model
  - ▶ RotationShield handles arbitrary fields (e.g. you can put a line current anywhere)
  - ▶ but before applying the field it builds an interaction matrix - requires azimuthal symmetry
  - ▶ would probably affect azimuthal symmetry in measured map

## other things to try

- ▶ metglas shearing - hard to model
  - ▶ RotationShield handles arbitrary fields (e.g. you can put a line current anywhere)
  - ▶ but before applying the field it builds an interaction matrix - requires azimuthal symmetry
  - ▶ would probably affect azimuthal symmetry in measured map
  - ▶ either the shear doesn't cause a peak shift



## other things to try

- ▶ metglas shearing - hard to model
  - ▶ RotationShield handles arbitrary fields (e.g. you can put a line current anywhere)
  - ▶ but before applying the field it builds an interaction matrix - requires azimuthal symmetry
  - ▶ would probably affect azimuthal symmetry in measured map
  - ▶ either the shear doesn't cause a peak shift
  - ▶ or it's aligned exactly along the  $y$ -axis

## other things to try

- ▶ metglas shearing - hard to model
  - ▶ RotationShield handles arbitrary fields (e.g. you can put a line current anywhere)
  - ▶ but before applying the field it builds an interaction matrix - requires azimuthal symmetry
  - ▶ would probably affect azimuthal symmetry in measured map
  - ▶ either the shear doesn't cause a peak shift
  - ▶ or it's aligned exactly along the  $y$ -axis
- ▶ rigorously check centering, dimensions of experimental setup