

MATH172 Galois Theory Notes

Xuehuai He
October 26, 2023

Contents

Rings! Or why $x^2 - 2$ has roots.	3
Ring	3
Phase I: ID, UFD, PID, ED, Fields	3
Zero divisors	3
Integral domain	4
Unit, irreducibles	4
Unique factorization domain	4
Ideals	4
Principal Ideal Domain	4
Prime Ideal	5
Maximal Ideal	5
Noetherian Rings	5
Quotient Rings	6
Ring homomorphism	7
First ring isomorphism theorem	7
Quotient by prime ideal is ID	7
Quotient by maximal ideal is field	8
Euclidean Domain	8
Why do we care about Euclidean domains?	9
Greatest common divisors & Euclidean algorithm	9
Prime elements	9
Prime implies irreducible in ID	10
Prime implies maximal in PID	10
Irreducible implies prime in UFD	10
PIDs are UFDs	11
Phase I summary	12
Field extensions	12
Phase II: Field extensions	12
$F[x]/(p(x))$ contains a copy of F	13

$F[x]/(p(x))$ field if $p(x)$ irreducible	13
Field extension, degree of extension	13
To summarize so far!	14
Irreducibility – a survey	15
Eisenstein’s Criterion	16
Gauss’ Lemma	16
Algebraic and transcendental	17
Algebraic extension	17
Finite extensions are algebraic	17
Minimal polynomial	17
Degree of extension is multiplicative	18
Algebraic elements form a field	19
Extension finite iff. adjoin finite algebraic elements	19
Composite field	20
Splitting field	20
Normal extensions	20
Splitting fields exist	20
Splitting fields are unique	21
Separable polynomials	22
Formal derivatives	22
Roots repeated iff. roots of derivative	22
Irreducible non-separables have $f'(x) = 0$	22
TODO	22
Irreducible polys are separable in finite or char 0 fields	22
Prime fields	23
Every field is an extension of \mathbb{Q} or \mathbb{F}_p	24
Finite fields are unique	24
Separable extensions	24
Galois extensions	24
Galois Theory	24
Automorphism group of field extensions	24
Fundamental Theorem of Galois Theory	24
Phase III: Galois Theory	25
Roots of minimal polys under automorphisms is still a root	25
Galois extensions (alternate def)	26

Rings! Or why $x^2 - 2$ has roots.

Definition 1. A **ring** is a set R together with associative binary *operations* $+$ and \times s.t.:

← map from
 $R \times R \mapsto R$

- $(R, +)$ is an **abelian** group with identity 0
- There exists $1 \in R$ s.t. $r \times 1 = 1 \times r = r$
- $r(s + t) = rs + rt$ and $(s + t)r = sr + tr \quad \forall s, r, t \in R$

← this is optional

Proposition 1. $0 \times 1 = 0$ (in fact, $0 \times r = 0 \quad \forall r \in R$)

Proof. Try it!

□

Definition 2. If \times is commutative, then R is a commutative ring.

Non-example 1. \mathbb{N} is not a ring.

Example 2. $\mathbb{Z} \subseteq \mathbb{Q} \subseteq \mathbb{R} \subseteq \mathbb{C}$ are all rings;

- $\mathbb{Z}/n\mathbb{Z}$ is a finite ring
- $M_n(\mathbb{R})$, the set of $n \times n$ real matrices, is a **noncommutative** ring
- Polynomial ring: $\mathbb{Q}[x] = \{a_0 + a_1x + \cdots + a_nx^n \mid a_i \in \mathbb{Q}\}$ is a commutative ring
- $\mathbb{Z}[\sqrt{-5}] = \{a + b\sqrt{-5} \mid a, b \in \mathbb{Z}\}$ is a commutative ring

← square brackets just mean "polynomials in..."

Phase I plan:

$$ID \supsetneq UFD \supsetneq PID \supsetneq ED \supsetneq Fields$$

Definition 3. Suppose R is a ring and $a, b \in R$ with $ab = 0$ but $a, b \neq 0$; then a, b are called **zero divisors**.

Example 3.

- In $\mathbb{Z}/6\mathbb{Z}$, $\bar{4} \times \bar{3} = \bar{0}$
- In $M_2(\mathbb{R})$, $\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}$

Definition 4. A commutative ring without zero divisors is called an integral domain (ID)

Why do we want ID? **Cancellation properties.**

- If R is an ID, $a, b, c \in R$, $a \neq 0$ and $ab = ac$, then

$$ab - ac = 0 \implies a(b - c) = 0 \implies b - c = 0 \implies b = c$$

Definition 5. Suppose R is an ID. An element $a \in R$ is called a **unit** if $a \neq 0$ and there exists $b \in R$ s.t. $ab = 1$.

← notation: $b = a^{-1}$

An element $r \in R$ is called **irreducible** if $r \neq 0$, r is NOT a unit, and whenever $r = ab$ for some $a, b \in R$ then a or b must be a unit.

- If r and s are irreducibles with $r = us$, then r and s are called **associates**.

Example 4.

- All “prime integers” are irreducibles in \mathbb{Z} ;
- $2, 3, 1 + \sqrt{-5}, 1 - \sqrt{-5}$ are irreducibles in $\mathbb{Z}[\sqrt{-5}]$.
 - Note: $2 \times 3 = (1 + \sqrt{-5})(1 - \sqrt{-5}) = 6$ says that 6 can be factored in more than one way. This means that $\mathbb{Z}[\sqrt{-5}]$ is NOT an UFD.

Definition 6. An integral domain R is called a unique factorization domain (UFD) if each nonzero, nonunit $a \in R$ can be written as a product of irreducibles **in a unique way** up to associates.

If a is a nonzero, nonunit element of UFD R and $a = r_1 r_2 \dots r_m = s_1 \dots s_n$ where r_i, s_j are irreducible, then after reordering $r_i = u_i s_i$ for any i and units u_i , and $m = n$.

← After reordering, there are the same amounts of factors and all factors are the same up to units.

Definition 7. Suppose R is a comm ring. A subset $I \subseteq R$ is called an **ideal** if $(I, +) \leq (R, +)$ and $ir, ri \in I$ for all $i \in I$ and for all $\boxed{r \in R}$.

Why do we want ideals? Such that R/I is a well-defined ring.

Example 5. $\{0\}$ and R are ideals of R .

Example 6. If R is commutative and $a \in R$, then $(a) = \{ar \mid r \in R\}$ is called the **principal ideal** generated by a .

← Prove this (be convinced)!
Also known as aR .

Definition 8. A **principal ideal domain** is an integral domain where all ideals are principal ideals.

Example 7. The only ideals of $(\mathbb{Z}, +)$ are of the form $n\mathbb{Z} = (n)$.

← Ideals generated by n

Non-example 8. $\mathbb{Z}[x]$ is a UFD but NOT a PID because the ideal $(2, x) = \{2r + xs \mid r, s \in \mathbb{Z}[x]\}$ is not principal.

← Observe that $(2, x)$ is an ideal made of polynomials with even constant terms. This cannot be principal, since if we only have 2

Lemma 2. If $I \subseteq R$ is an ideal and $1 \in I$, then $I = R$.

Proof. Try it!

Proposition 3. If $I \subseteq R$ is an ideal containing a unit of R then $I = R$.

Proof. If $u \in I$ is a unit then $u^{-1} \in R$, so $uu^{-1} = 1 \in I$. Then the result follows from Lemma 2. \square

Definition 9. A **field** is a commutative ring whose each nonzero element is a unit.

Corollary 4. If R is an ID whose ideals are (0) and R , then R is a **field**.

Proof. Suppose $a \in R \setminus \{0\}$ and consider (a) . Since $a \in (a)$, $(a) = R$. Hence, we must have that $1 \in (a)$, which means $1 = ar$ for some $r \in R$. \square

Definition 10. Suppose R is an integral domain. A *proper* ideal $P \subsetneq R$ is called **prime** if whenever $ab \in P$ for some $a, b \in R$, then a or $b \in P$.

Non-example 9. (6) is not a prime ideal of \mathbb{Z} since $2 \times 3 \in (6)$ but neither $2, 3 \in (6)$.

Non-example 10. (2) is not a prime ideal of $\mathbb{Z}[\sqrt{-5}]$ since $6 \in (2)$, but we observe that $6 = (1 + \sqrt{-5})(1 - \sqrt{-5})$ while $1 \pm \sqrt{-5} \notin (2)$.

Example 11. (2) is a prime ideal of \mathbb{Z} .

Definition 11. A proper ideal $M \subsetneq R$ is called **maximal** if whenever $I \subseteq R$ such that $M \subseteq I \subseteq R$ is an ideal containing M , then either $I = M$ or $I = R$.

Proposition 5. Every proper ideal is contained in a maximal ideal.

Proof. Axiom of choice. \square

Proposition 6. Suppose R is a commutative ring.

- (0) is prime *if and only if* R is an integral domain.
- (0) is maximal *if and only if* R is a field.

(The following is kind of on a tangent)

Definition 12. A commutative ring R with unity is called **Noetherian** if, whenever $I_1 \subseteq I_2 \subseteq \dots$ is an ascending sequence of (proper) ideals of R , there exists an $n > 0$ such that $I_n = I_{n+1} = \dots$ are the same ideals thereafter.

Theorem 7. R is Noetherian *if and only if* all ideals of R are finitely generated.

← The converse is also true. **The only ideals in a field are 0 and the field.**

← Observe that in $\mathbb{Z}[\sqrt{-5}]$, we have $6 = (1 + \sqrt{-5})(1 - \sqrt{-5}) = 2 \times 3$, so it is not a UFD!

← This might not be unique in non-local rings.

← By def of prime, if $ab = 0$, then either $a = 0$ or $b = 0$, which means there are NO zero divisors.

← The chain stops ascending!

Corollary 8. All Principal Ideal Domains are Noetherian.

← Since all ideals are generated by 1 elt.

(Tangent ends here)

Definition 13. Suppose R is a commutative ring with $1 \neq 0$ and $I \subseteq R$ is an ideal. Then the quotient ring of R by I is the set

$$R/I = \{r + I \mid r \in R\}$$

with addition and multiplication defined representative-wise.

Remark. The **coset criterion** of ideals: let I be an ideal; the cosets $r + I, s + I$ are the same *if and only if* $r - s \in I$.

Example 12.

- In $\mathbb{Z}/(6)$ aka. $\mathbb{Z}/6\mathbb{Z}$, we have $2 + (6) = \{\dots, -10, -4, 2, 8, 14, \dots\} = 26 + (6)$ due to $2 - 26 \in (6)$;
- In $\mathbb{Q}[x]/(x^2 - 2)$, we have

$$\{3x^2 - 47x + 1 + q(x)(x^2 - 2) \mid q(x) \in \mathbb{Q}[x]\} = \{-47x + 7 + q(x)(x^2 - 2) \mid q(x) \in \mathbb{Q}[x]\}$$

$$\text{due to } 3x^2 - 47x + 1 - (-47x + 7) \in (x^2 - 2).$$

Remark. Let I be an ideal of R . Then $(I, +) \trianglelefteq (R, +)$.

Definition 14. R/I is a group under $(r + I) + (s + I) = (r + s) + I$ and the operation $+$ is well-defined. We also define that $(r + I)(s + I) = (rs) + I$. We claim that multiplication in R/I is also well-defined.

Proof. Let $r_1 + I = r_2 + I$ and $s_1 + I = s_2 + I$. By coset criterion, $r_1 - r_2 = i, s_1 - s_2 = j$ for some $i, j \in I$. Hence $r_1 s_1 = (r_2 + i)(s_2 + j) = r_2 s_2 + i s_2 + j r_2 + ij$ where the latter three terms are all in the ideal I . Thus, $(r_1 s_1) + I = (r_2 s_2) + I$. \square

From R , R/I inherits nice properties:

- $0 + I = 0_{R/I}$
- $1 + I = 1_{R/I}$
- Multiplication is commutative and distributive over addition in R/I , so it is also a comm. ring with identity.

Definition 15. A function $\varphi : R \rightarrow S$ between rings is called a **ring homomorphism** if the following are satisfied:

- $\varphi(r_1 + r_2) = \varphi(r_1) + \varphi(r_2)$
- $\varphi(r_1 r_2) = \varphi(r_1) \varphi(r_2)$

Theorem 9. First ring isomorphism theorem

If $\varphi : R \mapsto S$ is a ring homomorphism, then $R / \ker(\varphi) \cong \varphi(R)$.

Example 13. If R is a ring and I is an ideal, then $\pi : R \rightarrow R/I$ where $r \mapsto r + I$ is a surjective homomorphism where $\ker(\pi) = I$. This is the *canonical projection* onto R/I .

Corollary 10. If I is a maximal ideal, then R/I is a field.

Recall Proposition 6. We now have a stronger statement:

Proposition 11. Suppose R is a commutative ring & $P \subseteq R$ is an ideal. Then R/P is an integral domain *if and only if* P is prime.

Proof. R/P is an integral domain *if and only if* whenever $(a+P)(b+P) = 0_{R/P}$ then one of $a+P$ or $b+P$ must already be $0_{R/P}$. This happens *if and only if* whenever $ab+P = P$ then $a+P$ or $b+P$ in P , which happens *if and only if* whenever $ab \in P$ then one of $a, b \in P$, which is the definition of a prime ideal. \square

Example 14. The map $\varphi : \mathbb{Z}[x] \rightarrow \mathbb{Z}$ where $p(x) \mapsto p(0)$ is a surjective ring homomorphism with $\ker(\varphi) = (x)$. By the the First Isomorphism Theorem 9, $\mathbb{Z}[x]/(x) \cong \mathbb{Z}$. As such, we conclude that (x) is a prime ideal since \mathbb{Z} is an integral domain.

Lemma 12. Suppose R is a comm. ring with $M \subseteq R$ being an ideal.

There is a bijective correspondence between the ideals of R/M and the ideals of R containing M .

Proof. Consider the projection $\pi : R \rightarrow R/M$ where $r \mapsto r + M$. It is enough to show:

$$\begin{aligned} \pi(\pi^{-1}(J)) &= J & \text{for all ideals } J \subseteq R/M, \text{ and} \\ \pi^{-1}(\pi(I)) &= I & \text{for all ideals } M \subseteq I \subseteq R \end{aligned}$$

To prove the first statement, observe that, if J is an ideal of R/M , then $\pi^{-1}(J) = \{r \in R \mid r + M \in J\}$ and so

$$\pi(\pi^{-1}(J)) = \{\pi(r) \in R/M \mid r + M \in J\} = \{r + M \mid r + M \in J\} = J$$

← Observe that kernels are ideals! And ideals are kernels of some homomorphism too.

← The *if and only if* version comes in Proposition 14.

← btw, $(x) \subsetneq (x, 2)$. the latter is the set of polynomials whose constant term is even, so it is also a proper ideal of $\mathbb{Z}[x]$. This is an excellent example where Prime \nRightarrow Maximal.

← To see why this is okay, see Homework 2 Sec. 7.3 P. 24

Next, to prove the second statement, first suppose $M \subseteq I \subseteq R$ is an ideal. Let $a \in I$. Then $a + M \in \{\alpha + M \mid \alpha \in I\} = \pi(I)$. This implies that $a \in \pi^{-1}(\pi(I))$, and so $I \subseteq \pi^{-1}(\pi(I))$.

Conversely, suppose $r \in \pi^{-1}(\pi(I))$. This is the same as saying $\pi(r) = r + M \in \pi(I) = \{\alpha + M \mid \alpha \in I\}$. Hence, for any $r \in \pi^{-1}(\pi(I))$, there exists some $a \in I$ such that $r + M = a + M$. Thus, $r - a \in M \subseteq I$ by coset conditions. Since $a \in I$, we have $a + (r - a) \in I$, meaning that $r \in I$ for any $r \in \pi^{-1}(\pi(I))$. This means that $\pi^{-1}(\pi(I)) \subseteq I$.

Hence, $I = \pi^{-1}(\pi(I))$.

Consequently, for any ideals $J \subseteq R/M$, we know that $\pi^{-1}(J) \subseteq R$ is an ideal containing M . And if $M \subseteq I \subseteq R$ is an ideal, we know $\pi(I) \subseteq R/M$ is an ideal. Since $\pi(\pi^{-1}(J)) = J$ and $I = \pi^{-1}(\pi(I))$ for any I, J , the correspondence is a bijection. \square

← Think about why this contains M !

Proposition 13. Suppose R is a comm. ring with an identity and $I \subseteq R$ is an ideal. Then R/I is a field if and only if I is maximal.

Proof. If I is maximal, then there are no other proper ideals strictly containing I . Hence, by Lemma 14, we have that R/I only have ideals (0) and R/I itself. This happens if and only if R/I is a field. \square

Corollary 14. If R is a commutative ring with identity and $M \subseteq R$ is maximal, then M is prime.

← Hence maximal implies prime, but prime does not necessarily implies maximal.

Proof. Maximal \implies quotient is a field \implies quotient is an ID \implies prime. \square

Definition 16. An integral domain R is an **Euclidean domain** if there exists a norm $N : R \rightarrow \mathbb{Z}_{\geq 0}$ with $N(0) = 0$ such that for all $a, b \in R$ with $b \neq 0$, there exists $q, r \in R$ for which

$$a = bq + r$$

with $N(r) < N(b)$ or $r = 0$.

Example 15. \mathbb{Z} is a ED with $N(a) = |a|$.

Example 16. $\mathbb{Q}[x]$ is a ED with $N(p(x)) = \deg(p(x))$.

Example 17. Every field F is a ED with $N(a) = 0 \forall a \in F$.

Non-example 18. $\mathbb{Z}\left[\frac{1+\sqrt{-19}}{2}\right]$ is a PID that is not an ED.

← Because in a field everything divides!

← This is one of the only good examples!

Why do we care about Euclidean domains?

Remark. Greatest common divisors exist and are relatively quick to compute.

← Using recursive application of Euclidean algorithm.

Definition 17. If $a, b \in R$, then $\gcd(a, b) = c$ means

1. c divides a and b ; that is, $a = cr, b = cs$ for some $r, s \in R$
2. If $c' \in R$ with $c'|a$ and $c'|b$, then it must be true that $c'|c$.

← All other common divisors divide the gcd.

Example 19. Say we want to compute the gcd of 47 and 10.

$$47 = 4 \times 10 + 7$$

$$10 = 1 \times 7 + 3$$

$$7 = 2 \times 3 + \textcircled{1}$$

$$3 = 3 \times 1$$

← circled is $\gcd(47, 10)$

← final line with no remainders

← This is a much faster algorithm than factoring!

This also works for finding gcds in $\mathbb{Q}[x]$ with polynomials long division and norm $\deg(p(x))$.

Remark. If F is a field, then $F[x]$ is a Euclidean domain.

← Just use long division!

Remark. Euclidean domains are PIDs.

Proof. Suppose R is a ED and $I \subseteq R$ is an ideal. Consider $\{N(a) \mid a \in I \setminus \{0\}\}$. This set has a minimal element by properties of natural numbers (or is an empty set if and only if $I = (0)$).

Let $d \in I$ be an element of minimum norm (hence $N(d) \leq N(a)$ for all $a \in I$). We claim that $\underline{(d) = I}$. Proof:

Since $d \in I$, we have $rd \in I$ for any $r \in R$. This implies that $\underline{(d) \subseteq I}$.

Then let $a \in I$. Since R is a ED, we first assume that there exists $q, r \in R, r \neq 0$ such that $a = qd + r$ and $N(r) < N(d)$. But we notice that $r = a - qd$ must be in I as both $a, qd \in I$, contradicting the minimality of $N(d)$. Thus, it must be that $r = 0$. This implies $a = qd$ and thus $a \in (d)$ for all $a \in I$. Consequently, $I \subseteq (d)$, and therefore $I = (d)$. \square

Definition 18. Suppose R is an integral domain and $p \in R \setminus \{0\}$. Then p is a **prime element** if (p) is a prime ideal.

Proposition 15. An element $p \in R$ is prime if and only if whenever $p|ab$ then $p|a$ or $p|b$.

Proof. p is prime means that (p) is a prime ideal. This is true *if and only if* whenever $ab \in (p)$ then $a \in (p)$ or $b \in (p)$. This is the same as saying if $ab = kp$ for some $k \in R$ then $a = lp$ or $b = lp$ for some $l \in R$. This is to say that whenever $p|ab$ then $p|a$ or $p|b$. \square

Proposition 16. In an integral domain, all prime elements are irreducibles.

Proof. Suppose R is an ID and $p \in R$ is prime. If $p = ab$ for some a, b in R , then, WLOG, $p|a$. That is, $a = pk$ for some $k \in R$. Hence, $p = pkb$. Since in an ID cancellation rule holds, $kb = 1$, meaning that b is a unit. Thus, p is irreducible by definition Definition 5. \square

Proposition 17. In PIDs, all *nonzero* prime ideals are maximal.

Proof. Suppose R is a PID and $(p) \subseteq R$ is a prime ideal. If $(p) \subseteq (m) \subseteq R$ is an ideal, then $p \in (p) \subseteq (m)$ hence $p = rm$ for some $r \in R$. Since $p|rm$, we have $p|r$ or $p|m$.

If $p|r$, this implies that $r = pk$ for some $k \in R$. Substituting into $p = rm$, we get $p = pkm$. By cancellation, we get $km = 1$, meaning that m is a unit. Hence, $(m) = R$.

If $p|m$, we have $m = pl$ for some $l \in R$, meaning that $m \in (p)$. Hence, $(m) \subseteq (p)$, but we also defined that $(p) \subseteq (m)$, so $(m) = (p)$.

Therefore, (p) has to be the maximal ideal. \square

Proposition 18. In an UFD, irreducible implies prime.

Proof. Let R be a UFD and $p \in R$ be irreducible. Let $a, b \in R$ such that $p|ab$. Hence, $pr = ab$ for some $r \in R$. Since R is a UFD, let $a = q_1 \dots q_n, b = s_1 \dots s_m$ be the factorization. Since the factorizations are unique and each of the q_i, s_j are irreducible, if $p|ab$, then p must be an associate with one of the q_i, s_j . Therefore, either $p|a$ or $p|b$, implying prime. \square

Example 20. \mathbb{Q} is a field, so $\mathbb{Q}[x]$ is a ED. Since EDs are UFDs, irreducible \implies prime. We see that $x^2 - 2 \in \mathbb{Q}[x]$ is an irreducible element, which means that $(x^2 - 2)$ is a prime ideal, meaning that it is a maximum ideal, meaning that $\mathbb{Q}[x]/(x^2 - 2)$ is a field. We observe that it is a field containing \mathbb{Q} and $(\sqrt{2})$.

← In fact, this is the smallest field containing \mathbb{Q} and $(\sqrt{2})$.

Lemma 19. In a PID, irreducible elements are prime.

Proof. Suppose $p \in R$ is irreducible in the principal ideal domain R . If $p|ab$ for some $a, b \in R$, we want to show that either $p|a$ or $p|b$, hereby showing that p is prime. Hence, we consider the ideal $(a, p) = d$, which is necessarily principal for some $d \in R$. Since $a, p \in (d)$, we have $a = dr$ and $p = ds$ for some $r, s \in R$. As p is irreducible, we get that one of d and s is a unit.

We first assume that s is a unit, in which case $d = ps^{-1}$, and so $a = ps^{-1}r$ implying that $p|a$.

In another case, d is a unit, in which case $(a, p) = (d) = R$ and so $1 = ak + pl$ for some $k, l \in R$. Multiplying by b , we get $b = abk + pbl$. Since $p|ab$, we have $b = abk + pbl = pmk + pbl$ for some $m \in R$. Hence, $b = p(mk + bl)$, meaning that $p|b$.

Therefore, whenever $p|ab$, either $p|a$ or $p|b$. Hence, in a PID, p is prime whenever it is irreducible. \square

Proposition 20. PIDs are UFDs.

Proof. Suppose R is a PID and $a \in R$ is nonzero, nonunit. If a is irreducible, we are done. If not, we write $a = p_1q_1$ for some $p_1, q_1 \in R$ nonunit. If p_1, q_1 are irreducibles, we are done. If not, then WLOG say $q_1 = p_2q_2$ for some nonunits p_2, q_2 . We would like to show that this splitting process terminates.

Observe that $(q_1) \subseteq (q_2)$ since $q_2|q_1$. Hence, the chain of splitting results in the chain of ideals $(q_1) \subseteq (q_2) \subseteq (q_3) \subseteq \dots$.

Now consider the ideal $\bigcup_{i=1}^{\infty} (q_i)$. Since this is a PID, we have $\bigcup_{i=1}^{\infty} (q_i) = (q)$ for some $q \in R$. Since $q \in \bigcup_{i=1}^{\infty} (q_i)$, it is contained in some (q_n) for some $n \geq 1$. This implies that $(q) \subseteq (q_n)$, but we also know that $(q_n) \subseteq (q)$, hence $(q) = (q_n)$. Hence, this process terminates, and there exists an n in this chain such that q_n is irreducible. Therefore, R is a factorization domain.

Now we want to prove the uniqueness. That is, if $p_1 \dots p_n = q_1 \dots q_m$ for irreducibles p_i, q_j and $n \leq m$ WLOG, then we want to show that $m = n$ and that $p_i = u_i q_i$ with units u_i up to reordering for all i . We do so by induction on n .

¹The proof that this is an ideal is as follows:

We first prove that $\bigcup_{n=1}^{\infty} I_n$ is a subgroup of R under addition. Let $r, s \in \bigcup_{n=1}^{\infty} I_n$, where $r \in I_k$ and $s \in I_{k+i}$ for some $k, i \in \mathbb{N}$. Since $I_1 \subseteq I_2 \subseteq \dots$ are ideals of R , we know that $r \in I_k$ implies that $r \in I_{k+i}$. Thus, $r - s \in I_{k+i}$ due to I_{k+i} being an ideal. As $I_{k+i} \subseteq \bigcup_{n=1}^{\infty} I_n$, we have $r - s \in \bigcup_{n=1}^{\infty} I_n$, which means that $\bigcup_{n=1}^{\infty} I_n$ is closed under additive inverse. Hence, $\bigcup_{n=1}^{\infty} I_n$ is a subgroup of R under addition.

Then, we prove that for any $t \in R, r \in \bigcup_{n=1}^{\infty} I_n$, we would have $tr, rt \in \bigcup_{n=1}^{\infty} I_n$. Since $r \in \bigcup_{n=1}^{\infty} I_n$, it must be true that $r \in I_k$ for some $k \in \mathbb{N}$. Hence, $tr, rt \in I_k$ due to I_k being an ideal. Therefore, $tr, rt \in \bigcup_{n=1}^{\infty} I_n$ for any $t \in R, r \in \bigcup_{n=1}^{\infty} I_n$.


In conclusion, since $\bigcup_{n=1}^{\infty} I_n$ is a subgroup of R under addition with the property that $tr, rt \in \bigcup_{n=1}^{\infty} I_n$ for any $t \in R, r \in \bigcup_{n=1}^{\infty} I_n$, it is an ideal of R . \square

(Base case) If $p_1 = q_1 \dots q_m$ and p_1 irreducible, then $q_2 \dots q_m$ are all units. Hence, $m = 1$ and $p_1 = q_1$.

(Inductive step) Say we have already proven the statement for $n = k$. Then consider $p_1 p_2 \dots p_{k+1} = q_1 q_2 \dots q_m$. Since R is a PID where irreducible implies prime, p_1 is a prime element dividing the product of primes $q_1 q_2 \dots q_m$, so we say WLOG $p_1 | q_1$. This means that $q_1 = u_1 p_1$ for some $u \in R$, but since q_1 is not reducible, it forces u_1 to be a unit. Hence, we apply cancellation on both sides and get $p_2 \dots p_{k+1} = (u_1 q_2) \dots q_m$.

By inductive hypothesis, $m - 1 = k$ and p_i, q_i are associates up to reordering for any i . Hence, the factorization must be unique. \square

We have shown:



Field extensions

We observe that the polynomial $x^2 - 2 \in \mathbb{Q}[x]$ is irreducible. If we have $x^2 - 2 = p(x)q(x)$ where p, q nonunits, then $\deg(p) + \deg(q) = 2$ and we cannot have any $0+2$ combinations due to constants being units, we only have $x^2 - 2 = (x - \sqrt{2})(x + \sqrt{2})$, but $x \pm \sqrt{2} \notin \mathbb{Q}[x]$!

Since $\mathbb{Q}[x]$ is a UFD, the irreducible element $(x^2 - 2)$ is prime, and since $\mathbb{Q}[x]$ is a PID, $(x^2 - 2)$ is maximal which means that $\mathbb{Q}[x]/(x^2 - 2)$ is a field.

Phase II plan: Field extensions!

Suppose F is a field and $p(x) \in F[x]$ nonzero. Recall that $F[x]$ is a ED with the norm function $\deg(a(x))$ and long division of polynomials. Let $a(x) + (p(x)) \in F[x]/(p(x))$. By the division algorithm, we have $a(x) = p(x)q(x) + r(x)$ for $q(x), r(x) \in F[x]$ and $\deg(r(x)) < \deg(p(x))$ or $r(x)$ is the zero polynomial.

← $a(x)$ is a coset rep
← We can do division algorithm since this is an ED

Now we see that since $a(x) - r(x) \in (p(x))$, they are in the same coset! Hence $a(x) + (p(x)) = r(x) + (p(x))$. We observe that every element of $F[x]/(p(x))$ can be represented by a polynomial of a degree less than $\deg(p(x))$. In other words, if $\deg(p(x)) = n$, then $F[x]/(p(x))$ is of the form

← The expression under the bar functions like $r(x)$! Also note that span is just the set of linear combinations.

$$\begin{aligned} F[x]/(p(x)) &= \left\{ \overline{a_0 + a_1x + \cdots + a_{n-1}x^{n-1}} \mid a_0, a_1, \dots, a_{n-1} \in F \right\} \\ &= \text{Span}_F\{\bar{1}, \bar{x}, \dots, \bar{x}^{n-1}\} \end{aligned}$$

In fact, $F[x]/(p(x))$ is (partly) just a **vector space** over F ...

← Why is this not the vector space over $F/(p(x))$ but just F ? See the next paragraph.

We shall observe that it does not matter if we are using F or \bar{F} .

Consider $\varphi : F \hookrightarrow F[x]/(p(x))$ where $a \mapsto \bar{a}$. We observe this is an **injective** map: whenever $\deg(p(x)) = n > 0$, we have $\varphi(a) = \varphi(b)$ if and only if $\bar{a} = \bar{b}$, which happens if and only if $a - b \in (p(x))$; but the difference of two constants always have $\deg 0$ and cannot be in $(p(x))$ unless it is a straight zero, which tells us that $\bar{a} = \bar{b}$ if and only if $a = b$. In other words, $F[x]/(p(x))$ contains an isomorphic copy of F , its field of scalars! Namely, $\overline{F \cong \varphi(F) = \{\bar{a} \in F[x]/(p(x)) \mid a \in F\}}$.

...Hence, $F[x]/(p(x))$ is a vector space **of dimension n** over the scalar field F that also contains an isomorphic copy of F .

Moreover, if $p(x)$ is irreducible, then $(p(x))$ is prime since this is an ED, and hence, it is also a maximal ideal, meaning that $F[x]/(p(x))$ is a field containing an isomorphic copy of F .

← all thanks to Euclidean domains!

Definition 19. Suppose $F \subseteq K$ are fields. Then K is called a **field extension** of F .

- Notation: K/F or $\begin{smallmatrix} K \\ \mid \\ F \end{smallmatrix}$ (the lattice notation)

← Please, this is NOT a quotient. DO NOT CONFUSE THOSE!!

The dimension of K as a vector space over F is called the **degree** of the extension.

- Notation: $[K : F]$

But does my field F always have an extension? Here is a systematic way to get extensions:

Example 21. If $p(x) \in F[x]$ is an irreducible polynomial of degree $n \geq 1$ over the field F , then $F[x]/(p(x))$ is a **field extension** of F of degree n .

← Since $\varphi(F) \cong F$, and $\varphi(F) \subseteq F[x]/(p(x))$

Furthermore, if $p(x) = a_0 + a_1x + \cdots + a_nx^n$, then \bar{x} is a **root** of

$$\varphi(p(x)) = \bar{a}_0 + \bar{a}_1y + \cdots + \bar{a}_ny^n \in (F[x]/(p(x)))[y]$$

because, plugging in $y = \bar{x}$, we get

$$\bar{a}_0 + \bar{a}_1\bar{x} + \cdots + \bar{a}_n\bar{x}^n = \overline{p(x)} = \bar{0} \in F[x]/(p(x))$$

Hence, the isomorphic copy of the polynomial $p(x)$ has **roots** in the field extension $F[x]/(p(x))$.

← We think about modding out by $(p(x))$ as making it equal to zero, which is how we find roots.

So, what the hell is $F[x]/(p(x))$? We have already shown that the field extension $F[x]/(p(x))$ does indeed contain a root of $p(x)$. Now we think about it **the other way around**: if we want to find an extension of F that contains a root of $p(x)$, we would eventually get this one!

Suppose $p(x) \in F[x]$ is irreducible. Let K/F be an extension, and $\alpha \in K$ a root of $p(x)$. Denote by $F(\alpha) \subseteq K$ the **smallest** subfield of K that contains both F and α . Consider the map $\varphi : F[x] \rightarrow F(\alpha) \subseteq K$ where $q(x) \mapsto q(\alpha)$ is simply the evaluation at α map. We note that $p(x) \in \ker(\varphi) = (d(x))$ since an ED is a PID; this implies that $p(x) = u(x)d(x)$. As $p(x)$ is irreducible, $u(x)$ must be a unit, which means $p(x)$ and $d(x)$ are associates and $\ker(\varphi) = (p(x))$. Therefore,

$$F[x]/(p(x)) = F[x]/\ker(\varphi) \cong \varphi(F[x]) \subseteq F(\alpha)$$

by first isomorphism theorem. However, $F(\alpha) \subseteq K$ the **smallest** subfield of K that contains both F and α , so $\varphi(F[x])$ cannot be smaller than that. Hence, it must be true that $\varphi(F[x]) = F(\alpha)$.

← Observe that $\varphi(F[x])$ is a field: $\ker(\varphi)$ is a maximal ideal

Therefore, $F(\alpha)$ is simply $F[x]/(p(x))$. □

To summarize so far!

Suppose $p(x) \in F[x]$ is an irreducible polynomial with coefficients in the field F .

- $F[x]/p(x)$ is a **field** containing an isomorphic copy of F in which $\bar{x} = x + (p(x))$ is a **root** of (the image of) $p(y) \in (F[x]/(p(x)))[y]$.

Example 22. In $\mathbb{Q}[x]/(x^2 - 2)$, we have $x + (x^2 - 2)$ is a root of $y^2 - \bar{2} \in (\mathbb{Q}[x]/(x^2 - 2))[y]$ because

$$\begin{aligned} & (x + (x^2 - 2))^2 - (2 + (x^2 - 2)) \\ &= x^2 - 2 + (x^2 - 2) && \text{by coset addition \& multiplication} \\ &= 0 + (x^2 - 2) && \text{since } x^2 - 2 \in (x^2 - 2) \\ &= \bar{0} \end{aligned}$$

Furthermore, if $\deg(p(x)) = n$, then

$$F[x]/(p(x)) = \left\{ \overline{a_0 + a_1x + \cdots + a_{n-1}x^{n-1}} \mid a_0, a_1, \dots, a_{n-1} \in F \right\}$$

is a vector space over F of dimension n .

Example 23. $\mathbb{Q}[x]/(x^2 - 2) = \{\bar{a}_0 + \bar{a}_1\bar{x} \mid a_0, a_1 \in \mathbb{Q}\} = \text{Span}_{\mathbb{Q}}\{\bar{1}, \bar{x}\}$

- If K/F is an extension and $\alpha \in K$ is a root of $p(x)$, denote by $F(\alpha)$ **the smallest field containing F and α** .

← Read ‘ F adjoint α ’




Figure 1: Field diagram

Then $F(\alpha) \cong F[x]/(p(x))$, and

$$\begin{aligned} F(\alpha) &= \left\{ \overline{a_0 + a_1x + \cdots + a_{n-1}x^{n-1}} \mid a_0, a_1, \dots, a_{n-1} \in F \right\} \\ &= F[\alpha] \quad \leftarrow \text{the polynomial of } \alpha \text{ over } F \end{aligned}$$

← The eval map
 $\varphi : F[x] \rightarrow F(\alpha)$
 where $f(x) \mapsto f(\alpha)$
 has in fact
 $\ker(\varphi) = (p(x))$
 when α is a root of
 $p(x)$.

Example 24. $\mathbb{Q}(\sqrt{2}) = \{a_0 + a_1\sqrt{2} \mid a_0, a_1 \in \mathbb{Q}\} = \mathbb{Q}[\sqrt{2}]$

Irreducibility – a survey

Proposition 21. If $p(x) \in F[x]$, then $\alpha \in F$ is a root *if and only if* $x - \alpha$ divides $p(x)$.

Proof. Write $p(x) = (x - \alpha)q(x) + r(x)$ with $q(x), r(x) \in F[x]$ and $\deg(r(x)) = 0$ or $r(x) = 0$. Then $0 = p(\alpha) = 0 + r(\alpha)$ which forces $r(x) = 0$. \square

Corollary 22. A degree-2 or -3 polynomial over a field F is irreducible *if and only if* it has no roots in F .

Proposition 23. Suppose $p(x) = a_0 + a_1x + \cdots + a_nx^n \in \mathbb{Z}[x]$ with root $\frac{c}{d}$ written in reduced form (i.e. $\gcd(c, d) = 1$). Then $\boxed{c|a_0 \text{ and } d|a_n}$.

Proof.

$$\begin{aligned} d^n \cdot p\left(\frac{c}{d}\right) &= 0 \\ 0 &= (a_0d^n + a_1d^{n-1}c + \cdots + a_{n-1}dc^{n-1}) + a_nc^n \end{aligned}$$

$$0 = a_0 d^n + (a_1 d^{n-1} c + \cdots + a_{n-1} d c^{n-1} + a_n c^n)$$

Looking at the 2nd line, since d divides all of the ones in the $()$, it must also divide the last term $a_n c^n$. However, since $\gcd(c, d) = 1$, it forces d to divide a_n .

Similarly, we make the same argument for c and a_0 using the 3rd line. □

Lemma 24. $(R/I)[x] \cong R[x]/(I)$ where $(I) = I[x]$.

Proof. Consider the surjective homomorphism $\pi : R[x] \rightarrow (R/I)[x]$. □

Proposition 25 (Eisenstein's Criterion). Suppose $f(x) = 1x^n + a_{n-1}x^{n-1} + \cdots + a_1x + a_0 \in \mathbb{Z}[x]$ is a monic polynomial and $p \in \mathbb{Z}$ is a **prime** such that $p \mid a_0, \dots, a_{n-1}$ but $p^2 \nmid a_0$. Then $f(x)$ is irreducible.

Proof. Assume BWOC that $f(x) = a(x)b(x)$ for some nonunit $a(x), b(x) \in \mathbb{Z}[x]$, then

$$x^n = \bar{f}(x) = \bar{a}(x)\bar{b}(x)$$

in $(\mathbb{Z}/p\mathbb{Z})[x] \cong \mathbb{Z}[x]/p\mathbb{Z}[x]$ since all other terms are divisible by p . Since $\mathbb{Z}/p\mathbb{Z}$ does not contain any zero divisors, $\bar{a}(x), \bar{b}(x)$ must have zero constant terms. Hence $a(x), b(x)$ have constant terms that are multiples of p , so $a(x)b(x)$ have constant term divisible by p^2 . This is a contradiction with $p^2 \nmid a_0$. □

Lemma 26 (Gauss' Lemma). If $p(x) \in \mathbb{Z}[x]$ is reducible in $\mathbb{Q}[x]$, then it is reducible in $\mathbb{Z}[x]$.

Proof. Suppose $p(x) = a(x)b(x)$ for $a(x), b(x) \in \mathbb{Q}[x]$. Then by multiplying by coefficient denominators, for some $m \in \mathbb{Z}$, we could write $m \cdot p(x) = c(x)d(x)$ for some $c(x), d(x) \in \mathbb{Z}[x]$. Now since $m \in \mathbb{Z}$, we could write $m = q_1 q_2 \cdots q_n$ be a product of irreducibles in \mathbb{Z} .

Now in $(\mathbb{Z}/q_1\mathbb{Z})[x] \cong \mathbb{Z}[x]/(q_1\mathbb{Z})[x]$, we observe that $m \cdot p(x) = c(x)d(x) = q_1(q_2 \cdots q_n)p(x)$, meaning that

$$\overline{c(x)} \overline{d(x)} = \overline{q_1(q_2 \cdots q_n)p(x)} = \bar{0}$$

Since $(\mathbb{Z}/q_1\mathbb{Z})[x] \cong \mathbb{Z}[x]/(q_1\mathbb{Z})[x]$ is an integral domain, WLOG, $\overline{c(x)} = \bar{0}$ if and only if $c(x) \in q_1\mathbb{Z}[x]$, meaning that all coefficients of $c(x)$ are multiples of q_1 . Therefore, $\frac{1}{q_1}c(x) \in \mathbb{Z}[x]$.

← since q_1 is irreducible and hence prime in UFD

Now we repeat the process for all q_1, q_2, \dots, q_n and we are done. □

Recall that if $F \subseteq K$ are fields, $\alpha \in K$ and $p(x) \in F[x]$ is irreducible with root α , then

$$F[\alpha] = F(\alpha) \cong F[x]/(p(x)) = \overline{\{a_0 + a_1x + \cdots + a_{n-1}x^{n-1} \mid a_0, a_1, \dots, a_{n-1} \in F\}}$$

We observe that this has a few implications. For instance, $F(\alpha)$ contains $\frac{1}{\alpha}$, meaning that it could also be written as a polynomial of α with coefficients in F (as in $F[\alpha]$)!

← since it is a field containing the mult. inverse of α

Definition 20. Suppose K/F is a field extension and $\alpha \in K$. We say that α is **algebraic over F** if there exists $p(x) \in F[x]$ such that $p(\alpha) = 0$. If not, α is **transcendental**.

Definition 21. The extension K/F is an **algebraic extension** if **every** element $\alpha \in K$ is algebraic over F .

Example 25. π is transcendental over \mathbb{Q} but algebraic over \mathbb{R} (since it is a root of $x - \pi$).

Proposition 27. If K/F is a **finite extension**, then it is an algebraic extension.

← finite extension just means finite degree
 $[K : F] < \infty$

Proof. Call $[K : F] = n$ and let $\alpha \in K$. Then the $n + 1$ elements $\{1, \alpha, \alpha^2, \dots, \alpha^n\}$ must be linearly dependent. Hence, by linear algebra, there exist $a_0, a_1, \dots, a_n \in F$ not all zero such that the linear combination $a_0 + a_1\alpha + a_2\alpha^2 + \cdots + a_n\alpha^n = 0$. Hence, α is a root of $a_0 + a_1x + a_2x^2 + \cdots + a_nx^n = 0$. \square

← since $n + 1 > \dim(K/F) = n$

Corollary 28. If K/F is an extension and $\alpha \in K$, then α is algebraic over F if and only if $[F(\alpha) : F] < \infty$.

Proof.

(\Leftarrow) Follows from prop.

(\Rightarrow) If α is algebraic, then there exists an irreducible polynomial $p(x)$ with α as a root and of degree $n < \infty$. Then $F(\alpha) \cong F[x]/(p(x))$ is a n -dimensional vector space over F .

Another perspective: $F(\alpha) = \text{Span}_F\{1, \alpha, \alpha^2, \dots, \alpha^{n-1}\}$.

← Review proof of $F(\alpha) \cong F[x]/(p(x))$.

\square

Proposition 29. Suppose K/F is an extension & $\alpha \in K$ is algebraic over F . Then there exists a unique, irreducible, and monic polynomial $m_{\alpha, F}(x) \in F[x]$ that has α as a root.

Remark. We observe that m does depend on the base field F ; $m_{\sqrt{2}, \mathbb{Q}}(x) = x^2 - 2$, but $m_{\sqrt{2}, \mathbb{Q}(\sqrt{2})}(x) = x^2 - \sqrt{2}$.

Proof. Since the subset of $F[x]$ satisfying α is a root is nonempty, we can pick one with a **minimal degree**. By multiplying by an element of F if necessary, we can assume WLOG this polynomial is **monic**. Call it $m_{\alpha,F}(x)$.

Assume BWOC that m is the product of two other polynomials of lesser degree such that $m_{\alpha,F}(x) = a(x)b(x)$, then we plug in $0 = m_{\alpha,F}(\alpha) = a(\alpha)b(\alpha)$. Since there are no zero divisors in $F[x]$, WLOG $a(\alpha) = 0$, contradicting the minimality of $m_{\alpha,F}$. Hence $m_{\alpha,F}$ is **irreducible**.

← So $F(\alpha) \cong F[x]/(m_{\alpha,F}(x))$

Then, BWOC if $p(x) \in F[x]$ with α as a root and is monic and irreducible, there exist $q(x), r(x) \in F[x]$ such that $p(x) = m_{\alpha,F}(x)q(x) + r(x)$ where $\deg(r) < \deg(m_{\alpha,F})$ or $r(x) = 0$. Then, we observe that $p(\alpha) = 0 = m_{\alpha,F}(\alpha)q(\alpha) + r(\alpha) = 0 + r(\alpha)$. Thus, $r(\alpha) = 0$, so $\deg(r) \geq \deg(m_{\alpha,F})$ unless $r(x) = 0$ by minimality. Hence we must have $r(x) = 0$, so $m_{\alpha,F} | p$. This contradicts the assumption that p is monic and irreducible. Therefore, $m_{\alpha,F}$ is the **only** minimal, monic and irreducible polynomial where α is a root. \square

Definition 22. $m_{\alpha,F}(x)$ is the **minimal** polynomial of α over F .

(The following is kind of on a tangent)

Some exam prep!

- In general, for subrings $R \subseteq S$, we have if $r \in R^\times$, then $r \in S^\times$.
- If we adjoin one root of an irreducible polynomial to a field, the fields are isomorphic no matter which root of that polynomial we adjoin.

(Tangent ends here)

To summarize, if K/F is a field extension and $\alpha \in K$, then α is **algebraic** over F if it is the root of some polynomials in $F[x]$. For each algebraic α , there exists a unique, monic, irreducible polynomial $m_{\alpha,F}(x) \in F[x]$ such that $m(\alpha) = 0$. In that case, the degree of extension $[F(\alpha) : F] = \deg(m_{\alpha,F}(x))$; and, if $p(\alpha) = 0$ for some $p(x) \in F[x]$, then $m_{\alpha,F} | p(x)$. In general, if $[K : F] < \infty$, then K/F is algebraic. Thus, $[F(\alpha) : F] < \infty$ if and only if α is algebraic over F .

Proposition 30. If $F \subseteq K \subseteq L$ are fields, then

$$[L : F] = [L : K] \cdot [K : F]$$

← $mn \begin{pmatrix} L \\ K \\ F \end{pmatrix} \begin{matrix} n \\ m \end{matrix}$

Proof. We first see that if $[K : F] = \infty$, then for any $N \in \mathbb{N}$, there exists $\alpha_1, \dots, \alpha_N \in K$ that are linearly independent over F . In that case, it is certainly true that $\alpha_1, \dots, \alpha_N \in L$ are linearly independent over F . Thus, $[L : F] = \infty$.

If $[L : K] = \infty$, then for any $N \in \mathbb{N}$, there exists $\beta_1, \dots, \beta_N \in L$ that are linearly independent over K . As a result, it also is linearly independent over F . Hence, $[L : F] = \infty$.

If $[K : F] = m$ and $[L : K] = n$, let $\alpha_1, \dots, \alpha_m \in K$ be a basis for K over F and $\beta_1, \dots, \beta_n \in L$ be a basis for L over K .

Claim: $\{\alpha_i \beta_j \mid 1 \leq i \leq m, 1 \leq j \leq n\}$ forms a basis for L over F . □

Some nice consequences:

Corollary 31. Suppose K/F is an extension and $\alpha, \beta \in K$ are algebraic over F . Then:

- $F(\alpha, \beta) = (F(\alpha)(\beta))$
- $[F(\alpha, \beta) : F] = [F(\alpha, \beta) : F(\beta)][F(\beta) : F] = \deg(m_{\alpha, F(\beta)}(x)) \cdot \deg(m_{\beta, F}(x))$.
However, note that the minimal polynomial

$$m_{\alpha, F(\beta)}(x) \mid m_{\alpha, F}(x) \in F(\beta)[x]$$

so $\deg(m_{\alpha, F(\beta)}(x)) \leq \deg(m_{\alpha, F}(x))$. Hence,

$$[F(\alpha, \beta) : F] \leq \deg(m_{\alpha, F}(x)) \deg(m_{\beta, F}(x)) < \infty$$

This means that whenever α, β are algebraic over F , we get that $F(\alpha, \beta)/F$ is an algebraic extension.

- As a result, $\alpha \pm \beta, \alpha\beta, \alpha/\beta$ are all algebraic over F . The algebraic elements hence form a **field**.

Proposition 32. Suppose K/F is an extension. Then $[K : F] < \infty$ if and only if $K = F(\alpha_1, \dots, \alpha_n)$ could be written where $\alpha_1, \dots, \alpha_n \in K$ are algebraic over F .

In other words, an extension is finite if and only if it is generated by adjoining a finite amount of algebraic elements.

Proof.

(\implies) If $[K : F] < \infty$, then suppose $\{\alpha_1, \dots, \alpha_n\}$ is a basis of K over F . Then $\alpha_1, \dots, \alpha_n$ are algebraic and every element of K is an F -linear combination of α_i s. Hence K must be the smallest field containing F and α_i s, which means $K = F(\alpha_1, \dots, \alpha_n)$.

← Linear independence implies that whenever $a_1\alpha_1 + a_2\alpha_2 + \dots + a_N\alpha_N = 0$ for some coefficients $a_1, \dots, a_N \in F$, then necessarily $a_1 = a_2 = \dots = a_N = 0$.

← Use linear combinations to prove this claim.

← the smallest subfield of K containing F, α, β
← since $p(\alpha) = 0 \iff m_{\alpha, F}(x) \mid p(x)$

(\Leftarrow) We observe that

$$\begin{aligned} [K : F] &= [(F(\alpha_1, \dots, \alpha_{n-1}))(\alpha_n) : F(\alpha_1, \dots, \alpha_{n-1})] \cdot \dots \cdot [F(\alpha_1) : F] \\ &\leq \prod_{i=1}^n \deg(m_{\alpha_i, F}(x)) < \infty \end{aligned}$$

□

Corollary 33. If L/K and K/F are algebraic extensions, then so is L/F .



← L/K and K/F need not be finite!

Proof. Suppose $\alpha \in L$. Since L/K is algebraic, there exists $p(x) \in K[x]$ such that $p(\alpha) = 0$. Let $\alpha_0, \dots, \alpha_n \in K$ be the coefficients of $p(x)$, necessarily algebraic over F since K/F algebraic. Therefore,

$$[F(a_0, \dots, a_n, \alpha) : F] = [(F(a_0, \dots, a_n))(\alpha) : F(a_0, \dots, a_n)][F(a_0, \dots, a_n) : F]$$

Since $p(\alpha) = 0$ has coefficients in $K \supseteq F(\alpha_0, \dots, \alpha_n)$, we have $[(F(a_0, \dots, a_n))(\alpha) : F(a_0, \dots, a_n)] < \infty$. The second term is also clearly $< \infty$. Therefore, $[F(a_0, \dots, a_n, \alpha) : F] < \infty$, meaning that α is algebraic over F . □

Definition 23. Suppose L/F is an extension & K_1 and K_2 are intermediate fields. The **composite** field $K_1 K_2$ is the smallest subfield of L containing K_1 and K_2 .



Definition 24. Suppose F is a field and $p(x) \in F[x]$. The **splitting field** of $p(x)$ over F is the smallest field extension of F over which $p(x)$ could be factored into **linear factors**.

Remark. If E is the splitting field of $p(x)$ over F then $[E : F] \leq n!$ where $n = \deg(p(x))$.

Remark. Such an extension is called **normal**.

Proposition 34. Splitting fields exist.

← Assuming that splitting fields **exist** and are **unique** up to isomorphism.

Proof outline. By induction on $\deg(p(x))$, whose base case, $\deg(p(x)) = 1$, yields F as a splitting field. More generally, any $p(x)$ has a root α in $F(\alpha) \cong F[x]/(q(x))$ for some irreducible $q(x)$ so $p(x) = (x - \alpha)f(x) \in F(\alpha)[x]$. We observe that $\deg(f(x)) = \deg(p(x)) - 1$. Induction takes care of the rest. □

← The splitting field of $p(x)$ over F is the same as the splitting field of $f(x)$ over $F(\alpha)$

Remark. K is a splitting field over F if and only if every irreducible $p(x) \in F[x]$ that has one root in K has **all** its roots in K .

Non-example 26. $\mathbb{Q}(\sqrt[3]{2})$ over \mathbb{Q} is not such an extension.

Lemma 35. Suppose $\varphi : F_1 \rightarrow F_2$ is a field isomorphism, $p_1(x) \in F_1[x]$, and $p_2(x) = \varphi(p_1(x))$ (φ applied to coeffs of $p_1(x)$). Let α_1 be a root of the irreducible factor $q_1(x)$ of $p_1(x)$, and let $q_2(x) = \varphi(q_1(x))$ and α_2 be a root of $q_2(x)$. Then there exists an isomorphism $\tau : F_1(\alpha_1) \rightarrow F_2(\alpha_2)$ such that $\tau(\alpha_1) = \alpha_2$ and $\tau|_{F_1} = \varphi$ (this means “ τ restricted to F_1 ”).

← In this way, φ induce a ring isomorphism $F_1[x] \rightarrow F_2[x]$.

Proof outline.

$$\begin{array}{ccccccc} F_1(\alpha_1) & \xrightarrow{\sim} & F_1[x]/(q_1(x)) & \xrightarrow{\sim} & F_2[x]/(q_2(x)) & \xrightarrow{\sim} & F_2(\alpha_2) \\ \alpha_1 & \xrightarrow{\quad} & \bar{x} & \xrightarrow{\quad} & \bar{x} & \xrightarrow{\quad} & \alpha_2 \\ \text{if } a \in F_1, & a & \xrightarrow{\quad} & \bar{a} & \xrightarrow{\quad} & \overline{\varphi(a)} & \xrightarrow{\quad} & \varphi(a) \end{array}$$

□

Proposition 36. Suppose $F_1, F_2, \varphi, p_1(x)$ and $p_2(x)$ are as in Lemma 35. Let E_1 & E_2 be splitting fields of p_1 and p_2 respectively. Then there exists an isomorphism $\sigma : E_1 \rightarrow E_2$ such that $\sigma|_{F_1} = \varphi$.

← if we set $F_1 = F_2$ and $p_1 = p_2$, we get corollary: splitting fields are unique up to isomorphism.

Proof. Proceed by induction on $\deg(p_1(x))$. For the base case, if $\deg(p_1(x)) = 1$, then $E_1 = F_1$ and $\sigma = \varphi$.

Assume the result is true for all polynomials of fixed degree $k \geq 1$ and suppose $\deg(p_1(x)) = k+1$. Let α_1 be a root of $p_1(x)$ and α_2 be a root of the φ -corresponding irreducible factor of $p_2(x)$. By Lemma 35, φ can be extended to $\tau : F_1(\alpha_1) \rightarrow F_2(\alpha_2)$ such that $\tau|_{F_1} = \varphi$.

In $(F_1(\alpha_1))[x]$, we can factor out $p_1(x) = (x - \alpha_1)g_1(x)$, and in $(F_2(\alpha_2))[x]$ we factor $p_2(x) = (x - \alpha_2)g_2(x)$ with $g_2(x) = \tau(g_1(x))$. We observe that E_1 and E_2 are the splitting fields of g_1 and g_2 over $F_1(\alpha_1)$ and $F_2(\alpha_2)$!

←

$$\begin{array}{ccc} \sigma : & E_1 & \xrightarrow{\sim} E_2 \\ | & \downarrow & \downarrow \\ \tau : & F_1(\alpha_1) & \xrightarrow{\sim} F_2(\alpha_2) \\ | & \downarrow & \downarrow \\ \varphi : & F_1 & \xrightarrow{\sim} F_2 \end{array}$$

By inductive hypothesis, τ could be extended to σ and $\sigma|_{F_1(\alpha_1)} = \tau$ and $\sigma|_{F_1} = \varphi$. □

Corollary 37. Splitting fields are unique.

Proof. Set $F_1 = F_2$, $\varphi = \text{id}$, $p_1(x) = p_2(x)$. □

(The following is kind of on a tangent)

Homework hint: the proof of existence & uniqueness of splitting fields relied on inductive arguments where we adjoin one root at a time. This is the same as saying $E = F(\alpha_1, \dots, \alpha_n)$ but this tends to overlook isomorphic ways to adjoin roots. In this context, it is convenient to start by considering a specific K containing F and all roots of $p(x)$. In that case, $E = F(\alpha_1, \dots, \alpha_n)$ becomes more rigorous.

(Tangent ends here)

Definition 25. A polynomial is called **separable** if it doesn't have repeated roots.

← First note that a poly of degree n over a field has exactly n roots.

Definition 26. Suppose $f(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0$. The **formal derivative** of $f(x)$ is the polynomial

$$D_x f(x) = f'(x) = n a_n x^{n-1} + (n-1) a_{n-1} x^{n-2} + \dots + a_1$$

From this definition, we can check that the usual differential rules hold.

Lemma 38. Suppose F is a field, $f(x)$ is a polynomial in $F[x]$, and E/F a field extension containing a root α of $f(x)$. Then α is a repeated root of $f(x)$ if and only if α is a root of the formal derivative $f'(x)$.

Proof. If α is a repeated root of $f(x)$ then $f(x) = (x - \alpha)^2 g(x)$ for some $g(x) \in E[x]$. In that case, $f'(x) = 2(x - \alpha)g(x) + (x - \alpha)^2 g'(x)$ and so $f'(\alpha) = 0$.

Conversely, if $f'(\alpha) = 0$, then differentiating $f(x) = (x - \alpha)h(x)$ (where $h(x) \in E[x]$) and plugging $x = \alpha$ yields $0 = f'(\alpha) = h(\alpha) + (\alpha - \alpha)h'(\alpha) = h(\alpha)$. This is saying that $h(x) = (x - \alpha)g(x)$ for some $g(x) \in E[x]$. \square

Lemma 39. If $f(x) \in F[x]$ is **irreducible and not separable**, then $f'(x) = 0$.

finish proof

\square

If $f(x)$ is not constant and $f'(x) = 0$, then $\text{char}(F) = p > 0$ and $f(x) = g(x^p)$.

← All powers in $f(x)$ are multiples of $p(x)$.

Proposition 40. If $\text{char}(F) = 0$, or $|F| < \infty$ and $\text{char}(F) = p$, then every irreducible polynomial in $F[x]$ is separable.

Proof. For the case of $\text{char}=0$, it follows from 39.

For the case of $\text{char}>0$, suppose F is a finite field of p^n elements². Then the map $F \rightarrow F$ where $\alpha \mapsto \alpha^p$ is a field isomorphism. Hence, every element of F is a p^{th} power.

← Use binomial theorem.

← Such fields are called **perfect**.

Now suppose BWOC $f(x) = \sum_{i=0}^n a_i x^i \in F[x]$ is an irreducible but not separable polynomial. Therefore, $f'(x) = 0$ must be true. This happens *if and only if* $f(x) = \sum_{j=0}^m a_{jp} x^{jp}$, that is, the x in all terms are of p^{th} degree. However, we know that all elements $a_{jp} \in F$ are already the p^{th} powers of sth else $(b_{jp})^p = a_{jp}$, so

$$f(x) = \sum_{j=0}^m (b_{jp}^p) x^{jp}$$

and by reverse Binomial Theorem, we get

$$f(x) = \sum_{j=0}^m (b_{jp}^p) x^{jp} = \left(\sum_{j=0}^m b_{jp} x^j \right)^p$$

is not irreducible! □

Non-example 27. Let $F = \mathbb{F}_p(t) = \left\{ \frac{f(t)}{p(t)} \mid f(t), g(t) \in \mathbb{F}_p[t], g(t) \neq 0 \right\}$.

← This is a field of $\text{char}>0$ but is infinite.

Then $p(x) = x^p - t$ is not separable (but it is irreducible). This can be seen if we suppose α is a root of $p(x)$ (so $\alpha = t$). Then, in $F(\alpha)[x]$, we have $p(x) = x^p - t = x^p - \alpha^p = (x - \alpha)^p$, which tells us $p(x)$ is not separable.

← The coefficients of $p(x)$ are ratios of polys in $\mathbb{F}_p(t)$.

(The following is kind of on a tangent)

Prime fields

Suppose R is a commutative ring with identity. The map $\mathbb{Z} \rightarrow R$ where $n \mapsto \underbrace{\pm(1_R + 1_R + \cdots + 1_R)}_{|n| \text{ times}}$ ($-$ if $n < 0$) is a ring homomorphism with kernel $n\mathbb{Z}$ where $n = \text{char}(R)$. So:

← check it!

- if $\text{char}(R) = 0$, then R contains \mathbb{Z} ;
- if $\text{char}(R) = n > 0$, then R contains $\mathbb{Z}/n\mathbb{Z}$.

²See Section 27

If F is a field, then:

- if $\text{char}(F) = 0$, then F contains \mathbb{Q} ;
- if $\text{char}(F) = p > 0$, then p prime and F contains $\mathbb{Z}/p\mathbb{Z}$.

← That is, F is an extension of $\mathbb{Z}/p\mathbb{Z}$!

In other words, every field is an extension of \mathbb{Q} or \mathbb{F}_p . Moreover, a finite field is a finite extension of \mathbb{F}_p : if $[F : \mathbb{F}_p] = n$, then $|F| = p^n$.

In addition, $|F - \{0\}| = p^n - 1 \implies$ if $\alpha \in F - \{0\}$ then $\alpha^{p^n-1} = 1$, implying that if $\alpha \in F$, then $\alpha^{p^n} = \alpha$, meaning that α is a root of $x^{p^n} - x \in F[x]$. Therefore, F is the splitting field of $x^{p^n} - x$. But splitting fields are unique, so we conclude that there is only one unique finite field for each order.

← By Lagrange's Theorem

(Tangent ends here)

Definition 27. An algebraic extension K/F is called (algebraically) **separable** if $m_{\alpha,F}(x)$ is separable for **all** $\alpha \in K$.

Definition 28. A finite extension K/F is called **Galois** if K/F is normal and separable.

← normal just means it is a splitting field of something

Galois Theory


Definition 29. If K/F is an extension, then the **automorphism group** of K/F is defined as

$$\text{Aut}(K/F) = \{\sigma \in \text{Aut}(K) \mid \sigma(a) = a \forall a \in F\}$$

That is, all the automorphisms of K that also fix the field F .

Theorem 41 (The Fundamental Theorem of Galois Theory).

If K/F is Galois, then



where H is a subgroup of $\text{Aut}(K/F)$ that fixes the field E , an extension of F contained in K .

Galois theory is concerned with the study of roots of polynomials by way of automorphisms of splitting fields (of separable polynomials). In particular, we are interested in what

$$\text{Aut}(K) = \{\sigma : K \rightarrow K \text{ isomorphisms}\}$$

(a group under composition) is. Naturally, such groups are *finite*.

← review MATH171 finite groups!

Last time, we showed that $K \supseteq \begin{cases} \mathbb{Q} & \text{if char}(K) = 0 \\ \mathbb{F}_p & \text{if char}(K) = p \end{cases}$, since $\sigma(1) = \sigma(1^2) = (\sigma(1))^2$ implies that $\sigma(1) = 1$ must always be true! Hence, $\sigma(n) = n$ must be true in char 0 fields, or $\sigma(\bar{n}) = \bar{n}$ if char > 0 for all $n \in \mathbb{Z}$. Therefore, σ **fixes** the prime subfield \mathbb{Q} or \mathbb{F}_p .

Remark. Why does Definition 29 have to fix the field F ? Because we've shown that $\text{Aut}(K) = \begin{cases} \text{Aut}(K/\mathbb{Q}) & \text{if char}(K) = 0 \\ \text{Aut}(K/\mathbb{F}_p) & \text{if char}(K) = p \end{cases}$.

Lemma 42. If K/F is an extension, $\alpha \in K$ is algebraic over F and $\sigma \in \text{Aut}(K/F)$, then $\sigma(\alpha)$ is a root of $m_{\alpha,F}(x)$.

Proof. Observe that $m_{\alpha,F}(\alpha) = 0 = \sigma(0) = \sigma(m_{\alpha,F}(\alpha))$. Hence, since $\sigma(\alpha) = \alpha$ for all $\alpha \in F$,

$$\sigma(a_0 + a_1\alpha + \cdots + a_n\alpha^n) = a_0 + a_1\sigma(\alpha) + \cdots + a_n\sigma(\alpha)^n = m_{\alpha,F}(\sigma(\alpha))$$

□

So if $f(x) \in F[x]$, then every $\sigma \in \text{Aut}(K/F)$ **permutes** the roots of $f(x)$ that lie in K . It would be nice if the roots of $f(x)$ all lived in K . This is why we consider K the splitting field of some polynomial over F .

← Recall: if irreducible $f(x)$ has one root in such K , then all roots lie in K .

Proposition 43. If K is the splitting field of some polynomial $f(x)$ over F (so $[K : F] < \infty$), then $|\text{Aut}(K/F)| \leq [K : F]$, with equality if $f(x)$ is separable.

Proof. We will prove a more general statement by induction. If $\sigma : F_1 \rightarrow F_2$ is an isomorphism, $f_1(x) \in F_1[x]$ and $f_2(x) = \sigma(f_1(x)) \in F_2[x]$ and E_1 and E_2 are the splitting fields of f_1 and f_2 over F_1 and F_2 respectively. Then we would like to show that there are at most $[E_1 : F_1]$ isomorphisms $\tau : E_1 \rightarrow E_2$ such that $\tau|_{F_1} = \sigma$ with equality if f_1 separable.

← The prop above fixes $F_1 = F_2$ and σ being identity.

Base case. If $[E_1 : F_1] = [E_2 : F_2] = 1$, then $E_1 = F_1$, $E_2 = F_2$ and $\tau = \sigma$ is our only choice.

Inductive step. Suppose we've proven the result for all extensions of degree $< n$ for some $n \geq 2$. Now consider $[E_1 : F_1] = [E_2 : F_2] = n$. Pick $\alpha \in E_1 \setminus F_1$ and let $\beta \in E_2$ be any root of $\sigma(m_{\alpha, F_1}(x))$. Then σ could be extended to $\rho : F_1(\alpha) \rightarrow F_2(\beta)$ such that $\rho(\alpha) = \beta$ and $\rho|_{F_1} = \sigma$. Observe that $[F_1(\alpha) : F_1] = \deg(m_{\alpha, F_1}(x))$. Moreover, the number of extensions of σ to ρ equals the number of *distinct* roots of $\sigma(m_{\alpha, F_1}(x))$. Thus, the number of extensions of σ to $F_1(\alpha)$ is *at most* the degree of $m_{\alpha, F_1}(x)$ which is $[F_1(\alpha) : F_1]$ with equality if $m_{\alpha, F_1}(x)$ is separable. Since $[E_1 : F_1] = [E_2 : F_2] = n$, we have $[E_1 : F_1(\alpha)] < n$, by inductive hypothesis, there are at most $[E_1 : F_1(\alpha)]$ ways of extending ρ to $\tau : E_1 \rightarrow E_2$. Hence,

$$\begin{aligned} |\{\text{extensions of } \sigma \text{ to } \tau\}| &= |\{\text{extensions of } \sigma \text{ to } \rho\}| |\{\text{extensions of } \rho \text{ to } \tau\}| \\ &\leq [F_1(\alpha) : F_1] [E_1 : F_1(\alpha)] \\ &= [E_1 : F_1] \end{aligned}$$

Looking at the case $F_1 = F_2$, $E_1 = E_2$, $\sigma = \text{id}$, we get our result. \square

Definition 30. If K/F is a normal extension, then the extension is **Galois** if $[K : F] = |\text{Aut}(K/F)|$.

$$\begin{array}{lcl} \leftarrow & & \\ \tau : & E_1 & \xrightarrow{\sim} E_2 \\ & | & | \\ & \vdots & \vdots \\ \rho : & F_1(\alpha) & \xrightarrow{\sim} F_2(\beta) \\ & | & | \\ \sigma : & F_1 & \xrightarrow{\sim} F_2 \end{array}$$