Chapter 3. Classical Demand Theory

3.A. Introduction: Take $\succsim$ as the primitive

(1) What assumption(s) on $\succsim$ do we need to represent $\succsim$ with a utility function?

(2) How to perform utility maximization and derive the demand function?

(3) How to derive the utility as a function of prices and wealth, or the indirect utility function?

(4) How to perform expenditure minimization and derive the expenditure function?

(5) How are demand function, indirect utility function, and expenditure function related?

3.B. Preference Relations: Basic Properties/Assumptions

In the classical approach to consumer demand, the analysis of consumer behavior begins by specifying the consumer's preferences over the commodity bundles in the consumption set. Consumption set is defined as $X \subset \mathbb{R}^L_+$. Consumer's preferences are captured by the preference relation $\succsim$.

Rationality We would assume *Rationality* (*Completeness and Transitivity*) throughout the chapter. Definition 3.B.1 below repeats the formal definition of *Rationality*.

Definition 3.B.1. The preference relation $\succsim$ on X is rational if it possesses the following two properties:

(i) Completeness: For all $x, y \in X$, we have $x \succsim y$ or $y \succsim x$ (or both).

(ii) Transitivity: For all $x, y, z \in X$, if $x \succsim y$ and $y \succsim z$, then $x \succsim z$.

Desirability Assumptions The first *Desirability* Assumption we consider is *Monotonicity*: larger amounts of commodities are preferred to smaller ones.

For Definition 3.B.2, we assume that the consumption of a larger amounts of goods is always feasible in principle; that is, if $x \in X$ and $y \geq x$, then $y \in X$.

Definition 3.B.2. The preference relation $\succsim$ on X is *monotone* if $x, y \in X$ and $y \gg x$ implies $y \succ x$. It is *strongly monotone* if $y \geq x$ & $y \neq x$ implies $y \succ x$.



Figure 1: Monotonicity



Figure 2: Strong Monotonicity

Remark.

- If $\succsim$ is monotone, we may have indifference with respect to an increase in the amount of some but not all commodities.
- If $\succsim$ is strongly monotone, y is strictly preferred to x if y is larger than x for some commodity and is no less for any other commodities.

Claim. If $\succsim$ is strongly monotone, then it is monotone.

Proof. For $x, y \in X$, if $y \gg x$, then $y \geq x$ and $y \neq x$. Since $\succsim$ is strongly monotone, we have $y \succ x$. Thus, $\succsim$ is monotone. $\square$

Example. Here is an example of a preference that is monotone, but not strongly monotone:

$$u(x_1, x_2) = x_1 \text{ in } \mathbb{R}^2_+.$$

1. The preference is monotone: if $y \gg x$, then $y_1 > x_1$. So, we must have $u(y_1, y_2) > u(x_1, x_2)$, which implies $y \succ x$.

2. The preference is not strongly monotone: For $x = (1, 2)$ and $y = (1, 3)$, we have $y \geq x$, but $u(y) = u(x)$, which implies $y \sim x$.

For much of the theory, a weaker desirability assumption, *local nonsatiation*, suffices.

Definition 3.B.3. The preference relation $\succsim$ on X is *locally nonsatiated* if for every $x \in X$ and every $\varepsilon > 0, \exists y \in X$ such that $\|y - x\| \leq \varepsilon$ and $y \succ x$.



Figure 3: Test for Local Nonsatiation

Claim. *Local nonsatiation* is a weaker desirability assumption compared to *monotonicity*. If $\succsim$ is monotone, then it is locally nonsatiated.

Proof. Fix some $\varepsilon > 0$, Let there be an arbitrary $x \in X$ and $e = (1, \cdots, 1)$. For any $\lambda > 0$, we also have $y = x + \lambda e \in X$. Since clearly $y = x + \lambda e \gg x$, by monotonicity, $y \succ x$. On the other hand, $\|y - x\| = \sqrt{L\lambda^2} = \lambda\sqrt{L}$. Thus, for $\lambda < \frac{\varepsilon}{\sqrt{L}}$, $\|y - x\| \leq \varepsilon$. Since x is arbitrary, the existence of the point $y = x + \lambda e$ where $\lambda < \frac{\varepsilon}{\sqrt{L}}$ implies that $\succsim$ is locally nonsatiated. $\square$

Example. Here is an example of a preference that is locally nonsatiated, but not monotone:

$$u(x_1, x_2) = x_1 - |1 - x_2| \text{ in } \mathbb{R}^2_+.$$

1. The preference is locally nonsatiated: Fix an $\varepsilon > 0$. We can find $\lambda > 0$ such that $\lambda < \varepsilon$. Denote $y = (y_1, y_2) = (x_1 + \lambda, x_2)$. Then $u(y) - u(x) = \lambda > 0$, which

implies $y \succ x$. On the other hand, $\|y - x\| = \sqrt{\lambda^2} = \lambda < \varepsilon$. Therefore, $\succsim$ is locally nonsatiated.

2. The preference is not monotone: For $x = (1, 1)$ and $y = (1.5, 2)$, we have $y \gg x$, but $u(x) = 1 > u(y) = 0.5$, which implies $x \succ y$.

Indifference sets Given $\succsim$ and x, we can define 3 related sets of consumption bundles.

1. The indifferent set is $\{y \in X : y \sim x\}$.
2. The upper contour set is $\{y \in X : y \succsim x\}$.
3. The lower contour set is $\{y \in X : x \succsim y\}$.

Implication of local nonsatiation One implication of local nonsatiation (and, hence, of monotonicity) is that it rules out "thick" indifference sets.



Figure 4: Violation of local nonsatiation



Figure 5: Compatible with local nonsatiation

Exercise 3.B.2

The preference relation $\succsim$ defined on the consumption set $X = \mathbb{R}^L_+$ is said to be *weakly monotone* if and only if $x \geq y$ implies that $x \succsim y$. Show that if $\succsim$ is transitive, locally nonsatiated, and weakly monotone, then it is monotone.

Convexity Assumptions

Definition 3.B.4. The preference relation $\succsim$ on X is *convex* if for every $x \in X$, the upper contour set of x, $\{y \in X : y \succsim x\}$ is convex; that is, if $y \succsim x$ and $z \succsim x$, then $\alpha y + (1-\alpha)z \succsim x$ for any $\alpha \in [0,1]$.



Figure 6: Convex Preference



Figure 7: Nonconvex Preference

Properties associated with convexity

(i) *Diminishing marginal rates of substitution*: with convex preferences, from any initial consumption X, and for any two commodities, it takes an increasingly larger amounts of one commodity to compensate for successive unit losses of the other.

(ii) Preference for diversity (implied by (i)): under convexity, if x is indifferent to y, then $\frac{1}{2}x + \frac{1}{2}y$ cannot be worse than x or y.

Definition 3.B.5. The preference relation $\succsim$ on X is *strictly convex* if for every $x \in X$, we have that $y \succsim x$ and $z \succsim x$, and $y \neq z$ implies $\alpha y + (1-\alpha)z \succ x$ for all $\alpha \in (0,1)$.

Homothetic and Quasilinear Preference In applications (particularly those of an econometric nature), it is common to focus on preferences for which it is possible to deduce the consumer's entire preference relation from a single indifference set. Two examples are the classes of *homothetic* and *quasilinear* preferences.

Definition 3.B.6. A monotone preference relation $\succsim$ on $X = \mathbb{R}^L_+$ is *homothetic* if all indifference sets are related by proportional expansion along rays; that is, if $x \sim y$, then $\alpha x \sim \alpha y$ for any $\alpha \geq 0$.



Figure 8: Homothetic Preference

Definition 3.B.7. The preference relation $\succsim$ on $X = (-\infty, \infty) \times \mathbb{R}^{L-1}_+$ is *quasilinear* with respect to commodity 1 (the *numeraire* commodity) if

(i) All the indifference sets are parallel displacements of each other along the axis of commodity 1. That is, if $x \sim y$, then $(x + \alpha e_1) \sim (y + \alpha e_1)$ for $e_1 = (1, 0, 0, ..., 0)$ and any $\alpha \in \mathbb{R}$.

(ii) Good 1 is desirable; that is $x + \alpha e_1 \succ x$ for all x and $\alpha > 0$.



Figure 9: Quasilinear Preference

3.C. Preference and Utility

Key Question. When can a rational preference relation be represented by a utility function?

Answer: If the preference relation is continuous.

Definition 3.C.1. The preference relation $\succsim$ on X is *continuous* if it is preserved in the limits. That is, for any sequence of pairs $\{(x^n, y^n)\}_{n=1}^{\infty}$ with $x^n \succsim y^n$ for all n, $x = \lim_{n\to\infty} x^n$, $y = \lim_{n\to\infty} y^n$, we have $x \succsim y$.

Interpretation: Consumer's preferences cannot exhibit jumps. For example, the consumer cannot prefer each elements in the sequence x^n to the corresponding element in the sequence y^n but suddenly reverse her preference at the limiting points of these sequences x and y.

Claim. $\succsim$ is continuous if and only if for all x, the upper contour set $\{y \in X : y \succsim x\}$ and the lower contour set $\{y \in X : x \succsim y\}$ are both closed.

Proof. We only provide the proof for "only if" part. "if" part is more advanced and not required by this course.

Definition 3.C.1 implies that for any sequence of points $\{y^n\}_{n=1}^{\infty}$ with $x \succsim y^n$ for all n and $y = \lim_{n\to\infty} y^n$, we have $x \succsim y$ (just let $x^n = x$ for all n). Hence the closedness of lower contour set is implied. Similarly, we could show the closedness of upper contour set. $\square$

Example 3.C.1. Lexicographic Preference Relation on $\mathbb{R}^2$

$x \succ y$ if either $x_1 > y_1$, or $x_1 = y_1$ and $x_2 > y_2$.

$x \sim y$ if $x_1 = y_1$ and $x_2 = y_2$.

Claim. Lexicographic Preference Relation on $\mathbb{R}^2$ is not continuous.

Proof. Consider sequence of bundles (x^n, y^n) where $x^n = (1 + \frac{1}{n}, 1)$ and $y^n = (1, 2)$. $x = \lim_{n\to\infty} x^n = (1, 1)$, $y = \lim_{n\to\infty} y^n = (1, 2)$. $x^n \succ y^n$ for all n but $x \prec y$. $\square$

Claim. Lexicographic Preference Relation on $\mathbb{R}^2$ cannot be represented by $u(\cdot)$.

Proof. Here, we only provide a sketch of proof with the help of Figure 10. The contradiction comes from the one-to-one mapping $r(\cdot) : \mathbb{R} \to \mathbb{Q}$. For detailed proof, please refer to Chapter 1 Lecture Notes.



Figure 10: Lexicographic Preference

Alternatively, we could use the fact that upper and lower contour sets of a continuous preference must be closed. It is shown in Figure 11 and 12 that the upper and lower contour set of Lexicographic preference are not closed.



Figure 11: Upper Contour Set



Figure 12: Lower Contour Set

□

Proposition 3.C.1. *Suppose that the preference relation $\succsim$ on X is continuous. Then there exists continuous utility function $u(x)$ that represents $\succsim$, i.e., $u(x) \geq u(y)$ if and only if $x \succsim y$.*

For this course, we will only prove a simplified version of Proposition 3.C.1, which assumes that the preference relation $\succsim$ is also monotone.

Proof. Define $Z := \{x \in \mathbb{R}^L_+ : x_l = x_k \text{ for all } k, l = 1, ..., L\}$ Define $e := (1, ..., 1)$ with L elements. Then, $\alpha e \in Z$ for $\alpha \geq 0$.

Step 1: Construction of a utility function



Figure 13: Construction of Utility Function

We want to show that for all $x \in \mathbb{R}^L_+$, $\exists$ unique $\alpha \geq 0$ such that $\alpha e \sim x$, $\alpha_1 e \succ x$ if $\alpha_1 > \alpha$, $x \succ \alpha_2 e$ if $\alpha_2 < \alpha$.

Monotonicity implies $x \succsim 0$. Also, for $\overline{\alpha}$ such that $\overline{\alpha}e \gg x$, we have $\overline{\alpha}e \succ x$. Monotonicity and continuity of preference can be used to show that there exists a unique $\alpha(x) \in [0, \overline{\alpha}]$, such that $\alpha(x)e \sim x$.

Formally, continuity of preference implies that $A^+ = \{\alpha \in \mathbb{R}_+ : \alpha e \succsim x\}$ and $A^- = \{\alpha \in \mathbb{R}_+ : x \succsim \alpha e\}$ are closed.

Define $\underline{\alpha}^+ := \min\{A^+\}$, and $\overline{\alpha}^- := \max\{A^-\}$.

Then by monotonicity, $\alpha e \succ x$, $\forall \alpha > \underline{\alpha}^+$ and $\alpha e \prec x$, $\forall \alpha < \overline{\alpha}^-$.

Now we show that $\underline{\alpha}^+ = \overline{\alpha}^-$.

Suppose $\underline{\alpha}^+ > \overline{\alpha}^-$. Then $\exists \alpha \in (\overline{\alpha}^-, \underline{\alpha}^+)$ such that $\alpha e \not\succsim x$ and $x \not\succsim \alpha e \implies$ incomplete preference;

Suppose $\underline{\alpha}^+ < \overline{\alpha}^-$. Then $\exists \alpha \in (\overline{\alpha}^-, \underline{\alpha}^+)$ such that

$$\alpha e \succ x (\because \alpha > \underline{\alpha}^+) \quad \& \quad \alpha e \prec x (\because \alpha < \overline{\alpha}^-) \implies \text{contradiction}$$

Thus, there exists a scalar α such that $\alpha e \sim x$.

Furthermore, by monotonicity, $\alpha_1 e \succ \alpha_2 e$ whenever $\alpha_1 > \alpha_2$. Hence, there can be at most one scalar satisfying $\alpha e \sim x$. This scalar is $\alpha(x)$.

Set $u(x) = \alpha(x)$. We have shown that $u(x)$ exists and is unique.

Step 2: $u(x)$ represents $\succsim$, i.e., $\alpha(x) \geq \alpha(y)$ iff $x \succsim y$.

Note that $\alpha(x)e \sim x$ (Step 1).

If $\alpha(x) \geq \alpha(y)$, then $x \sim \alpha(x)e \succsim \alpha(y)e \sim y (\because$ preference is monotone$) \implies x \succsim y$.

If $x \succsim y$, then $\alpha(x)e \succsim \alpha(y)e$, monotonicity implies $\alpha(x) \geq \alpha(y)$.

Since $u(x) = \alpha(x)$, $u(x)$ represents $\succsim$.

Step 3: $u(x) = \alpha(x)$ is continuous, i.e., take any sequence $\{x^n\}_{n=1}^{\infty}$ with $x = \lim_{n\to\infty} x^n$, we have $\lim_{n\to\infty} \alpha(x^n) = \alpha(x)$.



Figure 14: Proof of continuity

Note that $x^n \geq 0$ so $\alpha(x^n) \geq 0$ for all n. Also, there exists $\alpha_1 > 0$ such that $\alpha_1 e \succ x$. For any sufficiently small $\varepsilon > 0$, for all $x' \in X$ such that $\|x' - x\| \leq \varepsilon$, we have $\alpha_1 e \succ x'$ and $\alpha(x') < \alpha_1$. In other words, $\exists N$ s.t. $\forall n > N$, $0 \leq \alpha(x^n) \leq \alpha_1$. Since $\alpha(x^1), ..., \alpha(x^N)$ are also finite, every element in the sequence $\{\alpha(x^n)\}_{n=1}^{\infty}$ is finite and belongs to some compact space $[0, \alpha_2]$ for some sufficiently large α_2. By Bolzano-Weierstrass Theorem, **the infinite sequence $\{\alpha(x^n)\}_{n=1}^{\infty}$ must have a convergent subsequence.**

Now we show that **all convergent subsequences of $\{\alpha(x^n)\}_{n=1}^{\infty}$ coverges to $\alpha(x)$.**

Suppose otherwise and that $\lim_{n\to\infty} \alpha\left(x^{m(n)}\right) = \alpha' > \alpha(x)$. [The proof for the case $\alpha(x) > \alpha'$ is similar.] Let $\hat{\alpha} = \frac{1}{2}(\alpha' + \alpha(x))$, so $\alpha(x) < \hat{\alpha} < \alpha', x \sim \alpha(x)e \prec \hat{\alpha}\varepsilon \prec \alpha'\varepsilon$.

Recall that $x^{m(n)} \sim \alpha\left(x^{m(n)}\right) e$ for all n by construction.

(a) Since $\lim_{n\to\infty} \alpha\left(x^{m(n)}\right) = \alpha', \exists \overline{N}$ s.t. $\forall n \geq \overline{N}, x^{m(n)} \sim \alpha\left(x^{m(n)}\right) e \succ \hat{\alpha} e$.

(b) On the other hand, since $\lim_{n\to\infty} x^{m(n)} = x \prec \hat{a}e$, continuity of preference implies $\exists \widetilde{N}$ s.t. $\forall n \geq \widetilde{N}, x^{m(n)} \prec \hat{a}e$.

Combining (a) and (b), $\forall n \geq \max\{\overline{N}, \widetilde{N}\}, \hat{a}e \succ x^{m(n)} \sim \alpha\left(x^{m(n)}\right) e \succ \hat{\alpha} e \implies$ Contradiction.

Therefore, we can conclude that $\{\alpha(x^n)\}_{n=1}^{\infty}$ converges to $\alpha(x)$. $\square$

Remark. $u(x)$ is not unique, any increasing transformation $v(x) = f(u(x))$ will represent $\succsim$. **We can also introduce countably many jumps in $f(\cdot)$.**

Assumptions of differentiability of $u(x)$ The assumption of differentiability is commonly adopted for technical convenience, but is not applicable to all useful models. Here is an example of preference that is not differentiable.

Example (Leontief Preference)**.** $x'' \succsim x'$ if and only if $\min\{x_1'', x_2''\} \geq \min\{x_1', x_2'\}$.



Figure 15: Leontief Preference

$u(x_1, x_2) = \min\{x_1, x_2\}$ represents Leontief preference. $u(x_1, x_2)$ is not differentiable because of the kink in the indifference curves when $x_1 = x_2$, i.e., when $x = (x_1, x_1)$.

To see this, for the first variable x_1 (similar argument applies to the second variable):

$$\lim_{\varepsilon\to 0^-}\frac{u(x_1+\varepsilon,x_1)-u(x_1,x_1)}{\varepsilon}=\lim_{\varepsilon\to 0^-}\frac{\min\{x_1+\varepsilon,x_1\}-\min\{x_1,x_1\}}{\varepsilon}=\lim_{\varepsilon\to 0^-}\frac{x_1+\varepsilon-x_1}{\varepsilon}=1;$$
$$\lim_{\varepsilon\to 0^+}\frac{u(x_1+\varepsilon,x_1)-u(x_1,x_1)}{\varepsilon}=\lim_{\varepsilon\to 0^+}\frac{\min\{x_1+\varepsilon,x_1\})-\min\{x_1,x_1\}}{\varepsilon}=\lim_{\varepsilon\to 0^+}\frac{x_1-x_1}{\varepsilon}=0.$$

Implications of $\succsim$ and u

(i) $\succsim$ is convex $\iff$ $u: X\to\mathbb{R}$ is quasi-concave.

(ii) continuous $\succsim$ on $\mathbb{R}^L_+$ is homothetic $\iff$ $u(x)$ is H.D.1.

(iii) continuous $\succsim$ on $(-\infty,\infty)\times\mathbb{R}^{L-1}_+$ is quasilinear with respect to Good 1 $\iff$ $u(x)=x_1+\phi(x_2,...,x_L)$

Definition. The utility function $u(\cdot)$ is *quasiconcave* if the set $\{y\in\mathbb{R}^L_+ : u(y)\geq u(x)\}$ is convex for all x or, equivalently, if $u(\alpha x+(1-\alpha)y)\geq\min\{u(x),u(y))\}$ for all x,y and all $\alpha\in[0,1]$. If $u(\alpha x+(1-\alpha)y)>\min\{u(x),u(y)\}$ for $x\neq y$ and $\alpha\in(0,1)$, then $u(\cdot)$ is *strictly quasiconcave.*

Proof.

(i) $\succsim$ is convex. $\iff$ If $y\succsim x, z\succsim x$, then $\alpha y+(1-\alpha)z\succsim x \quad \forall\alpha\in[0,1]$.

$\iff$ If $u(y)\geq u(x), u(z)\geq u(x)$, then $u(\alpha y+(1-\alpha)z)\geq u(x) \quad \forall\alpha\in[0,1]$.

$\iff$ $\{y\in\mathbb{R}^L_+ : u(y)\geq u(x)\}$ is convex.

$\iff$ $u: X\to\mathbb{R}$ is quasi-concave.

(ii) (a) "$\Leftarrow$": Suppose $u(x)$ is H.D.1., i.e., $u(\alpha x)=\alpha u(x)$ and $u(\alpha y)=\alpha u(y)$.

Also suppose $x\sim y \iff u(x)=u(y)$.

Then $u(\alpha y)=\alpha u(y)=\alpha u(x)=u(\alpha x) \implies \alpha x\sim\alpha y \implies \succsim$ is homothetic.

(b) "$\Rightarrow$": Suppose $\succsim$ is homothetic. Suppose $u(x)e\sim x$. It is W.L.O.G to take $u(x)$ as a utility function.

Homothetic $\succsim$ implies $\alpha u(x)e\sim\alpha x$. We also have $u(\alpha x)e\sim\alpha x$.

Then $\alpha u(x)e\sim u(\alpha x)e \implies u(\alpha x)=\alpha u(x) \implies u(x)$ is H.D.1.

(iii) a) "$\Leftarrow$": Suppose $u(x) = x_1 + \phi(x_2, ..., x_L)$. Then, $u(x + \alpha e_1) = x_1 + \alpha + \phi(x_2, \cdots, x_L) = \alpha + u(x)$. Similarly, $u(y + \alpha e_1) = \alpha + u(y)$.

Therefore, $x \sim y \implies u(x) = u(y) \implies u(x + \alpha e_1) = u(y + \alpha e_1) \implies (x + \alpha e_1) \sim (y + \alpha e_1) \implies \succsim$ is quasilinear.

b) "$\Rightarrow$": In general, for some consumption bundle $(0, x_2, \cdots, x_L)$, there exists a consumption bundle $(x_1^*, 0, \cdots, 0)$, such that $(0, x_2, \cdots, x_L) \sim (x_1^*, 0, \cdots, 0)$.

We therefore define the mapping from $(x_2, \cdots, x_L)$ to x_1^* by a function $x_1^* = \phi(x_2, \cdots, x_L)$. From $\succsim$ being quasilinear, we have

$$(x_1, x_2, \cdots, x_L) \sim (x_1 + \phi(x_2, \cdots, x_L), 0, \cdots, 0).$$

Therefore, $\succsim$ admits a utility function of the form $u(x) = x_1 + \phi(x_2, ..., x_L)$. $\square$

Exercise 3.C.6

Suppose that in a two-commodity world, the consumer's utility function takes the form $u(x) = [\alpha_1 x_1^\rho + \alpha_2 x_2^\rho]^{1/\rho}$. This utility function is known as the *constant elasticity of substitution* (or *CES*) utility function.

(a) Show that when $\rho = 1$, indifference curves become linear.

(b) Show that as $\rho \to 0$, this utility function comes to represent the same preferences as the (generalized) Cobb-Douglas utility funciton $u(x) = x_1^{\alpha_1} x_2^{\alpha_2}$.

(c) Show that as $\rho \to -\infty$, indifference curves become "right angles"; that is, this utility funciton has in the limit the indifference map of the Leontief utility function $u(x_1, x_2) = \min\{x_1, x_2\}$.

3.D. Utility Maximization Problem (UMP)

We assume throughout that preference is rational, continuous, and locally nonsatiated, and we take $u(x)$ to be a continuous utility function representing these preferences. We also assume that the consumption set is $X = \mathbb{R}_+^L$.

The consumer's problem of choosing her most preferred consumption bundle $x(p, w)$ can be stated as a *Utility Maximization Problem (UMP)*:

$$\max_{x \in \mathbb{R}^L_+} \quad u(x)$$
$$\text{s.t. } p \cdot x \leq w$$

Proposition 3.D.1. *If $p \gg 0$ and $u(\cdot)$ is continuous, then the utility maximization problem has a solution.*

Proof. $B_{p,w} = \{x \in \mathbb{R}^L_+ : p \cdot x \leq w\}$ is compact, i.e,

(i) bounded: $0 \leq x_l \leq w/p_l$, for $p_l > 0$

(ii) closed (it contains all the limit points): Proof by contradiction.

Consider a sequence $\{x^n\}_{n=1}^{\infty}$ where $x^n \in B_{p,w}$, or $p \cdot x^n \leq w$ for all n and $x = \lim_{n \to \infty} x^n \notin B_{p,w}$ or $p \cdot x > w$. There exists $\varepsilon > 0$ such that for all y satisfying $\|y - x\| < \varepsilon, p \cdot y > w$. Therefore, $\exists N > 0$, s.t. $\forall n \geq N$, $p \cdot x^n > w$. This contradicts $p \cdot x^n \leq w$ for all n.

By *Extreme Value Theorem*, a continuous function always has a maximum value on any compact set. □

Here, we provide two counter examples where the solution of UMP does not exists.

Counter Examples.

(i) $B_{p,w}$ is not closed: $p \cdot x < w$

(ii) $u(x)$ is not continuous: $u(x) = \begin{cases} p \cdot x & \text{for } p \cdot x < w \\ 0 & \text{for } p \cdot x = w \end{cases}$

Properties of the Walrasian demand correspondence/functions The solution of UMP, denoted by $x(p, w)$, is called *Walrasian (*or *ordinary* or *market) demand correspondence.* When $x(p, w)$ is single valued for all (p, w), we refer to it as *Walrasian (*or *ordinary* or *market) demand function.*



Figure 16: Single solution



Figure 17: Multiple solutions

Proposition 3.D.2. *Suppose that $u(x)$ is a continuous utility function representing a locally nonsatiated preference relation $\succsim$ defined on the consumption set $X = \mathbb{R}^L_+$. Then the Walrasian demand correspondence $x(p,w)$ possesses the following properties:*

(i) Homogeneity of degree zero in (p,w) : $x(\alpha p, \alpha w) = x(p,w)$ for any p, w and scalar $\alpha > 0$.

(ii) Walras' Law: $p \cdot x = w$ for all $x \in x(p,w)$.

(iii) Convexity/uniqueness: If $\succsim$ is convex, so that $u(\cdot)$ is quasiconcave, then $x(p,w)$ is a convex set. Moreover, if $\succsim$ is strictly convex, so that $u(\cdot)$ is strictly quasiconcave, then $x(p,w)$ consists of a single element.

Recall,

Definition. The preference relation $\succsim$ on X is *convex* if for every $x \in X$, the upper contour set of x, $\{y \in X : y \succsim x\}$ is convex; that is, if $y \succsim x$ and $z \succsim x$, then $\alpha y + (1-\alpha)z \succsim x$ for any $\alpha \in [0,1]$.

Definition. The utility function $u(\cdot)$ is *quasiconcave* if the set $\{y \in \mathbb{R}^L_+ : u(y) \geq u(x)\}$ is convex for all x or, equivalently, if $u(\alpha x + (1-\alpha)y) \geq \min\{u(x, u(y))\}$ for all x, y and all $\alpha \in [0,1]$.

Proof.

(i) H.D.$\emptyset$: $\{x \in \mathbb{R}^L_+ : p \cdot x \leq w\} = \{x \in \mathbb{R}^L_+ : \alpha p \cdot x \leq \alpha w\}$.

The set of feasible consumption bundles in the UMP is unaffected by α. Therefore, $x(p, w) = x(\alpha p, \alpha w)$.

(ii) Walras' Law: Suppose $p \cdot x(p, w) < w$. Then $\exists \varepsilon > 0$ such that $\forall y$ such that $\|y - x(p, w)\| < \varepsilon$, $p \cdot y < w$. Local nonsantiation implies $\exists y$ with $y \in X$ and $\|y - x(p, w)\| < \varepsilon$ such that $y \succ x(p, w) \Longrightarrow$ contradiction with $x(p, w)$ being optimal.

(iii) Suppose $x, x' \in x(p, w)$ and $x \neq x'$. Then $u(x) = u(x')$. Quasiconcavity of u implies $u(\alpha x + (1 - \alpha)x') \geq \min\{u(x), u(x')\} \implies \alpha x + (1 - \alpha)x' \in x(p, w)$

Now suppose $u(x)$ is strictly quasiconcave. Suppose $x, x' \in x(p, w)$ and $x \neq x'$. Then $u(x) = u(x')$. Strict quasiconcavity implies $u(\alpha x + (1 - \alpha)x') > u(x) = u(x')$. This contradicts that $x, x' \in x(p, w)$. Therefore, $x(p, w)$ is single valued.

Alternative proof for (iii): Suppose $x, x' \in x(p, w)$. Then $x \sim x' \succsim y$, $\forall y \in B_{p,w}$. $\overline{x} = \alpha x + (1 - \alpha)x' \succsim x \sim x' \implies \overline{x} \succsim y, \forall y \in B_{p,w}$. Strict convexity $\overline{x} \succ x \sim x'$ which contradicts $x, x' \in x(p, w)$. $\square$

We will take a break to review some mathematical results before proceeding with this Chapter. Read "Math Review: Maximization Problem" for details.