Enhanced Langlands parameters of regular principal series

Xiaohang Yang

Joint work with Haoming Li, Jizhi Zhang, and Yelei Zhao

Beijing International Center for Mathematical Research
Peking University

4th National Doctoral Forum on Frontiers of Mathematics

November 5, 2026

Advised by Prof. Dongwen Liu

Let G be a connected reductive group over a finite extension $F/\mathbb{Q}_p$ with residue field of order q_F.

The local Langlands correspondence

$$\left\{\begin{array}{c}\text{irreducible smooth}\\ \text{representations}\\ \pi \text{ of } G(F)\end{array}\right\}/\text{iso.} \xrightarrow{\mathcal{L}_G} \left\{\begin{array}{c}L\text{-parameters for } G\\ \phi : W_F \times \mathrm{SL}_2(\mathbb{C}) \to {}^L G\end{array}\right\}/G^\vee\text{-conj.}$$

Write $\Phi(G)$ for the set on the right. Each fibre is a finite L-packet.

$$\Pi_\phi = \mathcal{L}_G^{-1}([\phi]), \qquad [\phi] \in \Phi(G).$$

How can we refine the local Langlands correspondence to obtain a bijection?

Several representations can share the same parameter ϕ. The enhanced LLC seeks a bijection by labelling each packet member on the Galois side.

$$\pi \quad \longleftrightarrow \quad (\phi_\pi, \rho_\pi).$$

The **enhancement** ρ_π labels π within its packet. It is an irreducible representation of a finite component group attached to ϕ_π.

The component group for quasi-split G

Let $C_\phi = Z_{G^\vee}(\operatorname{im}\phi)$ be the centralizer of the parameter. Then

$$\mathcal{R}_\phi = \pi_0\Big(C_\phi / Z(G^\vee)^{W_F}\Big), \qquad \rho_\pi \in \operatorname{Irr}(\mathcal{R}_\phi) \quad (\pi \in \Pi_\phi).$$

Here π_0 denotes the group of connected components.

The category of smooth complex representations has the **Bernstein decomposition**

$$\mathrm{Rep}(G(F)) \simeq \prod_{\mathfrak{s}\in\mathfrak{B}(G)} \mathrm{Rep}(G(F))^{\mathfrak{s}}.$$

Every representation decomposes canonically as $\pi \simeq \bigoplus_{\mathfrak{s}} \pi^{\mathfrak{s}}$.

Here $\mathfrak{B}(G)$ consists of inertial classes $\mathfrak{s} = [M, \sigma]_G$, where M is a Levi subgroup and σ is supercuspidal. Inertial equivalence allows conjugacy and unramified twists of σ.

The block $\mathrm{Rep}(G(F))^{\mathfrak{s}}$ consists of representations whose irreducible subquotients have supercuspidal support in $\mathfrak{s}$. For principal series, $M = T$ and $\sigma = \chi$ is a torus character.

Example for $G = \mathrm{GL}_2$

$$\mathbf{1}, \mathrm{St} \in \mathrm{Irr}(G(F))^{[T,\mathbf{1}]_G}, \qquad \phi_{\mathbf{1}} \not\sim \phi_{\mathrm{St}}.$$

Representations in one Bernstein block can belong to different L-packets.

Solleveld (2025)

For quasi-split G with a fixed Whittaker datum, the principal-series LLC is a bijection.

$$\left\{\begin{array}{c}\text{irreducible smooth}\\ \text{principal-series}\\ \text{representations of } G(F)\end{array}\right\}/\text{iso.} \xrightarrow[\sim]{\mathcal{L}^e_{\mathrm{ps}}} \left\{\begin{array}{c}\text{principal-series enhanced}\\ L\text{-parameters } (\phi,\rho)\\ \rho \in \mathrm{Irr}(\mathcal{R}_\phi)\end{array}\right\}/G^\vee\text{-conj.}$$

Forgetting the enhancement

$$\pi \longmapsto (\phi_\pi, \rho_\pi) \longmapsto \phi_\pi.$$

Projection onto the first entry recovers the ordinary packet map.

A trivial enhancement $\rho = 1$ need not mean a trivial component group.

Let ${}^L G = G^\vee \rtimes W_F$. An L-**parameter** is an admissible homomorphism

$$\phi : W_F \times \mathrm{SL}_2(\mathbb{C}) \longrightarrow {}^L G, \qquad \text{up to } G^\vee\text{-conjugacy.}$$

It is continuous on W_F and algebraic on $\mathrm{SL}_2(\mathbb{C})$.

Equivalent Weil–Deligne form

$$(\widetilde{\phi}, N), \qquad \widetilde{\phi} : W_F \longrightarrow {}^L G, \qquad N \in \mathrm{Lie}(G^\vee) \text{ nilpotent},$$
$$\widetilde{\phi}(w) := \phi\Big(w, \mathrm{diag}(\|w\|^{1/2}, \|w\|^{-1/2})\Big).$$
$$\mathrm{Ad}(\widetilde{\phi}(w))N = \|w\| N.$$

An **enhanced** L-**parameter** can therefore be written as a triple $(\widetilde{\phi}, N, \rho)$. Here $\widetilde{\phi}$ is the **infinitesimal parameter**, N the **monodromy**, and ρ the **enhancement**.

The setting: regular principal series

Let G/F be connected, quasi-split, and reductive, with an F-Borel $B = TU$. For a smooth character $\chi : T(F) \to \mathbb{C}^\times$, form

$$I(\chi) = \mathrm{Ind}_B^G \chi.$$

The character χ is **regular** if

$$\mathrm{Stab}_W(\chi) = \{1\}.$$

Here $W = N_G(S)(F)/T(F)$ is the relative Weyl group, where $S \subset T$ is maximal F-split.

- The torus LLC determines the common infinitesimal parameter of the constituents. It attaches $\widehat{\chi} : W_F \to {}^L T \hookrightarrow {}^L G$ to χ, and

$$[\widetilde{\phi}_\pi] = [\widehat{\chi}] \quad \text{for every irreducible constituent } \pi \text{ of } I(\chi).$$

- We want the monodromy and enhancement of *each* irreducible constituent of $I(\chi)$.

Every constituent of $I(\chi)$ has the common infinitesimal parameter

$$\widetilde{\phi} = \widehat{\chi}.$$

We seek its monodromy N and enhancement ρ.

For regular χ, the unique irreducible subrepresentation is $\operatorname{soc} I(\chi)$.

The motivating question

When is $N_{\mathrm{soc}} = 0$? Can this be decided from reducibility over F?

Bellaïche–Chenevier (2009) gave a sufficient condition after passing to a splitting field. We seek an intrinsic necessary and sufficient condition over F.

Earlier results on regular principal series

- **Rodier (1981), split groups.**
  Classified the irreducible constituents by chambers of the reducibility arrangement.
- **Rodier (1984), split groups.**
  Described their L-parameters, including a zero-monodromy criterion for the socle.
- **Luo (2021), general reductive groups.**
  Extended the chamber classification to regular generalized principal series, with Jacquet modules, multiplicity one, and independent reducibility coroots.
- **Solleveld (2025).**
  His enhanced principal-series LLC and joint work on geometric Hecke algebras provide a framework in which the enhancement can be computed.

Our aim: compute the monodromy N and enhancement ρ of every constituent of regular principal series for arbitrary quasi-split groups.

Reducibility directions and oriented walls

For each relative root line, test the rank-one induction

$$\mathrm{Ind}_{B\cap M_\alpha}^{M_\alpha}(\chi).$$

On a reducible root line, choose an integral coroot map $H_a : F^\times \to S(F) \subset T(F)$. Choose the sign of a so that

$$|\chi(H_a(t))| = |t|^{s_a}, \qquad s_a > 0.$$


Collect these directions in $\mathcal{S}(\chi)$. Set

$$\mathfrak{a}_T^* := \mathrm{Hom}\big(T(F)/T(F)_{\mathrm{cpt}}, \mathbb{R}\big),$$

where $T(F)_{\mathrm{cpt}}$ is the maximal compact subgroup of $T(F)$.


$$\mathcal{H}_a = \{x \in \mathfrak{a}_T^* : \langle x, a\rangle = 0\},$$
$$\mathcal{H}_a^+ = \{x \in \mathfrak{a}_T^* : \langle x, a\rangle > 0\}.$$

The connected regions are the chambers.



Rodier–Luo

The directions in $\mathcal{S}(\chi)$ are independent. Chambers label the pairwise nonisomorphic constituents of $I(\chi)$, each with multiplicity one.



Use $x_i = \langle x, a_i \rangle$. Two independent directions give four sign patterns and four constituents. In general: $2^{|\mathcal{S}(\chi)|}$ constituents.

Main theorem: positive sides determine monodromy, with trivial enhancement

Each reducibility direction $a \in \mathcal{S}(\chi)$ determines a one-dimensional monodromy root space. Choose a nonzero vector $X_{-\iota(a)}$ in that space.

Li–Yang–Zhang–Zhao


The enhanced parameter of π_Γ has a representative with

$$\widetilde{\phi}_\Gamma = \widehat{\chi}, \qquad N_\Gamma \sim \sum_{\substack{a \in \mathcal{S}(\chi) \\ \Gamma \subset \mathcal{H}_a^+}} X_{-\iota(a)}, \qquad \rho_\Gamma = \mathbf{1}.$$

Here $\sim$ denotes conjugacy under $Z_{G^\vee}(\widehat{\chi}(W_F))^\circ$.

- All irreducible constituents of the principal series $I(\chi)$ share the same infinitesimal parameter.
- The chamber determines the monodromy support.
- Every enhancement is trivial; the component group need not be.

Reading the formula: four chambers, four supports



Here $X_i = X_{-\iota(a_i)}$. All four constituents share $\widetilde{\phi} = \widehat{\chi}$. All enhancements are trivial, even though the component groups may differ.

A positive sign includes that root direction in the support.

When does the socle have zero monodromy?

Let C^+ be the dominant Weyl chamber determined by B, and let Γ_{soc} be the arrangement chamber containing it.

$$\operatorname{soc} I(\chi) = \pi_{\Gamma_{\mathrm{soc}}}.$$

A criterion using reducibility over F

$$N_{\mathrm{soc}} = 0 \quad \Longleftrightarrow \quad \langle x, a \rangle < 0 \quad (\forall x \in C^+,\ \forall a \in \mathcal{S}(\chi)).$$

Equivalently, C^+ lies on the negative side of every oriented reducibility wall. In this case,

$$\phi_{\mathrm{soc}}(w, g) = \widehat{\chi}(w), \qquad \rho_{\mathrm{soc}} = \mathbf{1}.$$

GL_2: two inducing orders

Let $\nu = |\cdot|_F$, let B be upper triangular, and put $s = \mathrm{diag}(q_F^{-1/2}, q_F^{1/2})$.

$$0 \longrightarrow \mathrm{St} \longrightarrow \mathrm{Ind}_B^{\mathrm{GL}_2}(\nu^{1/2} \boxtimes \nu^{-1/2}) \longrightarrow \mathbf{1} \longrightarrow 0.$$

Constituent	Wall side	$(\widetilde{\phi}(\mathrm{Fr}_F), N)$	ρ
St (socle)	$+$	(s, E_{12})	$\mathbf{1}$
$\mathbf{1}$ (quotient)	$-$	$(s, 0)$	$\mathbf{1}$

$$0 \longrightarrow \mathbf{1} \longrightarrow \mathrm{Ind}_B^{\mathrm{GL}_2}(\nu^{-1/2} \boxtimes \nu^{1/2}) \longrightarrow \mathrm{St} \longrightarrow 0.$$

Constituent	Wall side	$(\widetilde{\phi}(\mathrm{Fr}_F), N)$	ρ
$\mathbf{1}$ (socle)	$-$	$(s^{-1}, 0)$	$\mathbf{1}$
St (quotient)	$+$	(s^{-1}, E_{21})	$\mathbf{1}$

In each order, the constituents share an infinitesimal character but belong to different L-packets.

The socle in the lower table is $\mathbf{1}$, with zero monodromy.

Proof roadmap

$$\underbrace{\pi_\Gamma}_{\text{representation}} \mapsto \underbrace{M_\Gamma}_{\substack{\text{affine Hecke}\\ \text{module}\\ \text{(representation side)}}} \mapsto \underbrace{M_\Gamma^\vee}_{\substack{\text{affine Hecke}\\ \text{module}\\ \text{(Galois side)}}} \mapsto \underbrace{V_\Gamma}_{\substack{\text{graded Hecke}\\ \text{module}}} \mapsto \underbrace{(\sigma_\Gamma, N_\Gamma, \rho = \mathbf{1})}_{\substack{\text{geometric}\\ \text{parameter}}} \mapsto \underbrace{(\widetilde{\phi}_\Gamma, N_\Gamma, \rho_\Gamma)}_{\substack{\text{enhanced}\\ L\text{-parameter}}}.$$

1. **Bernstein block: pass to an affine Hecke module.**
2. **Hecke algebra comparison: transfer to the Galois side.**
3. **Fix the central character and reduce to a graded Hecke module.**
4. **Geometry: identify the triple attached to V_Γ.**
5. **Translate the triple into the enhanced Langlands parameter.**

1. **The space of possible monodromy.** Fix the semisimple datum. For $a \in \mathcal{S}(\chi)$, put $\alpha_a := -v^{-1}\iota(a)$. Then

$$\text{Possible Monodromy} = \bigoplus_{a \in \mathcal{S}(\chi)} \mathbb{C} X_{\alpha_a}.$$

   Every element of this space is nilpotent.

2. **Conjugacy classes are parametrized by subsets.** Independent root characters allow the centralizing torus to rescale nonzero coordinates:

$$\{\text{possible monodromy conjugacy classes}\} \longleftrightarrow \{J \subseteq \mathcal{S}(\chi)\}, \qquad N_J = \sum_{a \in J} X_{\alpha_a}.$$

3. **Which subset corresponds to the representation π_Γ?**

Proof: the chamber determines monodromy

Use weights to determine the subset: Let σ_Γ be a selected weight of V_Γ.

$$\pi_\Gamma \longmapsto M_\Gamma \longmapsto M_\Gamma^\vee \longmapsto V_\Gamma, \qquad \dim(V_\Gamma)_{\sigma_\Gamma} = 1.$$

1. **Geometric classification.** For an eligible triple $(\sigma_\Gamma, N_J, \rho)$, the standard module E_J has a unique irreducible quotient L_J.
2. **Weights and Möbius inversion.** Weight calculations and Möbius inversion determine the subset

$$J_\Gamma = \{a \in \mathcal{S}(\chi) : \Gamma \subset \mathcal{H}_a^+\}.$$

In particular,

$$\boxed{V_\Gamma \simeq L_{J_\Gamma}, \qquad N_\Gamma \sim \sum_{a \in J_\Gamma} X_{\alpha_a}.}$$

Proof: why the enhancement is trivial

Use generalized Springer theory for the monodromy N_Γ.

1. **Possible enhancements.** A possible enhancement $\rho \in \mathrm{Irr}(\pi_0 Z_H(\sigma_\Gamma, N_\Gamma))$ must occur in $H_*(\mathcal{P}_{N_\Gamma}^{\sigma_\Gamma}; \mathbb{C})$ after specialization.
2. **Only trivial enhancements appear.** Regularity makes the fixed locus $\mathcal{P}_{N_\Gamma}^{\sigma_\Gamma}$ finite and discrete. The group $\pi_0 Z_H(\sigma_\Gamma, N_\Gamma)$ acts trivially on its homology.

Thus the possible enhancement for the triple $(\sigma_\Gamma, N_\Gamma, \rho)$ is trivial:

$$\boxed{\rho = \mathbf{1}.}$$

The component group itself may still be nontrivial.

Hecke modules and enhanced L-parameters

At the chosen bounded datum and central character, the parametrizations complete the chain:

$$\pi_\Gamma \longmapsto M_\Gamma \longmapsto M_\Gamma^\vee \longmapsto V_\Gamma \longmapsto (\sigma_\Gamma, N_\Gamma, \rho = \mathbf{1}) \longmapsto (\widetilde{\phi}_\Gamma, N_\Gamma, \rho_\Gamma).$$

- **Infinitesimal character.** In the chosen coordinates, $\widetilde{\phi}_\Gamma = \widehat{\chi^v}$.
- **Monodromy.** The nilpotent entry N_Γ is the parameter's monodromy operator.
- **Enhancement.** The representation ρ_Γ is determined by inflation:

$$\rho = q^* \rho_\Gamma, \qquad q : \pi_0 Z_H(\sigma_\Gamma, N_\Gamma) \twoheadrightarrow \mathcal{R}_{\phi_\Gamma}.$$

References. AMS, *Affine Hecke algebras for Langlands parameters*;
Solleveld, *On principal series representations of quasi-split reductive p-adic groups*.

What the formula adds to the LLC dictionary

1. **An explicit constituent-by-constituent answer.**
   The inducing character fixes $\widetilde{\phi}$; chamber signs determine N; every enhancement is trivial.

2. **A criterion over the original field.**
   The socle has zero monodromy exactly when its chamber lies on every negative side.

3. **A uniform quasi-split formula.**
   It combines known chamber labels with the established principal-series LLC, including nonsplit and ramified cases.

Thank you.