
# 1 Programming Setup


#### 1.1 Setup Datacamp




#### 1.2 R

For installation process and HelloWorld example, please refer to the file located at "/homework/homework1/R\_code"

## 1.3 Debugger



### 1.4 Setup Github




### 2 Sign up NBER working paper series

- 1. The title of the second paper listed on the NBER weekly working paper series is "Why Survey-Based Subjective Expectations are Meaningful and Important", working paper number 32199
- 2. I have downloaded a paper interesting me most at the path "/homework/homework1/path" with title "Why Survey-Based Subjective Expectations are Meaningful and Important"

## 3 Sign up SRDA

- 1. The year of the PSFD dataset I downloaded is 2000
- 2. The working rate against age in the dataset on 2000 year is



# 4 Roy Model

#### 4.1 Review

1. We first consider  $\mathbb{E}[w_0|\text{Migrate}]$ , which can be expressed as

$$\mathbb{E}[w_0|\text{Migrate}] = \mu_0 + \mathbb{E}[\epsilon_0|w_1 > w_0 + C]$$

$$= \mu_0 + \mathbb{E}[\epsilon_0|\mu_1 + \epsilon_1 > \mu_0 + \epsilon_0 + C]$$

$$= \mu_0 + \mathbb{E}[\epsilon_0|\nu > \mu_0 - \mu_1 + C]$$

$$= \mu_0 + \mathbb{E}[\epsilon_0|\frac{\nu}{\sigma_\nu} > z]$$

$$= \mu_0 + \sigma_0 \mathbb{E}[\frac{\epsilon_0}{\sigma_0}|\frac{\nu}{\sigma_\nu} > z]$$

next, to compute the conditional expectation of  $\epsilon_0$ , we have to derive the following formulas

(a) Under normality, we have the regression coefficient formula

$$\mathbb{E}[\epsilon_0|\nu] = \frac{\sigma_{0,\nu}}{\sigma_{\nu}^2}\nu$$

(b) Use the formula after changing terms

$$\mathbb{E}\left[\frac{\epsilon_0}{\sigma_0}\middle|\frac{\nu}{\sigma_\nu}\right] = \frac{1}{\sigma_0} \underbrace{\frac{\sigma_{0,\nu}}{\sigma_\nu^2} \frac{\nu}{\sigma_\nu}}_{\text{by above formula}} \underbrace{\frac{1}{\sigma_0\sigma_\nu} \frac{1}{\sigma_\nu}}_{\text{by above}} = \underbrace{\frac{\sigma_{0,\nu}}{\sigma_0\sigma_\nu} \frac{\nu}{\sigma_\nu}}_{\text{formula}} = \rho_{0,\nu} \frac{\nu}{\sigma_\nu}$$
(1)

(c) From the preceding expectation formula, we can simplify the conditional expectation

$$\mathbb{E}[w_0|\text{Migrate}] = \mu_0 + \sigma_0 \mathbb{E}\left[\frac{\epsilon_0}{\sigma_0} \middle| \frac{\nu}{\sigma_\nu} > z\right]$$

$$\stackrel{(1)}{=} \mu_0 + \rho_{0,\nu} \sigma_0 \mathbb{E}\left(\frac{\nu}{\sigma_\nu} \middle| \frac{\nu}{\sigma_\nu} > z\right)$$

$$= \mu_0 + \rho_{0,\nu} \sigma_0 \frac{\phi(z)}{1 - \Phi(z)}$$

where  $\Phi(z)$  and  $\phi(z)$  are respectively defined as the pdf and cdf of z.

Besides, note that the coefficient of Inverse Mill Ratio can be rewritten as

$$\rho_{0,\nu}\sigma_0 = \underbrace{\frac{\sigma_{0,\nu}}{\sigma_{\nu}} = \frac{\sigma_{0,1} - \sigma_0^2}{\sigma_{\nu}}}_{\sigma_{0,\nu} = \text{Cov}(\epsilon_0,\epsilon_1 - \epsilon_0)} = \frac{\sigma_0\sigma_1}{\sigma_{\nu}} \left(\frac{\sigma_{0,1} - \sigma_0^2}{\sigma_0\sigma_1}\right) = \frac{\sigma_0\sigma_1}{\sigma_{\nu}} \left(\rho - \frac{\sigma_0}{\sigma_1}\right)$$

$$= \underbrace{\frac{\sigma_0\sigma_1}{\sigma_0\sigma_1} - \frac{\sigma_0\sigma_1}{\sigma_0\sigma_1}}_{\sigma_0\sigma_1} \left(\frac{\sigma_0\sigma_1 - \sigma_0^2}{\sigma_0\sigma_1}\right) = \frac{\sigma_0\sigma_1}{\sigma_{\nu}} \left(\frac{\sigma_0\sigma_1 - \sigma_0^2}{\sigma_0\sigma_1}\right)$$

Follow the same procedure, we can derive  $\mathbb{E}[w_1|\text{Migrate}]$  as well. Therefore, the conditional expectation can be expressed in a quite symmetric way

$$\begin{cases} \mathbb{E}[w_0|\text{Migrate}] = \mu_0 + \frac{\sigma_0 \sigma_1}{\sigma_\nu} (\rho - \frac{\sigma_0}{\sigma_1}) (\frac{\phi(z)}{1 - \Phi(z)}) \\ \mathbb{E}[w_1|\text{Migrate}] = \mu_1 + \frac{\sigma_0 \sigma_1}{\sigma_\nu} (\frac{\sigma_1}{\sigma_0} - \rho) (\frac{\phi(z)}{1 - \Phi(z)}) \end{cases}$$

2. If  $Q_0 > 0$  and  $Q_1 < 0$ , then we have

$$\begin{cases} \rho - \frac{\sigma_0}{\sigma_1} > 0 \\ \frac{\sigma_1}{\sigma_0} - \rho < 0 \end{cases} \Leftrightarrow \rho = \frac{\sigma_{0,1}}{\sigma_0 \sigma_1} > \max\{\frac{\sigma_0}{\sigma_1}, \frac{\sigma_1}{\sigma_0}\} \Leftrightarrow \sigma_{0,1} > \max\{\sigma_0^2, \sigma_1^2\} \end{cases}$$

which is impossible, hence such situation doesn't occur given people choose to migrate. In economics sense, this contradiction suggest that people would not leave for the lower proportion of  $w_1$  income distribution when they are in the upper proportion of  $w_0$  income distribution

#### 4.2 Simulation

1. First, let's pick some values for the parameters  $\{\mu_0, \mu_1, \sigma_0, \sigma_1, \sigma_{0,1}, C\}$ 

```
1 # 1. Pick my favorite value for set of parameters
2 mu_0 <- 100
3 mu_1 <- 100
4 sigma_0 <- 3
5 sigma_1 <- 6
6 sigma_01 <- 5
7 cost <- 50
8 n <- 1000000</pre>
```

2. After simulation, store the result of  $\{\epsilon_0, \epsilon_1\}$  in data.table

```
# 2. Simulate the epsilon_0 and epsilon_1
set.seed(123)
reps <- 1000000
par.est.DT <- data.table(matrix(0, reps, 6))
epsilon_0 <- rnorm(n, mean = 0, sd = sigma_0**2)
epsilon_1 <- rnorm(n, mean = 0, sd = sigma_1**2)</pre>
```

3. and create the columns for  $w_0$  and  $w_1$ 

```
# 3. Create the columns for w0 and w1
w_0 <- mu_0 + epsilon_0
w_1 <- mu_1 + epsilon_1
par.est.DT[, 1] <- epsilon_0
par.est.DT[, 2] <- epsilon_1
par.est.DT[, 3] <- w_0
par.est.DT[, 4] <- w_1</pre>
```

4. also, generate a column I that takes binary value

```
# 4. Generate the column I that take binary value.

2 par.est.DT[, 5] <- ifelse(par.est.DT[, 2] > par.est.DT[, 1], 1, 0)

3
```

5. Then compute  $\mathbb{E}[w_0|\mathbb{I}], \mathbb{E}[w_1|\mathbb{I}], Q_0, Q_1$  from data

```
1 # 5. Calculate E[w0|I], E[w1|I], Q0, Q1 from data
2 E_epsilon0_D1 <- par.est.DT[V5 == 1, mean(V1)]
3 E_epsilon1_D1 <- par.est.DT[V5 == 1, mean(V2)]
4 E_w0_D1 <- par.est.DT[V5 == 1, mean(V3)]
5 E_w1_D1 <- par.est.DT[V5 == 1, mean(V4)]
6</pre>
```

6. Use equation (1) and (2) to compute conditional expectation of  $w_0$  and  $w_1$ 

```
# 6. Calculate RHS of equation (1) and (2)
nu <- epsilon_1 - epsilon_0
par.est.DT[, 6] <- nu
sigma_nu <- par.est.DT[,sd(V5)]
Z <- (mu_0 - mu_1 + cost)/sigma_nu
mill <- function(x) {
pnorm(x, lower.tail=FALSE, log.p=TRUE) - dnorm(x, log=TRUE)
}</pre>
```

7. The column  $\mathbb{I}$  and  $\mathbb{E}[w_1|\mathbb{I}=1]$  are observable in the real world, but the other variable  $\{\epsilon_0, \epsilon_1, Q_0, Q_1, \phi(Z), \Phi(Z)\}$  and  $\mathbb{E}[w_0|\mathbb{I}=1]$  are unobservable

## 5 Roy Model is Everywhere

- 1. An example that immediately comes to my mind is getting master (or ph.D) degree or not. The time spent on studying a degree or the risk of not finishing it are costly, nonetheless, the degree itself is also desirable when seeking a job. It might need argue that  $Q_0 < 0$  or  $Q_0 > 0$ , but I think most people will agree on  $Q_1 > 0$ .
- 2. Consider the Roy model

$$\begin{cases} w_0 = \mu_0 + \epsilon_0 \\ w_1 = \mu_1 + \epsilon_1 \end{cases}$$

where studying a higer degree represents 1, while not studying represents 0. It's assumed that  $\epsilon_i \sim \mathcal{N}(0, \sigma_i^2)$ , where  $i \in \{0, 1\}$ . Note that the pain or time studying takes are capture by constant cost C and one choose studying higher degree only if  $\mathbb{E}[w_1] - C > \mathbb{E}[w_0]$ .