

Discrete Mathematics: Lecture 23

Part IV. Graph Theory

Hamilton Paths and Circuits, Shortest Paths and Dijkstra's Algorithm, Traveling Salesperson Problem

Xuming He





Associate Professor

School of Information Science and Technology
ShanghaiTech University

Spring Semester, 2025


Notes by Prof. Liangfeng Zhang

Q1 Which of the following graphs is bipartite but not complete bipartite?



Q2

What is the vertex connectivity of the graph pictured below?



A. 1

B. 2


C. 3

D. 4


Hamilton Paths and Circuits

DEFINITION: Let $G = (V, E)$ be a graph.


- **Hamilton Path:** A **simple path** that passes through every vertex exactly once.
- **Hamilton Circuit:** A **simple circuit** that passes through every vertex exactly once.




✓ Hamilton path
✓ Hamilton circuit



✓ Hamilton path
✓ Hamilton circuit



✓ Hamilton path
× Hamilton circuit



× Hamilton path
× Hamilton circuit

Hamilton Circuits

Determine if there is a Hamilton circuit in a given graph G ?

- This problem is NP-Complete. //that means very difficult

Necessary conditions on Hamilton circuit.



- If G has a vertex of degree 1, then G cannot have a Hamilton circuit.
- If G has a vertex of degree 2, then a Hamilton circuit of G traverses both edges.

Sufficient conditions on Hamilton circuit.





- **Ore's Theorem:** Let $G = (V, E)$ be a simple graph of order $n \geq 3$. If $\deg(u) + \deg(v) \geq n$ for all $\{u, v\} \notin E$, then G has a Hamilton circuit.
- **Dirac's Theorem:** Let $G = (V, E)$ be a simple graph of order $n \geq 3$. If $\deg(u) \geq n/2$ for every $u \in V$, then G has a Hamilton circuit.
 - This is a corollary of Ore's Theorem
 - $\forall u \in V, \deg(u) \geq n/2 \Rightarrow \forall u, v \in V, \deg(u) + \deg(v) \geq n$

Hamilton Circuits


- Examples (sufficient condition)



Hamilton Circuits



Remark: Dirac's and Ore's Theorems do not give a necessary condition for the existence of a Hamilton circuit!



Position of a rotating pointer encoded by a bit string of length n

Gray code: Labeling of the arcs of the circle such that adjacent arcs are labeled with bit string that differ exactly in one bit.

\Rightarrow Hamilton cycle in Q_n .





FIGURE 14 A Hamilton Circuit for Q_3 .

Shortest Path Problem

Definition

A **weighted graph** is a graph $G = (V, E)$ such that each edge is assigned with a strictly positive number.

The **length** of a path in weighted graph is the sum of the weights of the edges of this path.




a, b, c is a path of length 7 and b, e, d, c is a path of length 6


Remark: Observe that in a non-weighted graph the length of a path is the number of edges in the path!

Shortest Path Problem

Examples



distance




time

What is the shortest path in air distance between cities A and E?
What combination of flights has the smallest total flight time?

Shortest Path Problem


Question: Find the shortest path from a to d .



Method: Find the closest vertex to a , then the second closest, the third closest... until we reach d .

⇒ Dijkstra's algorithm

Dijkstra's Algorithm



- 1 Find the closest vertex to $a \rightsquigarrow$ analyse all the edges starting from a :
 a, b of length 4
 a, f of length 2
 $\Rightarrow f$ is the closest vertex to a . The shortest path from a to f has length 2.
- 2 Find the second closest vertex to $a \rightsquigarrow$ shortest paths from a to a vertex in $\{a, f\}$ followed by an edge from a vertex in $\{a, f\}$ to a vertex not in this set:
 a, b of length 4
 a, f, e of length 5
 $\Rightarrow b$ is the second closest vertex to a . The shortest path from a to b has length 4.

Dijkstra's Algorithm



- 3** Find the third closest vertex to $a \rightsquigarrow$ shortest path from a to a vertex in $\{a, f, b\}$ followed by an edge from a vertex in $\{a, f, b\}$ to a vertex not in this set:

a, b, c of length 7

a, b, e of length 7

a, f, e of length 5

$\Rightarrow e$ is the third closest vertex to a . The shortest path from a to e has length 5.

- 4** Find the fourth closest vertex to $a \rightsquigarrow$ shortest path from a to a vertex in $\{a, f, b, e\}$ followed by an edge from a vertex in $\{a, f, b, e\}$ to a vertex not in this set:

a, b, c of length 7


a, f, e, d of length 6

$\Rightarrow d$ is the fourth closest vertex to a . The shortest path from a to d has length 6.

Shortest Path Problem


Remarks:

- Of course in the example above, we could have looked at all the paths between a and d and compute their length, but too complicated if the graph has a lot of edges.



- Advantage of Dijkstra's algorithm: we can compute the length of a shortest path from one vertex to all other vertices of the graph.

Dijkstra's Algorithm



Goal: find the length of a shortest path from a to z with a series of iterations.

- A distinguished set of vertices is constructed by adding one vertex at each iteration.
- A labeling procedure is carried out at each iteration: a vertex w is labeled with the length of a shortest path from a to w that contains only vertices in the distinguished set.
- The vertex added to the distinguished set is one with minimal label among those vertices not already in the set.

Dijkstra's Algorithm


Notations: S_k := distinguished set after k iterations, $L_k(v)$:= length of a shortest path from a to v containing only vertices in S_k ("label" of v).

Initialization: $L_0(a) = 0$,
 $L_0(v) = \infty$ for every vertex $v \neq a$,
 $S_0 = \emptyset$.


k th iteration:

- S_k is formed from S_{k-1} by adding a vertex u not in S_{k-1} with smallest label,
- Update the labels of all vertices not in S_k so that $L_k(v)$ is the length of a shortest path from a to v containing only vertices in S_k , i.e.

$L_k(v) = \min\{L_{k-1}(v), L_{k-1}(u) + w(u, v)\}$ (with $w(u, v)$ length of the edge (u, v))



Dijkstra's Algorithm



- **k=0 (initialization):** $S_0 = \emptyset$,
 $L_0(a) = 0$, $L_0(b) = L_0(c) =$
 $L_0(d) = L_0(e) = L_0(z) = \infty$

2


3

4

5

6 $\{a, c, b, d, e, z\}$


Dijkstra's Algorithm



- **k=0 (initialization):** $S_0 = \emptyset$,
 $L_0(a) = 0$, $L_0(b) = L_0(c) =$
 $L_0(d) = L_0(e) = L_0(z) = \infty$

- **k=1:** $u := a \rightsquigarrow S_1 = \{a\}$,
 $L_0(a) + w(a, b) = 4 < L_0(b) \rightsquigarrow L_1(b) = 4$
 $L_0(a) + w(a, c) = 2 < L_0(c) \rightsquigarrow L_1(c) = 2$
- **k=2:** $u := c \rightsquigarrow S_2 = \{a, c\}$,
 $L_1(c) + w(c, b) = 3 < L_1(b) \rightsquigarrow L_2(b) = 3$
 $L_1(c) + w(c, d) = 10 < L_1(d) \rightsquigarrow L_2(d) = 10$
 $L_1(c) + w(c, e) = 12 < L_1(e) \rightsquigarrow L_2(e) = 12$
- **k=3:** $u := b \rightsquigarrow S_3 = \{a, c, b\}$,
 $L_2(b) + w(b, d) = 8 < L_2(d) \rightsquigarrow L_3(d) = 8$
- **k=4:** $u := d \rightsquigarrow S_4 = \{a, c, b, d\}$,
 $L_3(d) + w(d, e) = 10 < L_3(e) \rightsquigarrow L_4(e) = 10$
 $L_3(d) + w(d, z) = 14 < L_3(z) \rightsquigarrow L_4(z) = 14$
- **k=5:** $u := e \rightsquigarrow S_5 = \{a, c, b, d, e\}$,
 $L_4(e) + w(e, z) = 13 < L_4(z) \rightsquigarrow L_5(z) = 13$
- **k=6:** $u := z \rightsquigarrow S_6 = \{a, c, b, d, e, z\}$
- **return:** $L(z) = 13$


Traveling Salesperson Problem



Traveling salesperson problem: a traveling salesperson wants to visit each of the cities once and return to his starting point. In which order should he visit these cities to travel the minimum total distance?

⇒ **Hamiltonian circuit with minimum total weight in the complete graph.**

Traveling Salesperson Problem



Route	Tot. dist.
A, B, C, D, E, A	610
A, B, C, E, D, A	516
A, B, E, D, C, A	588
A, B, E, C, D, A	458
A, B, D, E, C, A	540
A, B, D, C, E, A	504
A, D, B, C, E, A	598
A, D, B, E, C, A	576
A, D, E, B, C, A	682
A, D, C, B, E, A	646
A, C, D, B, E, A	670
A, C, B, D, E, A	728



Traveling salesperson problem: a traveling salesperson wants to visit each of the cities once and return to his starting point. In which order should he visit these cities to travel the minimum total distance?

⇒ **Hamiltonian circuit with minimum total weight in the complete graph.**

Planar Graph

DEFINITION: Let $G = (V, E)$ be an undirected graph. G is called a **planar graph** 平面图 if it can be drawn in the plane without any edges crossing.

- Crossing of edges: an intersection other than endpoints (vertices)
- **planar representation** 平面表示: a drawing w/o edge crossing; **nonplanar** 非平面的



- K_1, K_2, K_3, K_4 are planar graphs
- $K_{1,n}, K_{2,n}$ are planar graphs
- C_n ($n \geq 3$), W_n ($n \geq 3$) are planar graphs
- Q_1, Q_2, Q_3 are planar graphs

