

Discrete Mathematics: Lecture 24

Part IV. Graph Theory

Planar Graph, Euler's Formula, Homeomorphic, Kuratowski's Theorem

Xuming He

Associate Professor

School of Information Science and Technology

ShanghaiTech University



Spring Semester, 2025

Notes by Prof. Liangfeng Zhang


Planar Graph

DEFINITION: Let $G = (V, E)$ be an undirected graph. G is called a **planar graph** 平面图 if it can be drawn in the plane without any edges crossing.

- Crossing of edges: an intersection other than endpoints (vertices)
- **planar representation** 平面表示: a drawing w/o edge crossing; **nonplanar** 非平面的




- K_1, K_2, K_3, K_4 are planar graphs
- $K_{1,n}, K_{2,n}$ are planar graphs
- C_n ($n \geq 3$), W_n ($n \geq 3$) are planar graphs
- Q_1, Q_2, Q_3 are planar graphs



Planar Graph

Examples






A graph may be planar even if it is usually drawn with crossings, because it may be possible to draw it in a different way without crossings.

Nonplanar Graph

Jordan Curve Theorem: Every simple closed planar curve Γ separates the plane into a bounded interior region and an unbounded exterior region. Any planar curve connecting the two regions must intersect Γ .


EXAMPLE: The bipartite graph $K_{3,3}$ is not planar.



- choose a simple circuit v_1, v_5, v_2, v_4, v_1 in $K_{3,3}$
- If $K_{3,3}$ is a planar, then the circuit forms a simple closed planar curve
- Add v_3, v_6 and the edges incident with them.
 - Intersection occurs (due to the Jordan curve Theorem).

Nonplanar Graph

EXAMPLE: The complete graph K_5 is not planar.




- $v_1, v_2, v_3, v_4, v_5, v_1$ is a simple closed curve in the planar representation of K_5
- Every remaining edge is in the interior region or in the exterior region
 - at least one is in the interior region
- No matter how you draw the remaining edges, crossing occurs.

Regions

DEFINITION: Let $G = (V, E)$ be a planar graph. Then the plane is divided into several **regions**面 by the edges of G .

- The infinite region is **exterior region**外部面. The others are **interior regions**内部面.
- The **boundary**边界 of a region is a subset of E .
- The **degree**度数 of a region is the number of edges on its boundary.
 - If an edge is shared by R_i, R_j , then it contributes 1 to $\deg(R_i), \deg(R_j)$
 - If an edge is on the boundary of a single region R_i , then it contributes 2 to $\deg(R_i)$



- The plane is divided into 5 regions R_0, R_1, R_2, R_3, R_4
 - R_0 is the exterior region
 - R_1, R_2, R_3, R_4 are interior regions
- The boundary of R_1 ; $\deg(R_1) = 4$
- There are 4 edges on the boundary of R_4
 - $\deg(R_4) = 1 + 1 + 1 + 2 = 5$ because one of the edges contribute 2 to $\deg(R_4)$
- $\deg(R_0) = 11, \deg(R_1) = 4, \deg(R_2) = 3, \deg(R_3) = 3, \deg(R_4) = 5$

Euler's Formula

THEOREM: Let $G = (V, E)$ be a connected planar simple graph with e edges and v vertices. Let r be the number of regions in a planar representation of G . Then $r = e - v + 2$.

THEOREM: Let G be a planar simple graph with p connected components. Then $|V(G)| - |E(G)| + |R(G)| = p + 1$.

- Let G_1, G_2, \dots, G_p be the connected components of G .
 - By Euler's formula, $|R(G_i)| = |E(G_i)| - |V(G_i)| + 2$ for all $i \in [p]$
- $|V(G)| = |V(G_1)| + |V(G_2)| + \dots + |V(G_p)|$
- $|E(G)| = |E(G_1)| + |E(G_2)| + \dots + |E(G_p)|$
- $|R(G)| = |R(G_1)| + |R(G_2)| + \dots + |R(G_p)| - p + 1$
- $|V(G)| - |E(G)| + |R(G)| = \sum_{i=1}^p (|V(G_i)| - |E(G_i)| + |R(G_i)|) - p + 1$
 $= 2p - p + 1 = p + 1$

Euler's Formula: Proof

Proof of Euler's formula by induction on the number e of edges

- A simple connected planar graph with 0 edges has only one vertex and one face (unbounded). The relation $f = e - v + 2$ is satisfied.
- Suppose the relation is satisfied for all simple connected planar graphs with k edges.

Consider a simple connected planar graph G with $k + 1$ edges, $k \geq 0$. This graph can be seen as a simple connected planar graph G' with k edges (satisfying the relation by induction hypothesis) to which we add one edge. There are two ways to add an edge to G' to get G :

- either the two endpoints of the edge are already in G' : in this case, adding the edge adds also one face,
- either only one of the endpoint is already in G' : in this case, adding the edge adds also one vertex but no other face.

In both cases, the relation $f = e - v + 2$ is satisfied by G .

Application

THEOREM: Let G be a **connected planar simple graph**. If every region has degree $\geq l$ in a planar representation of G , then

then $|E(G)| \leq \frac{l}{l-2} (|V(G)| - 2)$.

- Let R_1, \dots, R_t be the regions given by a planar representation of G // $t = |R(G)|$
 - $\deg(R_i) \geq l$ for every $i = 1, 2, \dots, t$
- Let $r = \deg(R_1) + \deg(R_2) + \dots + \deg(R_t)$. Then $r = 2|E(G)|$.
 - Every edge contributes 2 to r
 - If $e \in E$ is on the boundary of a single region R_i , then e contributes 2 to $\deg(R_i)$;
 - If $e \in E$ is shared by R_i and R_j , then e contributes 1 to $\deg(R_i)$ and 1 to $\deg(R_j)$;
- $2|E(G)| = r = \deg(R_1) + \deg(R_2) + \dots + \deg(R_t) \geq lt = l|R(G)|$
- $|R(G)| = |E(G)| - |V(G)| + 2$
- Hence, $|E(G)| \leq \frac{l}{l-2} (|V(G)| - 2)$

Application

COROLLARY: Let G be a **connected planar simple graph**. If $|V(G)| \geq 3$, then $|E(G)| \leq 3|V(G)| - 6$.

- Every region has degree ≥ 3 in a planar representation of G
- Let $l = 3$ in the previous theorem
 - $|E(G)| \leq \frac{3}{3-2} (|V(G)| - 2) = 3|V(G)| - 6$.

EXAMPLE: The complete graph K_5 is not planar.

- $|E(K_5)| = \binom{5}{2} = 10, |V(K_5)| = 5, K_5$ is connected simple and of order ≥ 3
- $|E(K_5)| > 3|V(K_5)| - 6$
 - Hence, K_5 cannot be planar

COROLLARY: Let G be a connected planar simple graph. Then G has a vertex of degree ≤ 5 .

- $|V(G)| < 3$: the statement is true.
- $|V(G)| \geq 3: \forall u \in V(G), \deg(u) \geq 6 \Rightarrow 2|E(G)| = \sum_u \deg(u) \geq 6|V(G)|$
 - G cannot be planar


Application

COROLLARY: Let G be a **connected planar simple graph**. If $|V(G)| \geq 3$ and there is no circuits of length 3 in G , then $|E(G)| \leq 2|V(G)| - 4$.

- Let R_1, \dots, R_t be the regions given by a planar representation of G // $t = |R(G)|$
 - $\deg(R_i) \geq 4$ for every $i = 1, 2, \dots, t$
- Hence, $|E(G)| \leq \frac{4}{4-2} (|V(G)| - 2) = 2|V(G)| - 4$

EXAMPLE: The complete bipartite graph $K_{3,3}$ is not planar.

- $|E(K_{3,3})| = 3 \times 3 = 9, |V(K_{3,3})| = 3 + 3 = 6 \geq 3$
- $K_{3,3}$ is connected, simple and of order ≥ 3 .
- There is no circuits of length 3 in $K_{3,3}$
- $|E(K_{3,3})| = 9 > 8 = 2|V(K_{3,3})| - 4$
- Hence, $K_{3,3}$ cannot be planar




REMARKS: K_5 and $K_{3,3}$ are fundamental nonplanar graphs.

Homeomorphic

DEFINITION: Let $G = (V, E)$ be a graph and $\{u, v\} \in E$.

- **elementary subdivision** 初等细分: $G' = (V \cup \{w\}, E - \{u, v\} + \{u, w\} + \{v, w\})$
- Two graphs are **homeomorphic** 同胚的 if they can be obtained from the same graph via elementary subdivisions






G_2 and G_3 are homeomorphic




Kuratowski's Theorem

THEOREM: A graph G is nonplanar if and only if it has a subgraph homeomorphic to $K_{3,3}$ or K_5 .

EXAMPLE: The following graph is nonplanar.




There is a subgraph
homeomorphic to K_5



There is a subgraph
homeomorphic to $K_{3,3}$

Dual Graph



Let G be a planar graph and assume we take a planar representation of G that we denote also G . The **dual of G** is the graph G^* that has a vertex for each face of G and an edge connecting two vertices if the corresponding faces in G have a common edge in their boundary.



Remark: The dual of a planar simple graph is not necessarily simple.

Dual Graph


- Properties of dual graph
 - The dual G^* of a planar graph G is a planar graph.
 - The dual of a planar graph is always connected.
 - The graphs G_1 and G_2 below are isomorphic. What about their duals?



Dual Graph

- Properties of dual graph

- The graphs G_1 and G_2 below are isomorphic. What about their duals?



⇒ The dual of a planar graph depends on the planar representation of the graph.

Dual Graph

- Properties of dual graph
 - If G is a planar connected graph, then $v^* = f$, $e^* = e$, and $f^* = v$
 - If G is a planar connected graph then $G^{**} = G$ (not true if the graph is disconnected!)

Dual Graph

- Properties of dual graph

- If G is a planar connected graph, then $v^* = f$, $e^* = e$, and $f^* = v$
- If G is a planar connected graph then $G^{**} = G$ (not true if the graph is disconnected!)

Definition

A planar graph is said **self-dual** if it is isomorphic to its dual.

Example: The wheels W_n are self-dual graphs.






Proposition

A self-dual graph with v vertices has $2v - 2$ edges.

Proof: We have $v^* = v = f^* = f$ because the graph is self-dual. By Euler's formula

$$f = e - v + 2 \Rightarrow v = e - v + 2 \Rightarrow 2v - 2 = e$$

Coloring a Map



Coloring regions of the map \Leftrightarrow Coloring vertices of the dual graph