Econometrics Guide

Xiang Ao

2026-10-17

Table of contents

Preface **1**
How to cite . . . 1

1 OLS **3**
1.1 Fitness of OLS . . . 3
1.2 Example . . . 4
1.3 The Geometry of Least Squares . . . 5
1.4 OLS as Conditional Expectation . . . 6
1.5 Properties of OLS estimator . . . 8
1.6 Summary . . . 10

2 The Method of Maximum Likelihood **11**
2.1 Basic Concepts . . . 11
2.2 MLE for a linear model . . . 11
2.3 Asymptotic Properties of MLE . . . 12
2.4 When the Model Is Wrong: Quasi-MLE and M-Estimation . . . 13
2.5 Example . . . 15
2.6 Hypotheses Testing . . . 16
2.7 Test of Structural Change (Chow test) . . . 18
2.8 Heteroskedasticity and Autocorrelation Consistent Standard Errors . . . 22
2.9 Summary . . . 25

3 GLS **27**
3.1 Introduction . . . 27
3.2 The GLS Estimator . . . 27
3.3 Weighted Least Squares . . . 28
3.4 Efficiency of GLS . . . 29
3.5 Summary . . . 30

4 Endogeneity **31**
4.1 Instrumental Variables (IV) . . . 31
4.2 Control Function Approach . . . 36
4.3 Durbin-Wu-Hausman Test . . . 38
4.4 Identification . . . 39
4.5 Weak Instruments . . . 40
4.6 When is endogeneity a problem and what can we say from an IV regression . . . 43
4.7 Summary . . . 44

5 Generalized Method of Moments (GMM) **45**
5.1 The Method of Moments (MOM) . . . 45
5.2 OLS as a moment problem . . . 45

5.3 IV as a moment problem . . . 46
5.4 The Generalized Method of Moments . . . 46
5.5 GMM . . . 47
5.6 A few concepts of conditioning . . . 53
5.7 Summary . . . 55

6 Censored, Truncated or Selected Data **57**
6.1 Compare three different cases . . . 57
6.2 An illustration for truncated or censored data . . . 58
6.3 Truncated Models . . . 58
6.4 Censored Models . . . 59
6.5 Sample Selection . . . 60
6.6 Switching Regression (Treatment-Effects Model) . . . 62
6.7 Summary . . . 62

7 Discrete and Limited Dependent Variables **65**
7.1 Binary Response Models . . . 65
7.2 Models for More than Two Discrete Responses . . . 66
7.3 Summary . . . 69

8 Count Data Models **71**
8.1 Poisson Model . . . 71
8.2 QMLE Poisson . . . 72
8.3 Fixed-effect Poisson model . . . 73
8.4 Negative Binomial model . . . 73
8.5 Models for truncated counts . . . 74
8.6 The hurdle regression model . . . 74
8.7 Zero-inflated count models . . . 75
8.8 An example of extra-zero count data model comparison . . . 76
8.9 Interpreting Coefficients and Marginal Effects in Nonlinear Models . . . 76
8.10 Summary . . . 79

9 Panel Data Models **81**
9.1 Background . . . 81
9.2 Random Effect Methods . . . 82
9.3 Fixed Effect Methods . . . 83
9.4 Fixed effects or random effects: the Hausman test . . . 84
9.5 Collapse to unit means, or keep the long format? . . . 84
9.6 What the long format adds . . . 85
9.7 Summary . . . 88

10 Survival Models **89**
10.1 Survival function . . . 89
10.2 Hazard function . . . 90
10.3 Log-likelihood function . . . 91
10.4 Proportional Hazard Models . . . 92
10.5 Discrete-time Survival Models . . . 96
10.6 Summary . . . 97

11 Dynamic Panel Data **99**
11.1 When Is It a Problem . . . 99
11.2 How Big Is the Bias . . . 100
11.3 Anderson and Hsiao estimator . . . 100
11.4 Arellano-Bond estimator . . . 101
11.5 Blundell and Bond estimator . . . 101
11.6 Diagnostics in practice . . . 102
11.7 Summary . . . 103

12 Missing Data **105**
12.1 Different cases under different assumptions . . . 105
12.2 Old methods . . . 106
12.3 Modern methods . . . 106
12.4 Summary . . . 110

13 Interpreting OLS: Effect Weights **111**
13.1 One formula . . . 111
13.2 Binary treatment: OLS weights overlap . . . 117
13.3 Smaller groups get larger weights . . . 118
13.4 Fixed effects: OLS weights movers . . . 121
13.5 When does the weighting change the answer? . . . 126
13.6 Start with the estimand . . . 128
13.7 Summary . . . 129

14 Interpreting OLS: Outcome Weights **131**
14.1 The other reading of FWL . . . 131
14.2 What the outcome weights look like . . . 132
14.3 When the sign flips . . . 134
14.4 Out of OLS: the same weights beyond the linear model . . . 137
14.5 The DiD setting . . . 142
14.6 Summary . . . 160

References **161**

Preface

The book covers the core estimation and inference methods used in applied econometrics. It begins with OLS — matrix derivation, the conditional expectation function, and the best linear predictor — then moves to maximum likelihood, quasi-MLE and M-estimation, and generalized least squares. Endogeneity, instrumental variables, and GMM follow. The second half treats models for specific data structures: censored and truncated data, discrete choice, count data, panel data, dynamic panels, survival analysis, and missing data. Two closing chapters develop the modern interpretation of OLS under heterogeneous treatment effects: effect weights and outcome weights.

Examples are in R or Stata, whichever has the cleaner implementation for the procedure at hand. For applied causal-inference material — difference-in-differences, matching, doubly robust methods, mediation, and causal discovery — see the companion books Introduction to Causal Econometrics and Topics on Econometrics and Causal Inference.

How to cite

Ao, Xiang. *Econometrics Guide*. https://xiangao.github.io/econometrics-guide/.

```
@book{ao_econometrics_guide,
  author = {Ao, Xiang},
  title  = {Econometrics Guide},
  url    = {https://xiangao.github.io/econometrics-guide/}
}
```

1 OLS

A linear model with n observations and k regressors can be written as (in vector forms)

$$\mathbf{y} = \mathbf{X}\boldsymbol{\beta} + \mathbf{u} \tag{1.1}$$

The idea of least-squares is to choose β to minimize the residual sum of squares (SSR),

$$SSR(\beta) = (\mathbf{y} - \mathbf{X}\boldsymbol{\beta})'(\mathbf{y} - \mathbf{X}\boldsymbol{\beta}) = \mathbf{y}'\mathbf{y} - \mathbf{2}\boldsymbol{\beta}'\mathbf{X}'\mathbf{y} + \boldsymbol{\beta}'\mathbf{X}'\mathbf{X}\boldsymbol{\beta} \tag{1.2}$$

The first-order condition to minimize SSR are:

$$\frac{\partial(SSR)}{\partial\beta} = -\mathbf{2}\mathbf{X}'\mathbf{y} + \mathbf{2}\mathbf{X}'\mathbf{X}\boldsymbol{\beta} = \mathbf{0} \tag{1.3}$$

This generates the so-called normal equations

$$(\mathbf{X}'\mathbf{X})\boldsymbol{\beta} = \mathbf{X}'\mathbf{y} \tag{1.4}$$

Therefore, Ordinary Least Squares (OLS) estimator of β is

$$\hat{\boldsymbol{\beta}} = (\mathbf{X}'\mathbf{X})^{-\mathbf{1}}\mathbf{X}'\mathbf{y} \tag{1.5}$$

1.1 Fitness of OLS

We can see y vector as the part explained by the regression and the unexplained (residual) part,

$$\mathbf{y} = \hat{\mathbf{y}} + \hat{\mathbf{u}} = \mathbf{X}\hat{\boldsymbol{\beta}} + \hat{\mathbf{u}} \tag{1.6}$$

where $\hat{u}$ is the OLS residual. Because the residual is orthogonal to the fitted values, $\hat{\mathbf{y}}'\hat{\mathbf{u}} = \mathbf{0}$, so the cross term drops:

$$\mathbf{y}'\mathbf{y} = (\hat{\mathbf{y}} + \hat{\mathbf{u}})'(\hat{\mathbf{y}} + \hat{\mathbf{u}}) = \hat{\mathbf{y}}'\hat{\mathbf{y}} + \hat{\mathbf{u}}'\hat{\mathbf{u}} = \hat{\boldsymbol{\beta}}'\mathbf{X}'\mathbf{X}\hat{\boldsymbol{\beta}} + \hat{\mathbf{u}}'\hat{\mathbf{u}} \tag{1.7}$$

Subtracting $n\bar{y}^2$ ($\bar{y}$ is the sample mean) from both sides, **and assuming the regression includes an intercept** (so that $\bar{\hat{y}} = \bar{y}$ and the residuals sum to zero, which is what makes this centered decomposition and the R^2 interpretation below hold),

$$\mathbf{y}'\mathbf{y} - \mathbf{n}\bar{\mathbf{y}}^{\mathbf{2}} = (\hat{\boldsymbol{\beta}}'\mathbf{X}'\mathbf{X}\hat{\boldsymbol{\beta}} - \mathbf{n}\bar{\mathbf{y}}^{\mathbf{2}}) + \hat{\mathbf{u}}'\hat{\mathbf{u}} \tag{1.8}$$

We decompose the total sum of squares into two parts: sum of squares due to error (noise), and sum of squares explained by the linear regression.

The R^2 is defined by

$$R^2 = 1 - \frac{SSR}{SST} \tag{1.9}$$

SST is the total (centered) sum of squares of y, $\mathbf{y'y - n\bar{y}^2} = \sum_{\mathbf{t}} (\mathbf{y_t - \bar{y}})^{\mathbf{2}}$.

R^2 is simply the proportion of the variation of Y that can be attributed to the variation of X. R^2, however, will never decrease with the addition of any variable to the set of regressors. R^2 stays exactly the same only in the knife-edge case that the added variable is orthogonal to the current residuals; in practice it always rises a little, even for an irrelevant variable. The adjusted R^2, however, takes account of the addition of any new regressors:

$$\bar{R}^2 = 1 - \frac{SSR/(n-k)}{SST/(n-1)} \tag{1.10}$$

1.2 Example

Let's look at an example of OLS of car's mpg on disp, hp and wt.

```
lm1 <- lm(mpg~disp+hp+wt, data=mtcars)
summary(lm1)
```


```
Call:
lm(formula = mpg ~ disp + hp + wt, data = mtcars)

Residuals:
   Min     1Q Median     3Q    Max
-3.891 -1.640 -0.172  1.061  5.861

Coefficients:
             Estimate Std. Error t value Pr(>|t|)
(Intercept) 37.105505   2.110815  17.579  < 2e-16 ***
disp        -0.000937   0.010350  -0.091  0.92851
hp          -0.031157   0.011436  -2.724  0.01097 *
wt          -3.800891   1.066191  -3.565  0.00133 **
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 2.639 on 28 degrees of freedom
Multiple R-squared:  0.8268,    Adjusted R-squared:  0.8083
F-statistic: 44.57 on 3 and 28 DF,  p-value: 8.65e-11
```

1.3 The Geometry of Least Squares



For simplicity, let's assume there are two explanatory variables x_1 and x_2. x_1 and x_2 form a plane. What OLS does is to project y onto this plane.

In mathematical terms, all linear combinations of these two vectors define a two-dimensional subspace of Euclidean space $\mathbf{E}^n$. This is called the column space of $\mathbf{X}$. The least-square principle is to choose β to make $\hat{y}$, which belongs to the subspace of $\mathbf{X}$, as close as possible to y.

When we estimate a linear regression model, we map the y into a vector of fitted values $\mathbf{X}\hat{\boldsymbol{\beta}}$ and a vector of residuals $\hat{\mathbf{u}} = \mathbf{y} - \mathbf{X}\hat{\boldsymbol{\beta}}$. Geometrically, these are examples of orthogonal projections. A projection is a mapping that takes each point of $\mathbf{E}^n$ into a point in a subspace of $\mathbf{E}^n$. An orthogonal projection maps any point into the point of the subspace that is closest to it.

An orthogonal projection can be performed by premultiplying the vector to be projected by a projection matrix. In the case of OLS, the two projection matrices that yield the vector of fitted values and the vector of residuals, are

$$\mathbf{P}_\mathbf{X} = \mathbf{X}(\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}' \tag{1.11}$$

$$\mathbf{M}_\mathbf{X} = \mathbf{I} - \mathbf{P}_\mathbf{X} = \mathbf{I} - \mathbf{X}(\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}' \tag{1.12}$$

From this, we see that the effects of two projection matrices.

$$\mathbf{P_X y} = \mathbf{X}\hat{\boldsymbol{\beta}} = \hat{\mathbf{y}} \tag{1.13}$$

$$\mathbf{M_X y} = (\mathbf{I} - \mathbf{P_X})\mathbf{y} = \mathbf{y} - \mathbf{P_X y} = \mathbf{y} - \mathbf{X}\hat{\boldsymbol{\beta}} = \hat{\mathbf{u}} \tag{1.14}$$

That is, $\mathbf{P_X y}$ projects $\mathbf{y}$ onto $\mathbf{X}$ and makes it $\hat{\mathbf{y}}$; $\mathbf{M_X y}$ makes it $\hat{\mathbf{u}}$.

In the picture, that corresponds to the fact that the projection of $\mathbf{y}$ onto the plane of $\mathbf{X}$ creates two parts: $\hat{\mathbf{y}}$ and $\hat{\mathbf{u}}$.

1.4 OLS as Conditional Expectation

The derivation above shows how to compute $\hat{\boldsymbol{\beta}}$ from a sample. We now ask a different question: what does OLS estimate at the population level?

1.4.1 The Conditional Expectation Function

For a random pair (Y, X), the conditional expectation function (CEF) is $E[Y \mid X]$, a function of X alone. Any random variable Y can be decomposed as

$$Y = E[Y \mid X] + \varepsilon, \qquad E[\varepsilon \mid X] = 0. \tag{1.15}$$

This is not an assumption; it is a definition. The residual $\varepsilon = Y - E[Y \mid X]$ is mean-independent of X by the construction of conditional expectation.

The CEF has an optimality property: among all functions $g(X)$, $E[Y \mid X]$ minimizes the mean squared prediction error,

$$E[Y \mid X] = \arg\min_{g} \; E\big[(Y - g(X))^2\big]. \tag{1.16}$$

That is, if we could use any function of X to predict Y, the CEF would be the best predictor in the mean squared error sense. The proof follows from the decomposition above: for any g,

$$E\big[(Y - g(X))^2\big] = E\big[(Y - E[Y \mid X])^2\big] + E\big[(E[Y \mid X] - g(X))^2\big] + 2\,E\big[\varepsilon\,(E[Y \mid X] - g(X))\big]. \tag{1.17}$$

The cross term vanishes by iterated expectations: $E[Y \mid X] - g(X)$ is a function of X, so $E\big[\varepsilon\,(E[Y \mid X] - g(X))\big] = E\big[E[\varepsilon \mid X]\,(E[Y \mid X] - g(X))\big] = 0$. The first term does not depend on g, and the second is nonnegative and equals zero when $g = E[Y \mid X]$.

1.4.2 The Best Linear Predictor

In practice we rarely know the CEF, and we often restrict attention to linear predictors. The best linear predictor (BLP) of Y given X is the coefficient vector β that solves

$$\beta = \arg\min_{b} \; E[(Y - X'b)^2]. \tag{1.18}$$

Taking the first-order condition,

$$\frac{\partial}{\partial b} E[(Y - X'b)^2] = -2\, E[X(Y - X'b)] = 0, \tag{1.19}$$

which gives the population normal equation

$$E[XX']\,\beta = E[XY]. \tag{1.20}$$

If $E[XX']$ is nonsingular, the BLP coefficient is

$$\beta = \left(E[XX']\right)^{-1} E[XY]. \tag{1.21}$$

This is the population analog of the sample formula $\hat{\boldsymbol{\beta}} = (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{y}$. The population version replaces sample cross-products with their expectations.

The first-order condition can equivalently be written as the population orthogonality condition:

$$E[X(Y - X'\beta)] = 0. \tag{1.22}$$

This says that the linear prediction error $Y - X'\beta$ is uncorrelated with every component of X. It is the population version of the sample condition $\mathbf{X}'\hat{\mathbf{u}} = \mathbf{0}$ that we derived from the matrix normal equations.

1.4.3 The BLP Approximates the CEF

The BLP and the CEF are connected by a key result: the BLP is the best linear approximation to the CEF. To see this, write

$$\beta = \left(E[XX']\right)^{-1} E[XY] = \left(E[XX']\right)^{-1} E[X\, E[Y \mid X]], \tag{1.23}$$

where the second equality uses the law of iterated expectations. The BLP coefficient depends on Y only through the CEF. OLS is fitting a linear function to the CEF, not directly to Y.

When the CEF is linear in X — that is, $E[Y \mid X] = X'\beta$ for some β — the BLP recovers the CEF exactly. The linear model is correctly specified and OLS estimates the true conditional expectation.

When the CEF is nonlinear, the BLP still provides the minimum mean squared error linear approximation to it. The OLS coefficient retains a well-defined interpretation: it is the slope of the best linear predictor, even if the world is nonlinear. This is why OLS remains useful in settings where the true CEF is unknown and possibly complicated.

1.4.4 From Population to Sample

The population formulas become sample formulas when we replace expectations with sample averages. For a sample of n observations,

$$E[XX'] \approx \frac{1}{n}\sum_{i=1}^{n} X_i X_i' = \frac{1}{n}\mathbf{X}'\mathbf{X} \tag{1.24}$$

$$E[XY] \approx \frac{1}{n}\sum_{i=1}^{n} X_i Y_i = \frac{1}{n}\mathbf{X}'\mathbf{y} \tag{1.25}$$

Substituting into the population BLP formula gives

$$\hat{\boldsymbol{\beta}} = \left(\frac{1}{n}\mathbf{X}'\mathbf{X}\right)^{-1}\frac{1}{n}\mathbf{X}'\mathbf{y} = (\mathbf{X}'\mathbf{X})^{-\mathbf{1}}\mathbf{X}'\mathbf{y}, \tag{1.26}$$

which is the OLS estimator derived from the matrix normal equations. The $1/n$ factors cancel. This shows what OLS does: it uses sample analogs to estimate the population BLP, which is the best linear approximation to the conditional expectation function $E[Y \mid X]$.

1.5 Properties of OLS estimator

1.5.1 Unbiasedness

In a linear model

$$\mathbf{y} = \mathbf{X}\boldsymbol{\beta} + \mathbf{u}, \tag{1.27}$$

the OLS estimator is

$$\hat{\boldsymbol{\beta}} = (\mathbf{X}'\mathbf{X})^{-\mathbf{1}}\mathbf{X}'\mathbf{y} \tag{1.28}$$

Since $\mathbf{y} = \mathbf{X}\boldsymbol{\beta} + \mathbf{u}$,

$$\hat{\boldsymbol{\beta}} = \boldsymbol{\beta} + (\mathbf{X}'\mathbf{X})^{-\mathbf{1}}\mathbf{X}'\mathbf{u}. \tag{1.29}$$

This makes

$$\mathrm{E}(\hat{\boldsymbol{\beta}}|\mathbf{X}) = \boldsymbol{\beta} + (\mathbf{X}'\mathbf{X})^{-\mathbf{1}}\mathbf{X}'\mathbf{E}(\mathbf{u}|\mathbf{X}). \tag{1.30}$$

The standard sufficient condition for unbiasedness is:

$$\mathrm{E}(\mathbf{u}|\mathbf{X}) = \mathbf{0}, \tag{1.31}$$

that is, all explanatory variables which form the columns of $\mathbf{X}$ are exogenous. This condition is weaker than the independence condition that u and X are independent. This says that given $\mathbf{X}$, the expected value of $\mathbf{u}$ is zero; it implies that the model is correctly specified. That is, $\mathbf{y}$ is a linear function of $\mathbf{X}$.

In the context of cross-sectional data, this assumption is plausible. However, when we have time series data, the assumption becomes strong, because it assumes that the entire series of $\mathbf{X}$ has no relationship with the error term. In a time series context, this is hard to satisfy. Strictly, unbiasedness needs only $X'E(u \mid X) = 0$, so a failure of Equation 1.31 does not by itself prove bias, but in time series it typically produces it.

For example, suppose we have a model

$$y_t = \beta_1 + \beta_2 y_{t-1} + u_t, \quad u_t \sim \text{IID}(0, \sigma^2). \tag{1.32}$$

In this simple model, even if we assume that y_{t-1} and u_t are uncorrelated, OLS estimator is still biased. That is because $\mathrm{E}(\mathbf{u}|\mathbf{X}) = \mathbf{0}$ is not satisfied. The problem is not that y_{t-1} contains the older errors $u_{t-1}, u_{t-2}, ...$; those are uncorrelated with u_t. It is that u_t enters y_t, and y_t is the regressor in the next row. Conditioning on the whole regressor matrix X means conditioning on future rows, and those depend on u_t, so $E(u_t \mid X) \neq 0$.

There is a weaker condition – weaker than the strict exogeneity above, and in fact the weakest of the conditions in this chapter:

$$\mathrm{E}(\mathbf{X}'\mathbf{u}) = \mathbf{0}, \tag{1.33}$$

This is a moment condition, and it is what GMM estimators are built on. It is a separate assumption from getting the functional form right: a correctly specified linear equation can still have $E(\mathbf{X}'\mathbf{u}) \neq 0$, which is exactly the endogeneity problem of a later chapter.

1.5.2 Consistency

Consistency does not need strict exogeneity. A much weaker condition suffices:

$$\mathrm{E}(u_t|X_t) = 0, \tag{1.34}$$

This condition is much weaker since it only assumes that the mean of current error term does not depend on the current predictors. Even a model with lagged dependent variable can easily satisfy this condition. This condition is called **contemporaneous exogeneity**. It is the weakest of the three conditional-mean conditions worth keeping apart. It is sufficient for consistency but not necessary: what consistency actually requires is the moment condition at the foot of the list below.

- **Strict exogeneity**, $E[u_t \mid X_1, \dots, X_n] = 0$: the error in every period is mean-independent of the regressors in *every* period. This is what unbiasedness requires, and what the lagged-dependent-variable example above violates.
- **Predeterminedness**, $E[u_t \mid \mathcal{F}_t] = 0$, where $\mathcal{F}_t = \sigma(X_1, \dots, X_t)$ is the regressor history through period t – equivalently, X_t is orthogonal to the current and all *future* errors, while being allowed to correlate with *past* errors. This is the relevant condition in models with a lagged dependent variable.
- **Contemporaneous exogeneity**, $E[u_t \mid X_t] = 0$.

- **Contemporaneous orthogonality**, $E[X_t u_t] = 0$ – mere zero correlation. This is weakest of all, and it is what consistency actually needs: together with a rank condition and a law of large numbers – which for dependent data needs the weak-dependence conditions of the MLE chapter – it gives $\operatorname{plim} n^{-1}\mathbf{X}'\mathbf{u} = \mathbf{0}$ and hence $\operatorname{plim} \hat{\beta} = \beta$. Contemporaneous *exogeneity* is a strictly stronger, sufficient condition – it constrains the whole conditional mean rather than a single covariance – so the $E[\mathbf{X}'\mathbf{u}] = 0$ introduced under unbiasedness above sits at the *bottom* of this ladder, not between the others.

Each condition implies the next, because conditioning on a larger information set is a more restrictive requirement, not a less restrictive one. The ladder needs the information sets to be nested, which is why $\mathcal{F}_t$ is defined above as regressor history: $\sigma(X_t) \subseteq \sigma(X_1, \dots, X_t) \subseteq \sigma(X_1, \dots, X_n)$. Widen $\mathcal{F}_t$ to include past errors or other series and strict exogeneity no longer implies predeterminedness, because then neither information set contains the other. By the law of iterated expectations,

$$E[u_t \mid X_t] = E[E[u_t \mid \mathcal{F}_t] \mid X_t] = 0, \tag{1.35}$$

so predeterminedness delivers contemporaneous exogeneity but not conversely. In the time series example, then, the OLS estimator is biased, but can still be consistent, if we are willing to assume no contemporaneous correlation.

Note: general linear hypothesis testing, the Chow test for structural change, and heteroskedasticity- and autocorrelation-consistent (HAC) standard errors (White, Newey-West) are all standard OLS extensions, but are covered in the Maximum Likelihood chapter alongside this guide's other estimation-inference material rather than repeated here.

1.6 Summary

OLS chooses the coefficients that minimize the sum of squared residuals, which gives the normal equations and the estimator $(X'X)^{-1}X'y$. Geometrically it projects y onto the column space of X. The fitted values are the projection and the residuals are orthogonal to it. In the population, OLS estimates the best linear predictor, the best linear approximation to the conditional expectation function. So its coefficients keep a clear meaning even when the true conditional mean is nonlinear.

The properties depend on what we assume about the error. Strict exogeneity, a zero mean of the error given every regressor in every period, makes OLS unbiased. A lagged dependent variable breaks it. Consistency needs much less, only zero correlation between the regressors and the current error, plus a rank condition and a law of large numbers. So OLS can be biased in a dynamic model and still consistent.

2 The Method of Maximum Likelihood

If we make the assumption that the error terms are normally distributed, the maximum likelihood estimators (MLE) coincide with the least square estimators. But MLE can also be applied to a wide variety of models other than regression models, and it generally generates estimators with excellent asymptotic properties. The major disadvantage of MLE is that it requires stronger distributional assumptions than does the method of moments.

2.1 Basic Concepts

Maximum likelihood estimation is to find the set of parameters which makes the current sample most likely given the statistical model.

Suppose we have a sample of n independent and identically distributed (IID) observations. Each of them comes from some distribution function $f(\cdot, \boldsymbol{\theta})$, where parameters $\boldsymbol{\theta}$ is fixed. The joint distribution of all the data points would be

$$f(x_1, x_2, \dots, x_n|\theta) = f(x_1|\theta) \times f(x_2|\theta) \times \cdots \times f(x_n|\theta). \tag{2.1}$$

It is referred to as the likelihood function of the model for the given data set.

Then it's natural to estimate θ by picking the θ that maximize this joint distribution function. A parameter vector $\hat{\boldsymbol{\theta}}$ at which the likelihood takes on its maximum value is called a maximum likelihood estimate, or MLE, of the parameters.

Usually the log form of the likelihood function is preferred for computational purposes.

2.2 MLE for a linear model

Consider the classical normal linear model:

$$\mathbf{y} = \mathbf{X}\boldsymbol{\beta} + \mathbf{u}, \quad \mathbf{u} \sim \mathbf{N}(\mathbf{0}, \boldsymbol{\sigma}^{\mathbf{2}}\mathbf{I}) \tag{2.2}$$

In this case, we have two parameters to estimate, σ and $\boldsymbol{\beta}$. The distribution of $\mathbf{y}|\mathbf{X}$ is

$$f(\mathbf{y} \mid \mathbf{X}, \beta, \sigma) = (2\pi\sigma^2)^{-n/2} \exp\left(-\frac{(\mathbf{y} - \mathbf{X}\boldsymbol{\beta})'(\mathbf{y} - \mathbf{X}\boldsymbol{\beta})}{2\sigma^2}\right) \tag{2.3}$$

The log-likelihood function is

$$l(\mathbf{y}, \boldsymbol{\beta}, \boldsymbol{\sigma}) = -\frac{n}{2}\log(2\pi) - \frac{n}{2}\log(\sigma^2) - \frac{1}{2\sigma^2}(\mathbf{y} - \mathbf{X}\boldsymbol{\beta})'(\mathbf{y} - \mathbf{X}\boldsymbol{\beta}) \tag{2.4}$$

Maximizing this function gives exactly the OLS estimator for $\boldsymbol{\beta}$. That is algebra, and it holds whatever the true error distribution happens to be: the Gaussian log-likelihood is a decreasing function of the sum of squared residuals, so whatever minimizes one maximizes the other. For σ^2 the MLE divides by n rather than $n - k$, so it differs in finite samples and agrees asymptotically. What normality adds is not the formula but its standing. Under normality the OLS estimator *is* the maximum likelihood estimator, and the usual t and F statistics have exact finite-sample distributions. Without normality the same formula is a quasi-MLE: still consistent for $\boldsymbol{\beta}$ under the usual conditions. Its model-based variance $\sigma^2(\mathbf{X}'\mathbf{X})^{-1}$ remains asymptotically valid as long as the errors are homoskedastic and serially uncorrelated. Heteroskedasticity or dependence, not nonnormality, is what calls for the sandwich correction discussed below.

2.3 Asymptotic Properties of MLE

1. Consistency

$$\text{plim}(\hat{\boldsymbol{\theta}}) = \boldsymbol{\theta}. \tag{2.5}$$

This is to say that MLE estimator can get arbitrarily close to the true parameter if the sample size gets to infinity.

2. Asymptotic Normality

$$\sqrt{\mathbf{n}}(\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}) \xrightarrow{d} N(\mathbf{0}, \mathbf{I^{-1}}(\boldsymbol{\theta})), \tag{2.6}$$

where $\mathbf{I}(\boldsymbol{\theta})$ below is the *per-observation* information. Equivalently, in finite-sample terms,

$$\hat{\boldsymbol{\theta}} \approx N(\boldsymbol{\theta}, \mathbf{I_n^{-1}}(\boldsymbol{\theta})), \qquad \mathbf{I_n}(\boldsymbol{\theta}) = \mathbf{n}\,\mathbf{I}(\boldsymbol{\theta}), \tag{2.7}$$

with $\mathbf{I_n}(\boldsymbol{\theta})$ the *full-sample* information. The $\sqrt{n}$ scaling (equivalently, dividing the full-sample information by n) is what makes the statement well-defined; writing $\hat{\theta} \sim N(\theta, I^{-1}(\theta))$ without specifying which information is meant is ambiguous.

This states that the asymptotic distribution of $\hat{\boldsymbol{\theta}}$ is normal with mean $\boldsymbol{\theta}$ and variance given by the inverse of the information, where the per-observation information matrix $\mathbf{I}(\boldsymbol{\theta})$ is defined using the single-observation log-likelihood contribution l_t (so that the full-sample log-likelihood is $l = \sum_{t=1}^{n} l_t$):

$$\mathbf{I}(\boldsymbol{\theta}) = E[(\frac{\partial l_t}{\partial \theta})(\frac{\partial l_t}{\partial \theta})'] = -E[(\frac{\partial^2 l_t}{\partial \theta \partial \theta'})] \tag{2.8}$$

The information matrix is the expected value of the outer product of the gradient of a single observation's log-likelihood contribution, or the negative value of the expected value of its Hessian (second derivative). Using the *full-sample* l here instead of l_t would give the full-sample information $I_n(\theta) = n\,I(\theta)$, not $I(\theta)$ itself.

3. Asymptotic efficiency.

If $\hat{\boldsymbol{\theta}}$ is the MLE estimator of θ and $\mathbf{V}$ denotes its *asymptotic* variance matrix – the variance of the limiting distribution of $\sqrt{n}(\hat{\theta} - \theta)$, not of $\hat{\boldsymbol{\theta}}$ itself, the two differing by a factor of n exactly as with the information matrix above – then

$$\sqrt{n}(\hat{\theta} - \theta) \to^{d} N(\mathbf{0}, \mathbf{V}), \qquad \mathbf{V} = \mathbf{I^{-1}}(\boldsymbol{\theta}). \tag{2.9}$$

If $\tilde{\mathbf{V}}$ denotes the asymptotic variance matrix of any other *regular* consistent, asymptotically normal estimator, then $\tilde{\mathbf{V}} - \mathbf{V}$ is a positive semidefinite matrix. That is, the MLE is the best estimator, in terms of asymptotic variance, in that class. The regularity qualifier is not pedantry: without it the statement is false, since superefficient estimators such as Hodges' can beat the bound at a single point of the parameter space, at the cost of much worse behaviour in a neighbourhood of it.

4. Invariance.

If $\hat{\theta}$ is the MLE of θ and $g(\theta)$ is a continuous function of θ, then $g(\hat{\theta})$ is the MLE of $g(\theta)$.

2.4 When the Model Is Wrong: Quasi-MLE and M-Estimation

2.4.1 What MLE Actually Minimizes

The asymptotic properties above assume that the true density belongs to the parametric family $f(y \mid \theta)$. In practice this is rarely literally true. What does MLE estimate when the model is wrong?

Write $g(y)$ for the true density and $f(y \mid \theta)$ for the parametric model. Maximizing the log-likelihood is equivalent to minimizing the Kullback-Leibler divergence from the true density to the model:

$$\mathrm{KL}(g \,\|\, f(\cdot \mid \theta)) = \int g(y) \log \frac{g(y)}{f(y \mid \theta)}\, dy = E_g[\log g(Y)] - E_g[\log f(Y \mid \theta)]\,. \tag{2.10}$$

The first term does not depend on θ, so minimizing KL over θ is the same as maximizing $E_g[\log f(Y \mid \theta)]$, which is the population version of the sample log-likelihood. The MLE converges to the value θ^* that makes $f(\cdot \mid \theta)$ closest to g in the KL sense, whether or not g belongs to the family. This θ^* is called the pseudo-true value.

2.4.2 The Information Matrix Inequality

Under correct specification, the two forms of the information matrix are equal:

$$J(\theta) = E\left[\frac{\partial l_t}{\partial \theta}\frac{\partial l_t}{\partial \theta}'\right] = -E\left[\frac{\partial^2 l_t}{\partial \theta\, \partial \theta'}\right] = H(\theta). \tag{2.11}$$

This is the information matrix equality. Under misspecification it fails: $J(\theta^*) \neq H(\theta^*)$ in general. The asymptotic variance of the MLE is no longer $I^{-1}(\theta)$ but the sandwich

$$V = H(\theta^*)^{-1}\, J(\theta^*)\, H(\theta^*)^{-1}. \tag{2.12}$$

When the model is correctly specified, $J = H$ and the sandwich collapses to $H^{-1} = I^{-1}$, recovering the standard result. When it is not, H^{-1} is simply the wrong variance – inconsistent, rather than merely too small. There is no general ordering between H^{-1} and the sandwich: it can be too small in some parameter directions and too large in others.

2.4.3 Quasi-Maximum Likelihood Estimation

The sandwich variance makes MLE usable even under misspecification, provided we are clear about what parameter is being estimated. A quasi-maximum likelihood estimator (QMLE) is an MLE computed from a likelihood that may be wrong, together with a sandwich variance that accounts for the misspecification.

The leading case in econometrics is Poisson QMLE for count or nonnegative data. The Poisson log-likelihood for observation t is

$$l_t = y_t \log \mu_t - \mu_t - \log(y_t!), \quad \mu_t = \exp(\mathbf{x}_t'\beta). \tag{2.13}$$

The score equation $\sum_t (y_t - \mu_t)\,\mathbf{x}_t = 0$ depends only on the conditional mean $E[Y_t \mid \mathbf{x}_t] = \mu_t$, not on whether Y_t is actually Poisson. If the conditional mean is correctly specified, the QMLE $\hat{\beta}$ is consistent for β regardless of the true distribution of Y_t — the data need not even be counts. The variance, however, must be estimated with the sandwich, since the Poisson variance assumption $\mathrm{Var}(Y_t \mid \mathbf{x}_t) = \mu_t$ will generally be wrong. A fuller treatment is in the Count Data chapter.

Binary response is a different case, and it is worth being exact about why. If $P(Y = 1 \mid X) = F(X'\beta)$ is correct, then the whole conditional distribution is correct: a Bernoulli law has nothing left to specify once its mean is fixed. Logit and probit are therefore correctly specified maximum likelihood here, not quasi-MLE. The latent-variable story is a representation rather than an extra assumption, and a different latent error distribution generating the same $F(X'\beta)$ is observationally equivalent. Keep the quasi-MLE label for the Poisson case above, where the criterion is used without the full conditional distribution being correct.

2.4.4 M-Estimators

The QMLE perspective generalizes further. An M-estimator is any estimator defined as the solution to

$$\sum_{t=1}^{n} \psi(y_t, \mathbf{x}_t, \theta) = 0, \tag{2.14}$$

where ψ is a known vector-valued function. MLE is the special case where $\psi = \partial l_t / \partial \theta$, the score. OLS is the case $\psi(y_t, \mathbf{x}_t, \beta) = \mathbf{x}_t(y_t - \mathbf{x}_t'\beta)$, the population orthogonality condition from the OLS chapter. GMM generalizes further by allowing more equations than parameters; the GMM chapter develops this.

All M-estimators share the same asymptotic structure under regularity conditions:

$$\sqrt{n}\,(\hat{\theta} - \theta^*) \xrightarrow{d} N(0,\ A^{-1}\,B\,A^{-1\prime})\,, \tag{2.15}$$

where $A = E[\partial\psi/\partial\theta']$ and $B = E[\psi\psi']$. This is the sandwich again. When ψ is a score from a correctly specified likelihood, $A = -H$, $B = J = H$, and the sandwich reduces to H^{-1}.

The M-estimator framework unifies OLS, MLE, QMLE, and GMM as special cases of the same asymptotic theory. The sandwich variance is the general formula; the simpler I^{-1} or $\sigma^2(X'X)^{-1}$ are special cases that hold under correct specification.

2.5 Example

Let's look at the example of regression of car's mpg on disp, hp and wt. This time we use MLE. (R has a mle() function which I cannot make it work on this example. So I am using optim().)

```
ols.lf <- function(theta, y, X) {
  beta <- theta[-1]
  sigma2 <- theta[1]
  # Return a large penalty rather than NA: L-BFGS-B's line search can
  # transiently evaluate points that violate the box constraint, and
  # `fn` returning NA there causes a fatal "L-BFGS-B needs finite values
  # of 'fn'" error. A finite penalty keeps the optimizer on track instead.
  if (sigma2 <= 0) return(1e10)
  n <- nrow(X)
  e <- y - X%*%beta
  logl <- ((-n/2)*log(2*pi)) - ((n/2)*log(sigma2)) - ((t(e)%*%e)/(2*sigma2))
  return(-logl)
}
X <- cbind(1,mtcars[,c('disp','hp','wt')])
y <- mtcars$mpg

# Deliberately naive starting values (sigma2 = 1, slopes at +/-1): the point
# of the exercise is that the optimizer finds the OLS solution without being
# handed it. For harder likelihoods than this one, good starting values matter
# a great deal -- they reduce the risk of converging to a spurious local
# maximum, or of failing to converge at all.
optim(c(1,1,-1,-1,-1), method="L-BFGS-B", fn=ols.lf,
 ↪  lower=c(1e-6,-Inf,-Inf,-Inf,-Inf), upper=rep(Inf,5), y=y, X=as.matrix(X))
```

```
$par
[1]  6.0935580702 37.1058998652 -0.0009352804 -0.0311594264 -3.8010038304

$value
[1] 74.32149

$counts
function gradient
     126      126
```

```
$convergence
[1] 0

$message
[1] "CONVERGENCE: REL_REDUCTION_OF_F <= FACTR*EPSMCH"
```

```r
# it should match the OLS estimator

lm1 <- lm(mpg~disp+hp+wt, data=mtcars)
summary(lm1)
```

```
Call:
lm(formula = mpg ~ disp + hp + wt, data = mtcars)

Residuals:
   Min     1Q Median     3Q    Max
-3.891 -1.640 -0.172  1.061  5.861

Coefficients:
             Estimate Std. Error t value Pr(>|t|)
(Intercept) 37.105505   2.110815  17.579  < 2e-16 ***
disp        -0.000937   0.010350  -0.091  0.92851
hp          -0.031157   0.011436  -2.724  0.01097 *
wt          -3.800891   1.066191  -3.565  0.00133 **
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 2.639 on 28 degrees of freedom
Multiple R-squared:  0.8268,    Adjusted R-squared:  0.8083
F-statistic: 44.57 on 3 and 28 DF,  p-value: 8.65e-11
```

2.6 Hypotheses Testing

2.6.1 Linear Hypotheses Testing

Linear hypothesis can be framed as

$$\mathbf{H_0}: \quad \mathbf{R}\boldsymbol{\beta} - \mathbf{r} = \mathbf{0}, \tag{2.16}$$

where $\mathbf{R}$ is a $q \times k$ matrix of known constants with $q \leq k$ and $\text{rank}(\mathbf{R}) = q$, and $\mathbf{r}$ is a q-vector of known constants. The restrictions may number as many as the parameters – testing that every coefficient is zero is the case $q = k$ – but they must be linearly independent.

We assume that u_i are normally distributed.

$$\mathbf{u} \sim \mathbf{N}(\mathbf{0}, \boldsymbol{\sigma}^{\mathbf{2}}\mathbf{I}) \tag{2.17}$$

Even if this assumption does not hold, we still have asymptotic normality for $\hat{\boldsymbol{\beta}}$, so the tests below remain valid asymptotically – provided the covariance estimator is valid for whatever heteroskedasticity or dependence is actually present. The model-based $\sigma^2(\mathbf{X}'\mathbf{X})^{-1}$ is not: it encodes the homoskedastic, serially uncorrelated structure of Equation 2.17. Dropping normality alone does not invalidate it: under homoskedastic, serially uncorrelated errors the tests remain asymptotically valid with that formula. Heteroskedasticity or serial correlation does invalidate it. Use a sandwich estimator under heteroskedasticity and a HAC estimator under serial correlation.

Since linear combination of normal variables are also normally distributed,

$$\begin{aligned} \hat{\boldsymbol{\beta}} &\sim \mathbf{N}(\boldsymbol{\beta}, \boldsymbol{\sigma^2}(\mathbf{X}'\mathbf{X})^{-1}) \\ \mathbf{R}\hat{\boldsymbol{\beta}} &\sim \mathbf{N}(\mathbf{R}\boldsymbol{\beta}, \boldsymbol{\sigma^2}\mathbf{R}(\mathbf{X}'\mathbf{X})^{-1}\mathbf{R}') \\ \mathbf{R}(\hat{\boldsymbol{\beta}} - \boldsymbol{\beta}) &\sim \mathbf{N}(\mathbf{0}, \boldsymbol{\sigma^2}\mathbf{R}(\mathbf{X}'\mathbf{X})^{-1}\mathbf{R}') \end{aligned} \tag{2.18}$$

Under null hypothesis,

$$(\mathbf{R}\hat{\boldsymbol{\beta}} - \mathbf{r}) \sim \mathbf{N}(\mathbf{0}, \boldsymbol{\sigma^2}\mathbf{R}(\mathbf{X}'\mathbf{X})^{-1}\mathbf{R}') \tag{2.19}$$

Therefore,

$$(\mathbf{R}\hat{\boldsymbol{\beta}} - \mathbf{r})'[\boldsymbol{\sigma^2}\mathbf{R}(\mathbf{X}'\mathbf{X})^{-1}\mathbf{R}']^{-1}(\mathbf{R}\hat{\boldsymbol{\beta}} - \mathbf{r}) \sim \chi^2(q) \tag{2.20}$$

Here we don't know σ^2. However, it can be shown that

$$\frac{\hat{\mathbf{u}}'\hat{\mathbf{u}}}{\boldsymbol{\sigma^2}} \sim \chi^2(n-k) \tag{2.21}$$

Then we have

$$\frac{(\mathbf{R}\hat{\boldsymbol{\beta}} - \mathbf{r})'[\mathbf{R}(\mathbf{X}'\mathbf{X})^{-1}\mathbf{R}']^{-1}(\mathbf{R}\hat{\boldsymbol{\beta}} - \mathbf{r})/\mathbf{q}}{\hat{\mathbf{u}}'\hat{\mathbf{u}}/(\mathbf{n}-\mathbf{k})} \sim F(q, n-k) \tag{2.22}$$

This is an example of the F test for any linear hypotheses testing. t test will be a special case of this.

2.6.2 The three classical tests: LR, Wald and LM

The F test above leans on normal errors. The general machinery for testing $\mathbf{H_0} : \mathbf{R}\boldsymbol{\theta} - \mathbf{r} = \mathbf{0}$ after maximum likelihood needs only the asymptotic results already established, and it comes in three forms. Write $\hat{\theta}_u$ for the unrestricted MLE, $\hat{\theta}_r$ for the MLE subject to the restriction, $s(\theta) = \partial\ell/\partial\theta$ for the score, and $\mathbf{I_n}(\boldsymbol{\theta}) = \mathbf{n}\,\mathbf{I}(\boldsymbol{\theta})$ for the full-sample information.

Likelihood ratio. Compare the two maximized log-likelihoods:

$$LR = 2\big[\ell(\hat{\theta}_u) - \ell(\hat{\theta}_r)\big] \xrightarrow{d} \chi^2(q). \tag{2.23}$$

Wald. Ask whether the *unrestricted* estimate is far from satisfying the restriction, measuring distance in units of its own estimated variance:

$$W = (\mathbf{R}\hat{\theta}_u - \mathbf{r})'\big[\mathbf{R}\,\mathbf{I}_n^{-1}(\hat{\theta}_u)\,\mathbf{R}'\big]^{-1}(\mathbf{R}\hat{\theta}_u - \mathbf{r}) \xrightarrow{d} \chi^2(q). \tag{2.24}$$

Lagrange multiplier (score). Ask whether the score, evaluated at the *restricted* estimate, is far from zero – if the restriction is harmless, the restricted point should already be nearly a maximum:

$$LM = s(\hat{\theta}_r)' \, \mathbf{I}_n^{-1}(\hat{\theta}_r) \, s(\hat{\theta}_r) \xrightarrow{d} \chi^2(q). \tag{2.25}$$

The three are asymptotically equivalent under the null and have the same local power, so the choice among them is one of convenience: **Wald needs only the unrestricted fit, LM needs only the restricted fit, and LR needs both.** That is why LM is the natural choice for specification testing, where the restricted model is the easy one to estimate (White's heteroskedasticity test and the Breusch-Godfrey serial-correlation test are both LM tests run as auxiliary regressions), while Wald is what software reports by default, since the unrestricted fit is what was just computed.

In the classical normal linear model the three are monotone transformations of the same F statistic above, so nothing is lost by using F there. Two cautions otherwise. In finite samples the three can disagree, sometimes enough to straddle a conventional critical value, and there is a known ordering $W \geq LR \geq LM$ in the linear model under normality. And the Wald statistic is **not invariant to nonlinear reparameterization**: testing $\gamma = 1$ and testing $\log \gamma = 0$ are the same hypothesis but give different Wald statistics, whereas LR is invariant because it compares likelihood *values*, which reparameterization does not change. When a hypothesis can be written several algebraically equivalent ways, prefer LR.

2.7 Test of Structural Change (Chow test)

Chow test is a test between two groups of observations. For example, we have data on consumption function of prewar period and postwar period. It is natural to think of a test on whether consumption function parameters differ between prewar and postwar periods. The null hypothesis is that there is no difference.

Let $\mathbf{y_i}$, $\mathbf{X_i}$ $(i = 1, 2)$ indicate the appropriate partitioning of the data. The unrestricted model may be written

$$\begin{bmatrix} \mathbf{y_1} \\ \mathbf{y_2} \end{bmatrix} = \begin{bmatrix} \mathbf{X_1} & \mathbf{0} \\ \mathbf{0} & \mathbf{X_2} \end{bmatrix} \begin{bmatrix} \boldsymbol{\beta_1} \\ \boldsymbol{\beta_2} \end{bmatrix} + u \quad u \sim N(0, \sigma^2 I) \tag{2.26}$$

The null hypothesis of no structural break is

$$\mathbf{H_0}: \quad \boldsymbol{\beta_1} = \boldsymbol{\beta_2}. \tag{2.27}$$

Running OLS on two equations separately, we have

$$\begin{bmatrix} \hat{\boldsymbol{\beta}}_1 \\ \hat{\boldsymbol{\beta}}_2 \end{bmatrix} = \begin{bmatrix} \mathbf{X_1'X_1} & \mathbf{0} \\ \mathbf{0} & \mathbf{X_2'X_2} \end{bmatrix}^{-1} \begin{bmatrix} \mathbf{X_1'y_1} \\ \mathbf{X_2'y_2} \end{bmatrix} = \begin{bmatrix} (\mathbf{X_1'X_1})^{-1}\mathbf{X_1'y_1} \\ (\mathbf{X_2'X_2})^{-1}\mathbf{X_2'y_2} \end{bmatrix} \tag{2.28}$$

Restricted model under null hypothesis

$$\begin{bmatrix} \mathbf{y_1} \\ \mathbf{y_2} \end{bmatrix} = \begin{bmatrix} \mathbf{X_1} \\ \mathbf{X_2} \end{bmatrix} \boldsymbol{\beta} + \mathbf{u} \tag{2.29}$$

The test of the null hypothesis is given by

$$\mathbf{F} = \frac{(\hat{\mathbf{u}}_*'\hat{\mathbf{u}}_* - \hat{\mathbf{u}}'\hat{\mathbf{u}})/\mathbf{k}}{\hat{\mathbf{u}}'\hat{\mathbf{u}}/(\mathbf{n}-\mathbf{2k})} \sim F(k, n-2k) \tag{2.30}$$

where $\hat{u}_*$ is the residual of the restricted model, $\hat{u}$ is the stacked residual of two unrestricted models.

For example, for prewar period (suppose there is n_1 observations):

$$y_1 = \beta_0 + \beta_1 x_1 + \beta_2 x_2 \tag{2.31}$$

For postwar period (suppose there is n_2 observations):

$$y_2 = \gamma_0 + \gamma_1 x_1 + \gamma_2 x_2 \tag{2.32}$$

The null hypothesis of no structural break is

$$\mathbf{H_0}: \quad \boldsymbol{\beta_0} = \boldsymbol{\gamma_0}, \boldsymbol{\beta_1} = \boldsymbol{\gamma_1}, \boldsymbol{\beta_2} = \boldsymbol{\gamma_2}. \tag{2.33}$$

Running OLS on two equations separately, we have two sums of squares of error (SSR_1 and SSR_2). The unrestricted error sum of squares is

$$SSR_U = SSR_1 + SSR_2 \tag{2.34}$$

Then run the regression on the stacked sample of $n_1 + n_2$ observations and get SSR_R.

Then

$$\frac{(SSR_R - SSR_U)/k}{SSR_U/(n_1 + n_2 - 2k)} \sim F_{k, n_1+n_2-2k}. \tag{2.35}$$

k is number of restrictions; in this example, it is 3.

The other way to do the same test is easier: simply include interaction terms between group dummy and explanatory variables in the regression.

For our earlier example, we would estimate the model:

$$y = \alpha_0 + \alpha_1 x_1 + \alpha_2 x_2 + \eta_0 d + \eta_1 x_1 d + \eta_2 x_2 d \tag{2.36}$$

Here d is a dummy variable for postwar period. The Chow test is simply a test on the hypothesis that all the coefficients involving d are zero. A regression of the above model with $n_1 + n_2$ observations and a F test are easy to do.

$$\mathbf{H_0}: \quad \boldsymbol{\eta_0} = \boldsymbol{\eta_1} = \boldsymbol{\eta_2} = \mathbf{0}. \tag{2.37}$$

2.7.1 An example of Chow test in R

Here we use an example from the "strucchange" library.

```
## Example 7.4 from Greene (1993), "Econometric Analysis"
## Chow test on Longley data
data("longley")
library(strucchange)
## use structural change test from the library.
sctest(Employed ~ Year + GNP.deflator + GNP + Armed.Forces, data = longley,
  type = "Chow", point = 7)
```

```
    Chow test

data:  Employed ~ Year + GNP.deflator + GNP + Armed.Forces
F = 3.9268, p-value = 0.06307
```

```
## which is equivalent to segmenting the regression via
fac <- factor(c(rep(1, 7), rep(2, 9)))
## here fac is the factor for segmenting it at point 7.
fm0 <- lm(Employed ~ Year + GNP.deflator + GNP + Armed.Forces, data = longley)
fm1 <- lm(Employed ~ fac/(Year + GNP.deflator + GNP + Armed.Forces), data =
 ↪  longley)
## Here anova is returning the F test (Chow test). Equivalent to sctest results.
anova(fm0, fm1)
```

```
Analysis of Variance Table

Model 1: Employed ~ Year + GNP.deflator + GNP + Armed.Forces
Model 2: Employed ~ fac/(Year + GNP.deflator + GNP + Armed.Forces)
  Res.Df    RSS Df Sum of Sq      F  Pr(>F)
1     11 4.8987
2      6 1.1466  5    3.7521 3.9268 0.06307 .
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
## estimates from Table 7.5 in Greene (1993)
summary(fm0)
```

```
Call:
lm(formula = Employed ~ Year + GNP.deflator + GNP + Armed.Forces,
    data = longley)

Residuals:
    Min      1Q  Median      3Q     Max
```

```
-0.9058 -0.3427 -0.1076  0.2168  1.4377

Coefficients:
               Estimate Std. Error t value Pr(>|t|)
(Intercept)   1.169e+03  8.359e+02   1.399  0.18949
Year         -5.765e-01  4.335e-01  -1.330  0.21049
GNP.deflator -1.977e-02  1.389e-01  -0.142  0.88940
GNP           6.439e-02  1.995e-02   3.227  0.00805 **
Armed.Forces -1.015e-04  3.086e-03  -0.033  0.97436
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 0.6673 on 11 degrees of freedom
Multiple R-squared:  0.9735,    Adjusted R-squared:  0.9639
F-statistic: 101.1 on 4 and 11 DF,  p-value: 1.346e-08
```

```
summary(fm1)
```

```
Call:
lm(formula = Employed ~ fac/(Year + GNP.deflator + GNP + Armed.Forces),
    data = longley)

Residuals:
     Min       1Q   Median       3Q      Max
-0.47717 -0.18950  0.02089  0.14836  0.56493

Coefficients:
                    Estimate Std. Error t value Pr(>|t|)
(Intercept)        1.678e+03  9.390e+02   1.787  0.12413
fac2               2.098e+03  1.786e+03   1.174  0.28473
fac1:Year         -8.352e-01  4.847e-01  -1.723  0.13563
fac2:Year         -1.914e+00  7.913e-01  -2.419  0.05194 .
fac1:GNP.deflator -1.633e-01  1.762e-01  -0.927  0.38974
fac2:GNP.deflator -4.247e-02  2.238e-01  -0.190  0.85576
fac1:GNP           9.481e-02  3.815e-02   2.485  0.04747 *
fac2:GNP           1.123e-01  2.269e-02   4.951  0.00258 **
fac1:Armed.Forces -2.467e-03  6.965e-03  -0.354  0.73532
fac2:Armed.Forces -2.579e-02  1.259e-02  -2.049  0.08635 .
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 0.4372 on 6 degrees of freedom
Multiple R-squared:  0.9938,    Adjusted R-squared:  0.9845
F-statistic: 106.9 on 9 and 6 DF,  p-value: 6.28e-06
```

2.8 Heteroskedasticity and Autocorrelation Consistent Standard Errors

2.8.1 Variance estimator under homoscedasticity

In OLS regression, the variance-covariance matrix of $\hat{\beta}$ is

$$\text{var}(\hat{\boldsymbol{\beta}}) = \mathbf{E}[(\hat{\boldsymbol{\beta}} - \boldsymbol{\beta})(\hat{\boldsymbol{\beta}} - \boldsymbol{\beta})'] = \mathbf{E}[(\mathbf{X'X})^{-1}\mathbf{X'uu'X}(\mathbf{X'X})^{-1}] = \boldsymbol{\sigma^2}(\mathbf{X'X})^{-1} \tag{2.38}$$

if

$$\text{var}(\mathbf{u}) = \boldsymbol{\sigma^2}\mathbf{I}. \tag{2.39}$$

That is, the error term has mean zero and constant variance (homoscedasticity). The pairwise correlation between error terms is always zero (no serial correlation).

The estimation of σ^2:

$$s^2 = \frac{\hat{\mathbf{u}}'\hat{\mathbf{u}}}{n-k} \tag{2.40}$$

is an unbiased estimator of σ^2 (proof omitted).

The standard estimate of the variance-covariance matrix of the OLS parameter estimates under the assumption of IID errors is

$$\hat{\text{Var}}(\hat{\beta}) = s^2(\mathbf{X'X})^{-1} \tag{2.41}$$

2.8.2 White's estimator

We made strong assumption that the error terms of the regression model are IID when we estimate the variance-covariance matrix of OLS estimator. Under this assumption, the usual estimator of variance-covariance matrix of $\hat{\boldsymbol{\beta}}$ is consistent. Now let's relax this assumption to only independent but not identically distributed. Again, the linear regression models is

$$\mathbf{y} = \mathbf{X}\boldsymbol{\beta} + \mathbf{u}, \quad \mathbf{E}(\mathbf{u}) = \mathbf{0}, \quad \mathbf{E}(\mathbf{uu}') = \boldsymbol{\Omega}, \tag{2.42}$$

where $\boldsymbol{\Omega}$ is the error variance-covariance matrix with diagonal elements being σ_t^2 for t^{th} element, off-diagonal elements being zero. In other words, the error terms are heteroscedastic.

The variance-covariance matrix of the OLS estimator $\hat{\boldsymbol{\beta}}$ is equal to

$$\begin{aligned} \text{Var}(\hat{\boldsymbol{\beta}}) &= \mathbf{E}[(\hat{\boldsymbol{\beta}} - \boldsymbol{\beta})(\hat{\boldsymbol{\beta}} - \boldsymbol{\beta})'] \\ &= [(\mathbf{X'X})^{-1}\mathbf{X}'(\mathbf{E}(\mathbf{uu}'))\mathbf{X}(\mathbf{X'X})^{-1}] \\ &= (\mathbf{X'X})^{-1}\mathbf{X'\Omega X}(\mathbf{X'X})^{-1} \end{aligned} \tag{2.43}$$

If we know σ_t^2, then we would be able to estimate this "sandwich covariance matrix". But we don't.

$$\text{Var}(\hat{\boldsymbol{\beta}}) = \frac{\mathbf{1}}{\mathbf{n}}[\frac{\mathbf{1}}{\mathbf{n}}(\mathbf{X'X})]^{-1}[\frac{\mathbf{1}}{\mathbf{n}}\mathbf{X'\Omega X}][\frac{\mathbf{1}}{\mathbf{n}}(\mathbf{X'X})]^{-1} \tag{2.44}$$

Let y_t denote the tth observation on the dependent variable, and $x_t' = [1 x_{2t} \cdots x_{kt}]$ denote the tth row of the $\mathbf{X}$ matrix. Then

$$\mathbf{X'\Omega X} = \sum_{\mathbf{t=1}}^{\mathbf{n}} \boldsymbol{\sigma}_\mathbf{t}^\mathbf{2} \mathbf{x_t x_t'} \tag{2.45}$$

The White estimator replaces the unknown σ_t^2 by $\hat{u}_t^2$, the estimated OLS residuals. This provides a consistent estimator of the variance matrix for the OLS coefficient vector and is particularly useful since it does not require any specific assumptions about the form of the heteroscedasticity.

Therefore,

$$\begin{aligned} \widehat{\mathrm{Var}}(\hat{\boldsymbol{\beta}}) &= \frac{\mathbf{1}}{\mathbf{n}}[\frac{\mathbf{1}}{\mathbf{n}}(\mathbf{X'X})]^{-\mathbf{1}}[\frac{\mathbf{1}}{\mathbf{n}}\mathbf{X'}\hat{\mathbf{\Omega}}\mathbf{X}][\frac{\mathbf{1}}{\mathbf{n}}(\mathbf{X'X})]^{-\mathbf{1}} \\ \hat{\mathbf{\Omega}} &= \mathbf{diag}(\hat{\mathbf{u}}_\mathbf{1}^\mathbf{2}, \hat{\mathbf{u}}_\mathbf{2}^\mathbf{2}, \cdots, \hat{\mathbf{u}}_\mathbf{n}^\mathbf{2}) \end{aligned} \tag{2.46}$$

2.8.3 Newey-West estimator

White's estimator deals with the situation that we have heteroskedasticity (a diagonal Ω) of unknown form. When we have serial correlation of unknown form (a non-diagonal Ω), we can estimate the variance-covariance matrix by a heteroskedasticity and autocorrelation consistent, or HAC, estimator. Newey-West estimator is the most popular HAC estimator.

Given a time series data set, suppose we are interested in estimating the mean vector (suppose we have more than one variable) and its variance. We know that given IID data, we can apply central limit theorem: sample mean is a consistent estimator of the population mean and it's variance can be calculated since asymptotically the sample mean conforms to a normal distribution and the variance can be estimated, relatively easily. However, in the case of time series data, autocorrelation usually exists. We may be concerned the CLT may not work in this case.

Fortunately, autocorrelation does not by itself defeat inference, though the conditions need care. Suppose $\mathbf{y_t}$ is covariance-stationary (the covariance is not a function of time) *and* its autocovariances are absolutely summable, $\sum_{v=-\infty}^{\infty} \|\Gamma_v\| < \infty$. Then, as in Hamilton (1994), the sample mean satisfies:

$$\bar{\mathbf{y}}_\mathbf{T} \to \boldsymbol{\mu}, \tag{2.47}$$

$$\mathbf{S} = \lim_{T\to\infty} \mathbf{T} \cdot \mathbf{E}[(\bar{\mathbf{y}}_\mathbf{T} - \boldsymbol{\mu})(\bar{\mathbf{y}}_\mathbf{T} - \boldsymbol{\mu})'] = \sum_{\mathbf{v}=-\infty}^{\infty} \mathbf{\Gamma_v}. \tag{2.48}$$

where $\mathbf{\Gamma_v}$ is the variance-covariance matrix for $\mathbf{y_t}$ and $\mathbf{y_{t-v}}$.

The first one says the law of large numbers still holds. The second gives the long-run variance used for the standard error, and absolute summability is what makes that sum exist.

Covariance stationarity on its own is not enough for either statement. It can fail even the law of large numbers. Let $y_t = z + \varepsilon_t$, where z is drawn once and then held fixed and ε_t is white noise. Every autocovariance equals $\mathrm{Var}(z)$, so the process is covariance-stationary, the autocovariances are not summable, and $\bar{y}_T$ converges to z rather than to μ. Asymptotic normality asks for more again: a mixing, near-epoch dependence, or martingale-difference condition, together with moment restrictions. Covariance stationarity gives the variance formula, not the central limit theorem.

If the data were generated by a vector MA(q) process, then

$$\mathbf{S} = \sum_{v=-q}^{q} \Gamma_v. \tag{2.49}$$

A natural estimate is

$$\hat{\mathbf{S}} = \hat{\mathbf{\Gamma}}_\mathbf{0} + \sum_{\mathbf{v=1}}^{\mathbf{q}} (\hat{\mathbf{\Gamma}}_\mathbf{v} + \hat{\mathbf{\Gamma}}'_\mathbf{v}), \tag{2.50}$$

where

$$\hat{\mathbf{\Gamma}}_\mathbf{v} = (\mathbf{1}/\mathbf{T}) \sum_{\mathbf{t=v+1}}^{\mathbf{T}} (\mathbf{y_t} - \bar{\mathbf{y}})(\mathbf{y_{t-v}} - \bar{\mathbf{y}})'. \tag{2.51}$$

This gives a consistent estimate of $\mathbf{S}$; however, it sometimes is not positive semidefinite.

Whitney K. Newey and West (1987) suggested putting in a weight:

$$\hat{\mathbf{S}} = \hat{\mathbf{\Gamma}}_\mathbf{0} + \sum_{v=1}^{q} (1 - \frac{v}{q+1})(\hat{\mathbf{\Gamma}}_\mathbf{v} + \hat{\mathbf{\Gamma}}'_\mathbf{v}), \tag{2.52}$$

Here q is a truncation bandwidth, not a true MA order. For an MA(q) process the two coincide, but the Newey-West estimator is used precisely when the dependence has no known finite order: the Bartlett weights $1 - v/(q+1)$ downweight higher lags and keep $\hat{S}$ positive semidefinite, and consistency comes from letting q grow with T, slowly enough that the poorly estimated high-order autocovariances do not dominate.

Consider a linear regression model:

$$y_t = \mathbf{x}'_\mathbf{t}\boldsymbol{\beta} + u_t \tag{2.53}$$

Suppose we have the OLS estimator $\mathbf{b_T}$, then

$$\sqrt{T}(\mathbf{b_T} - \boldsymbol{\beta}) = [(1/T)\sum_{t=1}^{T} \mathbf{x_t}\mathbf{x}'_\mathbf{t}]^{-1}[(1/\sqrt{T})\sum_{t=1}^{T} \mathbf{x_t}\mathbf{u_t}] \tag{2.54}$$

The first term converges in probability to some constant. The second term is $\sqrt{T}$ times the sample mean of the vector $\mathbf{x_t}\mathbf{u_t}$, which is exactly the form to which a central limit theorem applies.

Under general conditions,

$$\sqrt{T}(\mathbf{b_T} - \boldsymbol{\beta}) \rightarrow^L N(0, Q^{-1}SQ^{-1}) \tag{2.55}$$

where $Q = \operatorname{plim}(1/T)\sum_{t=1}^{T} \mathbf{x_t}\mathbf{x}'_\mathbf{t} = \mathrm{E}[\mathbf{x_t}\mathbf{x}'_\mathbf{t}]$, and S can be estimated by

$$\hat{\mathbf{S}}_\mathbf{T} = \hat{\mathbf{\Gamma}}_\mathbf{0T} + \sum_{v=1}^{q} (1 - \frac{v}{q+1})(\hat{\mathbf{\Gamma}}_\mathbf{v,T} + \hat{\mathbf{\Gamma}}'_\mathbf{v,T}), \tag{2.56}$$

where

$$\hat{\Gamma}_{v,T} = (1/T) \sum_{t=v+1}^{T} (x_t \hat{u}_{t,T} \hat{u}_{t-v,T} x'_{t-v}), \tag{2.57}$$

where $\hat{u}_{t,T}$ is the OLS residual for data t in a sample of size T.

Overall, the variance of $\mathbf{b_T}$ is approximated by

$$\hat{\Sigma}_{NW} = [\sum_{t=1}^{T} x_t x_t']^{-1} [\sum_{t=1}^{T} \hat{u}_t^2 x_t x_t' + \sum_{v=1}^{q} (1 - \frac{v}{q+1}) \sum_{t=v+1}^{T} (x_t \hat{u}_{t,T} \hat{u}_{t-v,T} x_{t-v}' + x_{t-v} \hat{u}_{t-v,T} \hat{u}_{t,T} x_t')] [\sum_{t=1}^{T} x_t x_t']^{-1} \tag{2.58}$$

This estimation obviously depends on the selection of q, the lag length beyond which the autocorrelation of $x_t u_t$ and $x_{t-v} u_{t-v}$ is treated as negligible. One common rule of thumb sets $q = 0.75 \cdot T^{\frac{1}{3}}$; it is a convention rather than a theorem, and other growth rates are in use. W. K. Newey and West (1994) has suggested a way to automatically select the bandwidth q. Here we omitted the discussion. Both Stata and R now also implement the W. K. Newey and West (1994) estimator, with no need to specify q.

2.9 Summary

Maximum likelihood picks the parameters that make the observed sample most likely under the assumed model. With normal errors in a linear model it gives the OLS coefficients. Under correct specification the MLE is consistent, asymptotically normal with variance equal to the inverse information, efficient among regular estimators, and invariant to reparameterization. When the model is wrong, the MLE converges to the pseudo-true value closest to the truth in Kullback-Leibler divergence, and the variance becomes a sandwich. Poisson QMLE is the leading case: it is consistent whenever the conditional mean is right. OLS, MLE, QMLE and GMM are all M-estimators with the same sandwich variance.

The rest of the chapter is about testing. The F test handles linear restrictions under normal errors. The likelihood-ratio, Wald and LM tests do the same in general and agree asymptotically. The Chow test checks for a structural break. White and Newey-West standard errors keep inference valid under heteroskedasticity and autocorrelation.

3 GLS

3.1 Introduction

Since heteroscedasticity and serial correlation affect both linear and nonlinear regression models in the same way, there is no harm in limiting our attention to the simpler linear case.

$$\mathbf{y} = \mathbf{X}\boldsymbol{\beta} + \mathbf{u}, \quad \mathbf{E}(\mathbf{u}) = \mathbf{0}, \quad \mathbf{E}(\mathbf{u}\mathbf{u}') = \boldsymbol{\Omega}, \tag{3.1}$$

where $\boldsymbol{\Omega}$, the variance-covariance matrix of the error term, is a positive definite $n \times n$ matrix. If $\boldsymbol{\Omega}$ is equal to $\sigma^2\mathbf{I}$, then it is just the linear regression model without heteroscedasticity and serial correlation. If $\boldsymbol{\Omega}$ is diagonal with nonconstant diagonal elements, then the error terms are still uncorrelated, but there is heteroscedasticity. If $\boldsymbol{\Omega}$ is not diagonal, then u_i and u_j are correlated. In next section, we obtain an efficient estimator for the vector β by transforming the regression so that it satisfies the conditions of Gauss-Markov Theorem. This efficient estimator is called the Generalized Least Squares, or GLS, estimator.

3.2 The GLS Estimator

Since $\boldsymbol{\Omega}$, the variance-covariance matrix of the error term, is a positive definite $n \times n$ matrix, there always exist full-rank $n \times n$ matrices (usually triangular) $\boldsymbol{\Psi}$ such that

$$\boldsymbol{\Omega}^{-1} = \boldsymbol{\Psi}\boldsymbol{\Psi}'. \tag{3.2}$$

Premultiplying (the equation above) by $\boldsymbol{\Psi}'$ gives

$$\boldsymbol{\Psi}'\mathbf{y} = \boldsymbol{\Psi}'\mathbf{X}\boldsymbol{\beta} + \boldsymbol{\Psi}'\mathbf{u}. \tag{3.3}$$

The OLS estimator from regression (the equation above) is

$$\hat{\boldsymbol{\beta}}_{\mathbf{GLS}} = (\mathbf{X}'\boldsymbol{\Psi}\boldsymbol{\Psi}'\mathbf{X})^{-1}\mathbf{X}'\boldsymbol{\Psi}\boldsymbol{\Psi}'\mathbf{y} = (\mathbf{X}'\boldsymbol{\Omega}^{-1}\mathbf{X})^{-1}\mathbf{X}'\boldsymbol{\Omega}^{-1}\mathbf{y} \tag{3.4}$$

The transformed error term has an identity variance-covariance matrix:

$$\mathbf{E}(\boldsymbol{\Psi}'\mathbf{u}\mathbf{u}'\boldsymbol{\Psi}) = \boldsymbol{\Psi}'\mathbf{E}(\mathbf{u}\mathbf{u}')\boldsymbol{\Psi} = \boldsymbol{\Psi}'\boldsymbol{\Omega}\boldsymbol{\Psi} = \mathbf{I} \tag{3.5}$$

Since the transformed model satisfies OLS assumptions, we have

$$\begin{aligned} \mathrm{Var}(\hat{\boldsymbol{\beta}}_{\mathbf{GLS}}) &= (\mathbf{X}'\boldsymbol{\Psi}\boldsymbol{\Psi}'\mathbf{X})^{-1} \\ &= (\mathbf{X}'\boldsymbol{\Omega}^{-1}\mathbf{X})^{-1} \end{aligned} \tag{3.6}$$

3.3 Weighted Least Squares

It is easy to obtain GLS estimates when the error terms are heteroscedastic but uncorrelated, which means $\mathbf{\Omega}$ is diagonal. Let ω_t^2 denote the tth diagonal element of $\mathbf{\Omega}$. Then $\mathbf{\Psi}$ can be chosen as the diagonal matrix with tth diagonal element ω_t^{-1}. For a typical observation, regression can be written as

$$\omega_t^{-1} y_t = \omega_t^{-1} \mathbf{x}_t' \beta + \omega_t^{-1} u_t. \tag{3.7}$$

This is called weighted least squares, or WLS. It is worth keeping two notions of "weight" distinct. The transformed variables y_t and $\mathbf{x_t'}$ are *scaled* by ω_t^{-1}; equivalently, the WLS objective weights each squared residual by the inverse variance ω_t^{-2} (the square of the scaling factor). Either way, observations whose error variance is large are given low weight, and observations for which it is small are given high weight.

3.3.1 Feasible Generalized Least Squares

In many cases it is reasonable to suppose that $\mathbf{\Omega}$ depends in a known way on a vector of unknown parameters $\boldsymbol{\gamma}$. If so, it may be possible to estimate consistently, so as to obtain $\mathbf{\Omega}(\hat{\boldsymbol{\gamma}})$. This type of procedure is called feasible generalized least squares, or feasible GLS. But we'll have to specify the error term as some function of some known variables.

A typical FGLS procedure has three steps. First, run OLS and obtain the residuals $\hat{u}_t$. Second, use those residuals to estimate the parameters γ of the assumed error structure – for example:

- **Heteroscedasticity of known form**, e.g. $\omega_t^2 = \exp(\mathbf{z}_t'\gamma)$: regress $\log(\hat{u}_t^2)$ on $\mathbf{z_t}$ to estimate γ, then set $\hat{\omega}_t^2 = \exp(\mathbf{z}_t'\hat{\gamma})$ and apply WLS as above. (The intercept of that auxiliary regression is inconsistent, because $E[\log(u_t^2/\omega_t^2)] \neq 0$ – it is about -1.27 for normal errors. This does not matter for WLS: a constant added to every $\log \hat{\omega}_t^2$ scales all weights by a common factor, which leaves $\hat{\beta}$ unchanged. It does matter if the fitted $\hat{\omega}_t^2$ are wanted as variance estimates in their own right.)

- **AR(1) serial correlation**, $u_t = \rho u_{t-1} + \epsilon_t$: regress $\hat{u}_t$ on $\hat{u}_{t-1}$ to get $\hat{\rho}$, then quasi-difference each variable as $\tilde{y}_t = y_t - \hat{\rho} y_{t-1}$ for $t = 2, \ldots, n$ (and similarly for X), so that the transformed error is approximately i.i.d.

  The first observation is where **Cochrane-Orcutt** (1949) and **Prais-Winsten** (1954) part company, and the $\mathbf{\Psi}$ framework above says which is right. For AR(1) errors $\mathbf{\Omega} = \frac{\boldsymbol{\sigma_\epsilon^2}}{\mathbf{1-\rho^2}} \left[\boldsymbol{\rho}^{|\mathbf{i-j}|}\right]$, and its Cholesky factor makes the first row of $\mathbf{\Psi}'$ equal to $\sqrt{1-\rho^2}$ times the first unit vector. Full GLS therefore keeps observation 1, rescaled as $\tilde{y}_1 = \sqrt{1-\hat{\rho}^2}\, y_1$; that is Prais-Winsten. Cochrane-Orcutt simply drops observation 1. Both are consistent and asymptotically equivalent, but Cochrane-Orcutt throws away a data point, and how much that costs takes some care. The direct weight on observation 1 is $\sqrt{1-\rho^2}$, which *shrinks* as $|\rho| \to 1$, so that row's mechanical contribution falls rather than rises. What grows is its relative importance: quasi-differencing strips variation out of the remaining rows as well, and with trending or highly persistent regressors the single untransformed level observation ends up carrying a disproportionate share of the information about the level. So the loss is worth caring about in short samples, and with trending regressors when ρ is near 1 – not because that row is weighted heavily, but because the others are weighted lightly. Prefer Prais-Winsten.

Third, run OLS on the transformed (weighted or quasi-differenced) data to get the FGLS estimator $\hat{\beta}_{FGLS}$. FGLS is consistent and asymptotically efficient under two requirements, not one: the assumed *form* of $\mathbf{\Omega}$ must be correct, and $\hat{\gamma}$ must be consistent for γ within that form – alongside the usual exogeneity, rank and moment conditions. A consistent $\hat{\gamma}$ inside a misspecified covariance model does not help. Either way FGLS is generally biased in finite samples, since $\hat{\Omega}$ depends on the first-stage OLS residuals.

3.4 Efficiency of GLS

3.4.1 The Gauss-Markov Theorem for GLS

The GLS estimator is the best linear unbiased estimator (BLUE) when $\mathbf{\Omega}$ is known. To see this, consider any other linear unbiased estimator $\tilde{\boldsymbol{\beta}} = \mathbf{C}\mathbf{y}$ for some matrix $\mathbf{C}$ satisfying $\mathbf{C}\mathbf{X} = \mathbf{I}$. Write $\mathbf{C} = (\mathbf{X}'\mathbf{\Omega}^{-1}\mathbf{X})^{-1}\mathbf{X}'\mathbf{\Omega}^{-1} + \mathbf{D}$, where $\mathbf{D}$ is the difference from the GLS weighting. Unbiasedness requires $\mathbf{C}\mathbf{X} = \mathbf{I}$, which means $\mathbf{D}\mathbf{X} = \mathbf{0}$. The variance of $\tilde{\boldsymbol{\beta}}$ is

$$\begin{aligned} \operatorname{Var}(\tilde{\boldsymbol{\beta}}) &= \mathbf{C}\,\mathbf{\Omega}\,\mathbf{C}' \\ &= (\mathbf{X}'\mathbf{\Omega}^{-1}\mathbf{X})^{-1} + \mathbf{D}\,\mathbf{\Omega}\,\mathbf{D}'. \end{aligned} \tag{3.8}$$

The cross terms vanish because $\mathbf{D}\mathbf{X} = \mathbf{0}$ implies $\mathbf{D}\,\mathbf{\Omega}\,\mathbf{\Omega}^{-1}\mathbf{X} = \mathbf{D}\mathbf{X} = \mathbf{0}$, so $\mathbf{D}\,\mathbf{\Omega}\,\mathbf{\Omega}^{-1}\mathbf{X}(\mathbf{X}'\mathbf{\Omega}^{-1}\mathbf{X})^{-1} = \mathbf{0}$. Since $\mathbf{D}\,\mathbf{\Omega}\,\mathbf{D}'$ is positive semidefinite,

$$\operatorname{Var}(\tilde{\boldsymbol{\beta}}) - \operatorname{Var}(\hat{\boldsymbol{\beta}}_{\mathbf{GLS}}) = \mathbf{D}\,\mathbf{\Omega}\,\mathbf{D}' \geq \mathbf{0}. \tag{3.9}$$

No linear unbiased estimator can have a smaller variance than GLS.

When $\mathbf{\Omega} = \boldsymbol{\sigma}^{\mathbf{2}}\mathbf{I}$, GLS reduces to OLS, and this is the ordinary Gauss-Markov theorem: OLS is BLUE under homoscedasticity and no serial correlation.

3.4.2 GLS versus OLS with Robust Standard Errors

The Gauss-Markov result says GLS is more efficient than OLS when $\mathbf{\Omega}$ is known. In practice $\mathbf{\Omega}$ is never known, and we face a choice: estimate $\mathbf{\Omega}$ and use FGLS, or use OLS with a sandwich variance estimator (White 1980; Whitney K. Newey and West 1987), as in the MLE chapter, that does not require specifying the form of $\mathbf{\Omega}$.

When the conditional mean is correctly specified and the regressors are suitably exogenous, both approaches give consistent estimates of β, and the difference is efficiency. The exogeneity needed is not always the same: with serially correlated errors, FGLS mixes adjacent periods and needs strict exogeneity, while OLS needs only contemporaneous exogeneity. If the linear model is only a best linear approximation, weighting can also change the target. For example, let $X \sim U(-1, 1)$ and $Y = X + X^2 - 1/3 + \varepsilon$, with ε mean-zero noise independent of X. The unweighted projection of Y on $(1, X)$ has intercept 0 and slope 1. With weights $1 + X/2$, the weighted covariance of X and Y is $7/20$ and the weighted variance of X is $11/36$, so the slope becomes $(7/20)/(11/36) \approx 1.15$. GLS exploits the structure of $\mathbf{\Omega}$ to weight observations optimally; OLS ignores that structure and compensates afterward with a robust variance estimate.

The tradeoff turns on how well $\mathbf{\Omega}$ can be modeled:

When the error structure is well understood — for example, heteroscedasticity that follows a known function of observables, or AR(1) serial correlation whose form is dictated by the data generating process — FGLS can deliver substantial efficiency gains. The efficiency improvement is largest when the heterogeneity in error variances is large, since GLS downweights noisy observations that OLS treats equally.

When the error structure is unknown or hard to model correctly, FGLS carries a specification risk. If $\hat{\mathbf{\Omega}}$ is badly estimated, the FGLS estimator can be less efficient than OLS, and the FGLS standard errors can be misleading because they assume the estimated error structure is correct. OLS with robust standard errors avoids this risk: the point estimates are the same regardless of $\mathbf{\Omega}$, and the sandwich variance is consistent under unknown heteroscedasticity, and under serial correlation weak enough to satisfy the HAC conditions – summable autocovariances and a bandwidth that grows slowly with the sample. "Robust" is not the same as robust to dependence of any form.

The modern default in applied work is to report OLS with robust standard errors, reserving GLS for settings where there is strong prior information about the error structure or where the efficiency gain is large enough to justify the specification risk. Panel data models are an important exception: the random effects estimator is a form of FGLS, and there the structure of $\mathbf{\Omega}$ (an equicorrelated block for each unit) is implied by the model rather than estimated freely.

3.5 Summary

When the error covariance Ω is not $\sigma^2 I$, GLS transforms the model so that the transformed errors are spherical, then applies OLS. The result, $(X'\Omega^{-1}X)^{-1}X'\Omega^{-1}y$, is the best linear unbiased estimator when Ω is known. Weighted least squares is the special case of heteroscedastic, uncorrelated errors. Feasible GLS estimates Ω from OLS residuals first. For AR(1) errors, Prais-Winsten keeps the first observation that Cochrane-Orcutt drops.

In practice Ω is unknown. FGLS is efficient only when the form of Ω is right. OLS with robust or HAC standard errors stays valid without modelling Ω, at some cost in efficiency. The modern default is OLS with robust standard errors, keeping GLS for cases where the error structure is well understood, as in the random effects model.

4 Endogeneity

```
# Packages used in this chapter
library(MASS)
library(ivreg)
```

When we discuss the linear model we assume that the regressors are exogenous. That word covers conditions of different strength – full independence, mean independence, and mere zero correlation – which the OLS chapter sets out as a ladder. What instrumental variables work with is the weakest of them, the moment condition $E[Zu] = 0$. However, there could be reasons to believe that some regressors are correlated with the error term. In that case we call those regressors endogenous.

Under the classical assumptions OLS estimators are unbiased and consistent. One key assumption is that the regressors have to be uncorrelated with the error term. If this condition does not hold, OLS estimators are biased and inconsistent.

When one independent variable does not satisfy this condition, we say this variable is endogenous. The estimates of other coefficients are then usually biased as well, but not always: the contamination travels through the correlation between regressors, so a coefficient on a regressor orthogonal to the endogenous one is left alone. With an orthogonal design the damage stays in the endogenous coefficient.

The most popular cure for endogeneity is to use instrumental variables.

4.1 Instrumental Variables (IV)

The basic idea of IV is to use an exogenous variable (or exogenous variables) which is correlated with the endogenous independent variable as an "instrument" for the endogenous variable.

Suppose the linear regression model

$$\mathbf{y} = \mathbf{X}\boldsymbol{\beta} + \mathbf{u}, \quad \mathbf{E}(\mathbf{u}\mathbf{u}') = \boldsymbol{\sigma}^{\mathbf{2}}\mathbf{I}, \tag{4.1}$$

at least one of the explanatory variables in the $n \times k$ matrix $\mathbf{X}$ is assumed not to be predetermined with respect to the error terms, or say, endogenous.

Suppose we have a set of variables $\mathbf{Z}$, an $n \times l$ matrix of instruments , which satisfies the moment condition

$$\mathrm{E}[\mathbf{Z}'(\mathbf{y} - \mathbf{X}\boldsymbol{\beta})] = \mathbf{0}. \tag{4.2}$$

That is, $\mathbf{Z}$ is uncorrelated with the error term.

Here is what we do for two-stage least squares (2sls):

Stage 1: Regress each of the variables in the $\mathbf{X}$ matrix on $\mathbf{Z}$ to obtain a matrix of fitted values $\hat{\mathbf{X}}$,

$$\hat{\mathbf{X}} = \mathbf{Z}(\mathbf{Z}'\mathbf{Z})^{-1}\mathbf{Z}'\mathbf{X} = \mathbf{P_Z X} \tag{4.3}$$

This is essentially to get the part of $\mathbf{X}$ that is correlated with $\mathbf{Z}$. Or to say, to project $\mathbf{X}$ on to $\mathbf{Z}$.

Stage 2: Regress $\mathbf{y}$ on $\hat{\mathbf{X}}$ to obtain the estimated $\boldsymbol{\beta}$

$$\hat{\boldsymbol{\beta}}_{\mathbf{2sls}} = (\hat{\mathbf{X}}'\hat{\mathbf{X}})^{-1}(\hat{\mathbf{X}}'\mathbf{y}) = (\mathbf{X}'\mathbf{P_Z X})^{-1}(\mathbf{X}'\mathbf{P_Z y}) = \hat{\boldsymbol{\beta}}_{\mathbf{IV}} \tag{4.4}$$

Standard errors:

$$Var[\hat{\boldsymbol{\beta}}_{\mathbf{2sls}}] = \hat{\boldsymbol{\sigma}}^{\mathbf{2}}(\hat{\mathbf{X}}'\hat{\mathbf{X}})^{-1} = \hat{\boldsymbol{\sigma}}^{\mathbf{2}}(\mathbf{X}'\mathbf{P_Z X})^{-1} \tag{4.5}$$

where $\hat{\sigma}^2 = \hat{\mathbf{u}}'\hat{\mathbf{u}}/N$. This is the homoskedastic large-sample variance; finite-sample formulas typically apply a degrees-of-freedom correction ($N - k$ in place of N), and when errors are heteroskedastic or clustered the robust/clustered IV covariance matrix replaces $\hat{\sigma}^2(X'P_Z X)^{-1}$ with a sandwich form.

Note that $\hat{\mathbf{u}} = \mathbf{y} - \mathbf{X}\hat{\boldsymbol{\beta}}_{\mathbf{2sls}}$.

If we do it manually by two steps, the second step will report a wrong standard error. That is because the second stage regression will report standard errors based on $\hat{\mathbf{u}} = \mathbf{y} - \hat{\mathbf{X}}\hat{\boldsymbol{\beta}}_{\mathbf{2sls}}$. Therefore, it's always recommended to ask the statistical program to do a 2sls for you, since presumably that will give you correct standard errors.

Geometry Illustration:




Figure 4.1: IV geometry: the instrument z is perpendicular to the error u, so projecting x and y onto z isolates the exogenous variation. The IV estimator $\hat{\beta}_{IV} = \hat{y}_z/\hat{x}_z$ recovers β without using the endogenous component $x^{\parallel}$.

Suppose the simplest case: $y = \beta_0 + \beta_1 x + u$. Take all variables in deviations from their means, so the intercept drops out and $z'\iota = 0$; this keeps the algebra below about the slope alone. Split x into the part that lies along the error and the part that does not,

$$x = x^{\perp} + x^{\parallel}, \tag{4.6}$$

where $x^{\parallel}$ is the orthogonal projection of x onto u and $x^{\perp}$ is what is left over. Two facts hold *by construction* and are what make the picture work: $x^{\perp}$ is perpendicular to u, and $x^{\perp}$ is perpendicular to $x^{\parallel}$. Endogeneity is simply the statement that $x^{\parallel} \neq 0$. (This is a decomposition of x, not an extra assumption about it – every x admits it.)

Now let z be a vector perpendicular to u with a non-zero projection onto x (relevance: z actually predicts the regressor). Project both y and x onto z, and write $\hat{y}_z$ and $\hat{x}_z$ for those two projections. The instrumental variables estimator is the ratio of their signed projection coefficients – signed, because lengths are never negative and β_1 may be,

$$\hat{\beta}_{IV} = \frac{\hat{y}_z}{\hat{x}_z} = \frac{z'y}{z'x}, \tag{4.7}$$

since $z'y = \beta_1 z'x + z'u$. The figure draws z exactly perpendicular to u, which sets $z'u = 0$ and removes the second term. Read that as the idealisation it is. Exogeneity is a statement about the population, $E[z_i u_i] = 0$, and it does not make $z'u$ vanish in a realised sample. What we have in a sample of size n is

$$\hat{\beta}_{IV} = \beta_1 + \frac{z'u}{z'x}, \tag{4.8}$$

so $\hat{\beta}_{IV}$ is consistent because $n^{-1}z'u \to 0$ while $n^{-1}z'x$ converges to something non-zero – not because the numerator is identically zero. The geometry shows why IV works; it does not make IV exact.

Staying inside the idealised picture for a moment: because the two right triangles involved are similar, $\hat{\beta}_{IV}$ is the same number you would get from the *infeasible* regression of y on the exogenous part alone: that regression returns

$$\frac{x^{\perp\prime}y}{x^{\perp\prime}x^{\perp}} = \beta_1 \frac{x^{\perp\prime}x}{x^{\perp\prime}x^{\perp}} + \frac{x^{\perp\prime}u}{x^{\perp\prime}x^{\perp}} = \beta_1, \tag{4.9}$$

where the second term vanishes because $x^{\perp} \perp u$ and the first collapses to β_1 because $x^{\perp} \perp x^{\parallel}$, so $x^{\perp\prime}x = x^{\perp\prime}x^{\perp}$. That infeasible regression is the one we would run if we could observe the decomposition. We cannot – which is exactly why we need the instrument.

```
## DGP: data$y <- data$x + data$z + data$u
library(MASS)
set.seed(66)
nobs=10000
nDim = 3
sdxx = 1
sdww=1
```

```
sdzz=1

## here we have three variables x,z,w.
## z is the omitted variable,x and z are correlated,  w is the instrument, which
 ↪  is correlated with x, but not z.  u is independent of everything else.
crxz=.6
crzw=0
crxw=.5

covarMat = matrix( c(sdxx^2, crxz, crxw, crxz, sdzz^2, crzw,  crxw, crzw, sdww^2
 ↪  ) , nrow=nDim , ncol=nDim )
covarMat
```

```
     [,1] [,2] [,3]
[1,]  1.0  0.6  0.5
[2,]  0.6  1.0  0.0
[3,]  0.5  0.0  1.0
```

```
data  = data.frame(mvrnorm(n=nobs, mu=rep(0,nDim), Sigma=covarMat ))
names(data) <- c('x','z','w')
data$u <- rnorm(nobs,0,1)
# dgp
data$y <- data$x +  data$z + data$u

lm <- lm(y~x, data=data)
lm.full <- lm(y~ x + z, data=data)
tsls.model <- ivreg(y ~ x | w, data=data)
# lm is biased
summary(lm)
```

```
Call:
lm(formula = y ~ x, data = data)

Residuals:
    Min      1Q  Median      3Q     Max
-5.3506 -0.8612  0.0123  0.8504  4.9096

Coefficients:
             Estimate Std. Error t value Pr(>|t|)
(Intercept) -0.004284   0.012843  -0.334    0.739
x            1.596114   0.013079 122.035   <2e-16 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 1.284 on 9998 degrees of freedom
```

```
Multiple R-squared:  0.5983,    Adjusted R-squared:  0.5983
F-statistic: 1.489e+04 on 1 and 9998 DF,  p-value: < 2.2e-16
```

```
# lm.full is good
summary(lm.full)
```

```

Call:
lm(formula = y ~ x + z, data = data)

Residuals:
    Min      1Q  Median      3Q     Max
-3.7267 -0.6633 -0.0179  0.6742  3.6018

Coefficients:
            Estimate Std. Error t value Pr(>|t|)
(Intercept) 0.001857   0.009973   0.186    0.852
x           0.987181   0.012629  78.166   <2e-16 ***
z           1.017559   0.012542  81.129   <2e-16 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 0.997 on 9997 degrees of freedom
Multiple R-squared:  0.7578,    Adjusted R-squared:  0.7577
F-statistic: 1.564e+04 on 2 and 9997 DF,  p-value: < 2.2e-16
```

```
# tsls is good.
summary(tsls.model)
```

```

Call:
ivreg(formula = y ~ x | w, data = data)

Residuals:
      Min        1Q    Median        3Q       Max
-6.144530 -0.962464  0.003414  0.944728  5.340560

Coefficients:
            Estimate Std. Error t value Pr(>|t|)
(Intercept)  0.01153    0.01433   0.804    0.421
x            0.95129    0.03007  31.637   <2e-16 ***

Diagnostic tests:
                 df1  df2 statistic p-value
Weak instruments   1 9998    3074.7  <2e-16 ***
Wu-Hausman         1 9997     807.8  <2e-16 ***
Sargan             0   NA        NA      NA
```

```
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 1.431 on 9998 degrees of freedom
Multiple R-Squared: 0.5007, Adjusted R-squared: 0.5006
Wald test:  1001 on 1 and 9998 DF,  p-value: < 2.2e-16
```

4.2 Control Function Approach

A second way to do an IV regression is also two-step approach. Write $\mathbf{X}_{en}$ for the endogenous columns of $\mathbf{X}$. Regress them on $\mathbf{Z}$ and keep the residual: $\hat{\mathbf{v}} = \mathbf{X_{en}} - \mathbf{Z}(\mathbf{Z}'\mathbf{Z})^{-1}\mathbf{Z}'\mathbf{X_{en}}$. Then regress y on all of $\mathbf{X}$ and $\hat{v}$ to get $\hat{\beta}_{IV}$. The exogenous columns of $\mathbf{X}$ are already in $\mathbf{Z}$, so their first-stage residuals would be zero and add nothing.

The difference between this approach and the 2sls approach is that in 2sls we regress y on $\hat{\mathbf{X}}$; in the control function approach, we regress y on $\mathbf{X}$ and $\hat{v}$. They should both give you the same coefficient estimates. There are advantages of using the control function approach.

An example to show how to do 2sls manually, or using control function approach.

```
## DGP: data$y <- data$x + data$z + data$u

lm1 <- lm(y~x, data=data)
lm2 <- lm(x~w, data=data)
tsls.manual <- lm(data$y ~ lm2$fitted.values)
summary(tsls.manual)
```

```
Call:
lm(formula = data$y ~ lm2$fitted.values)

Residuals:
    Min      1Q  Median      3Q     Max
-8.6485 -1.3401 -0.0203  1.3543  7.6974

Coefficients:
                  Estimate Std. Error t value Pr(>|t|)
(Intercept)        0.01153    0.01977   0.583     0.56
lm2$fitted.values  0.95129    0.04147  22.937   <2e-16 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 1.974 on 9998 degrees of freedom
Multiple R-squared:  0.04999,   Adjusted R-squared:  0.04989
F-statistic: 526.1 on 1 and 9998 DF,  p-value: < 2.2e-16
```

```r
# control function approach
tsls.control <- lm(data$y ~ data$x + lm2$residuals)
summary(tsls.control)
```

```
Call:
lm(formula = data$y ~ data$x + lm2$residuals)

Residuals:
    Min      1Q  Median      3Q     Max
-5.1997 -0.8329  0.0131  0.8370  4.7457

Coefficients:
              Estimate Std. Error t value Pr(>|t|)
(Intercept)    0.01153    0.01237   0.932    0.351
data$x         0.95129    0.02594  36.670   <2e-16 ***
lm2$residuals  0.84313    0.02966  28.423   <2e-16 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 1.235 on 9997 degrees of freedom
Multiple R-squared:  0.6284,    Adjusted R-squared:  0.6283
F-statistic:  8451 on 2 and 9997 DF,  p-value: < 2.2e-16
```

The coefficient on `data$x` is the 2SLS estimate, but its default OLS standard error is not the 2SLS standard error. It takes the residual variance from $y - X\hat{\beta} - \hat{v}\hat{\gamma}$, which is smaller than the structural residual variance, and it ignores that $\hat{v}$ is estimated. Here it reports 0.026 against the 2SLS standard error of 0.030. For inference on β, use the 2SLS standard error or bootstrap both stages.

This approach can be used to do a simple endogeneity test. First, you can do a simple test of endogeneity of $\mathbf{X}$. For example:

```
reg x_endog x* z*
predict v_x, resid
reg y x_endog x* v_x
test v_x
```

Obviously, this test is based on $\mathbf{Z}$ being exogenous.

Residual inclusion extends to *some* non-linear models – logit, probit, Poisson – under additional control-function assumptions, but it is not automatically valid for "any glm". The construction is model-specific: the appropriate first-stage residual (a plain linear residual or a generalized residual), the identification conditions, and the standard errors all differ for continuous, binary, and nonlinear endogenous regressors, and inserting a linear first-stage residual into an arbitrary GLM is generally not correct. The generated-regressor step also inflates uncertainty, so standard errors need to account for the estimated first stage (e.g. by bootstrap). For example, the "eteffects" procedure in Stata (version 14) uses a control-function approach to get endogenous treatment effects for several outcome types.

4.3 Durbin-Wu-Hausman Test

4.3.1 Idea

In econometric modeling, there are often questions on endogeneity. Do we know how to test whether an independent variable is endogenous statistically? The answer is: sort of, but not really. We cannot do endogeneity test without a valid instrument. Therefore, we have to have strong argument for a valid instrument first before we can do endogeneity test.

With endogenous variables on the right-hand side of the equation, we need to use instrumental variable (IV) regression for consistent estimation. However, with IV regression, we lose efficiency: the asymptotic variance of the IV estimator is larger, and can be much larger than the OLS estimator. Therefore, we gain consistency, but lose efficiency, by using IV estimator when there is an endogeneity problem.

Now we have a familiar scenario (if you are familiar with Hausman test for fixed effect and random effect estimator for panel data): Suppose we have the null hypothesis as the regressor being exogenous. We have an efficient estimator under null hypothesis yet inconsistent under alternative hypothesis (OLS estimator). We also have a consistent estimator under both null and alternative (IV estimator).

Similar to panel data setting, we have the Hausman test statistic as:

$$H = (\hat{\beta}_c - \hat{\beta}_e)' D^{-} (\hat{\beta}_c - \hat{\beta}_e) \tag{4.10}$$

where $D = \text{Var}[\hat{\beta}_c] - \text{Var}[\hat{\beta}_e]$, $^{-}$ is the generalized inverse, $\hat{\beta}_c$ is the consistent estimator (in this case the IV estimator) and $\hat{\beta}_e$ is the efficient estimator (in this case OLS estimator).

H conforms to χ^2_k asymptotically, where k is the number of endogenous variables.

This test is to compare the IV estimator and the OLS estimator: if it's close, then OLS estimator is fine (fail to reject null that OLS is consistent, or say the variable is exogenous). If it's large, then IV estimator is needed, although we lose some efficiency. This test is based on the assumption that the instruments are exogenous. If that is in question, then it's pointless to do the test, since the IV estimator cannot guarantee consistency either.

4.3.2 Implementation in Stata

In Stata, there are different ways to do it:

Do a regular Hausman test:

```
ivreg y x1 (x2=x3 x4)
estimates store iv
reg y x1 x2
hausman iv ., constant sigmamore
```

Or, simply use "ivendog" in Stata.

4.4 Identification

Identification in a regression equation means that all parameters can be uniquely estimated. The relevant counting rule (the *order condition*) is most cleanly stated in terms of *excluded* instruments. Let k_e be the number of endogenous regressors and l_e the number of excluded instruments (instruments not already included as exogenous regressors). A necessary condition is $l_e \geq k_e$: there must be at least as many excluded instruments as endogenous regressors. (Equivalently, counting the included exogenous regressors as their own instruments, the total number of instruments must be at least the total number of regressors.) If $l_e = k_e$ we have exact-identification. If $l_e > k_e$, we have over-identification.

Additional valid instruments that carry independent information can improve asymptotic precision under a suitable weighting scheme. The gain is not automatic: a redundant instrument adds nothing, a weak one need not help, and a large number of instruments worsens finite-sample behaviour even when every one of them is valid. The other advantage of over-identification is that over-identification tests can be done to test the adequacy of instruments.

Under the null hypothesis that all the instruments are uncorrelated with the error term, an LM statistic $N \times R^2$ conforms to $\chi^2(r)$ distribution, $r = l_e - k_e$, the number of excess instruments, or say, the number of overidentifying restrictions. If we reject the null, then we should be concerned about the exogeneity of the whole set of the instruments. This test is called Sargan's (1958) test in IV context, and (Hansen's) J test in GMM context. The two names are not quite interchangeable: Sargan's $N \times R^2$ form assumes homoskedastic errors, while Hansen's J, built from the efficient GMM weighting, is the heteroskedasticity-robust counterpart and the one to use whenever the standard errors are robust.

It is worth being exact about what the test delivers, because it is easy to over-read. It is a joint test of the overidentifying restrictions. Failing to reject does not certify the instruments: the test has no power against violations that all of them share, because instruments wrong in the same way agree on the same wrong answer. Rejection, for its part, does not say *which* moment condition failed. A passing J is the absence of one particular kind of evidence against the instruments, not evidence for them.

What the J test or Sargan's test does is to test the whole set of instruments being exogenous or not. There is another test for testing exogeneity for a subset of instruments. It's call a C test or a difference-in-Sargan test. The idea is to calculate the difference between two Sargan's statistics (or Hansen's J in GMM setting); one is with the whole set of instruments, the other one without the suspected instruments. The null is that the suspect instruments are exogenous; or orthogonal to the error term. Obviously to conduct the C test, we'll have to have at least one extra instrument more than the number of endogenous variables.

To understand it better, we look at how to implement Sargan's test manually: For the 2SLS estimator, the test statistic is Sargan's statistic, typically calculated as $N \times R^2$ from a regression of the IV residuals on the full set of instruments.

```
. ivregress 2sls rent pcturban (hsngval = faminc i.region)
. estat overid

  Tests of overidentifying restrictions:
```

```
  Sargan (score) chi2(3) =  11.2877  (p = 0.0103)
  Basmann chi2(3)        =  12.8294  (p = 0.0050)

. predict res, residual

. reg res pcturban faminc i.region

. disp e(N)*e(r2)
11.287665
```

4.4.1 Implementation in Stata

In Stata, there are different ways to do over-identification test, ivreg2 reports a comprehensive set of tests; overid command does the over-identification test after the ivreg command.

ivreg2 with gmm option returns J test; it reports Sargan's test without this option.

ivreg2 also reports the C test statistic, with ortho(). If the C test rejects and the J test without the suspect instruments does not, the evidence points at the suspect instruments – conditional on the retained ones being valid. That condition does real work: a difference test cannot tell an invalid suspect from an invalid maintained instrument, so it locates the problem only when the maintained set is sound.

4.5 Weak Instruments

4.5.1 Problem with the cure

An instrument needs to satisfy two criteria: orthogonality and relevance. We need instruments to be orthogonal to the error term. With extra instruments, Sargan's test lets us test the joint compatibility of the overidentifying restrictions. As stressed above, that is a test of the set as a whole and not a verification of orthogonality: passing it leaves the instruments untested in every direction they share.

It turns out instrument relevance is important too: if instruments are weak, the usual large-sample approximations are poor. It helps to keep three cases apart. With fixed, nonzero first-stage relevance, IV remains *consistent* but can be badly *biased* in finite samples (and biased toward OLS), with the normal approximation working poorly. Under weak-instrument (local-to-zero) asymptotics, the limiting distribution is *nonstandard* and not centered in the usual way, so conventional standard errors and t-statistics mislead. Only in the limiting case of complete irrelevance ($\Pi = 0$) is the parameter unidentified.

To see the problem, suppose

$$\mathbf{y} = \mathbf{X}\boldsymbol{\beta} + \mathbf{u}, \quad \mathbf{E}(\mathbf{u}\mathbf{u}') = \boldsymbol{\sigma}_{\mathbf{u}}^{\mathbf{2}}\mathbf{I}, \tag{4.11}$$

$$\mathbf{X} = \mathbf{Z}\boldsymbol{\Pi} + \mathbf{v}, \quad \mathbf{E}(\mathbf{v}\mathbf{v}') = \boldsymbol{\sigma}_{\mathbf{v}}^{\mathbf{2}}\mathbf{I}, \tag{4.12}$$

and

$$\mathrm{E}(\mathbf{Zu}) = \mathbf{0}. \tag{4.13}$$

We can see here $\mathbf{Z}$ is exogenous. However, the model does not say anything about relevance. To illustrate the problem caused by weak instrument, suppose we have only one endogenous variable and one instrument.

$$\hat{\beta}_{2sls} = \frac{\mathbf{Z'y}}{\mathbf{Z'X}} = \frac{\mathbf{Z'}(\mathbf{X}\boldsymbol{\beta} + \mathbf{u})}{\mathbf{Z'X}} = \beta + \frac{\mathbf{Z'u}}{\mathbf{Z'X}}. \tag{4.14}$$

If $\mathbf{Z}$ is irrelevant, or, $\Pi = 0$, then

$$\hat{\beta}_{2sls} - \beta = \frac{\mathbf{Z'u}}{\mathbf{Z'v}} = \frac{\frac{1}{\sqrt{n}}\sum_{i=1}^{n} Z_i u_i}{\frac{1}{\sqrt{n}}\sum_{i=1}^{n} Z_i v_i} \xrightarrow{d} \frac{z_u}{z_v}, \tag{4.15}$$

where (normalizing $\mathrm{Var}(Z_i) = 1$, which is harmless since any common scale factor cancels in the ratio)

$$\begin{bmatrix} z_u \\ z_v \end{bmatrix} \sim N(0, \begin{bmatrix} \sigma_u^2 & \sigma_{uv} \\ \sigma_{uv} & \sigma_v^2 \end{bmatrix}). \tag{4.16}$$

Therefore, if $\mathbf{Z}$ is irrelevant, β_{2sls} is inconsistent. Also, the distribution of the bias is Cauchy-like (the ratio of correlated normals).

This is a case where the cure might be worse than the disease itself: the bias can be big comparing to the bias an OLS estimate suffers.

4.5.2 Weak Instrument Tests

There are a variety of weak-instruments tests proposed. Most of them are based on so-called weak-instruments asymptotics and a new parameter called "concentration parameter" $\mu^2 = \Pi' Z' Z \Pi / \sigma_v^2$. Sample size only enters the distribution through μ^2.

With weak-instruments asymptotics, IV estimators are no longer consistent, and they are not normal asymptotically. Most test statistics (J test, etc.) do not have normal or χ^2 distributions anymore.

Now I list the following tests in the order of recommended level by James Stock:

1. Moreira (2003) conditional likelihood ratio test (CLR).

Advantages of this test: a. Nearly optimal among invariant similar tests (Andrews, Moreira, and Stock 2006). b. Implemented in Stata as condivreg.

Disadvantages:

a. Complicated. Only developed so far for one endogenous variable case.

2. Stock-Yogo bias method and size method.

Stock and Yogo (2005) provide critical values for both methods: one is to control the size of bias, the other one is to control the size of a Wald test of $\beta = \beta_0$. Bias method is more frequently used. In the case of multiple endogenous variables, the Cragg-Donald statistics is used to compare with the critical values. It is implemented in Stata as part of the *ivreg2* command. The two tables have different coverage. The 2SLS relative-bias critical values exist only when there are at least two overidentifying restrictions, $l_e - k_e \geq 2$ (three excluded instruments for one endogenous regressor), because the finite-sample mean of 2SLS exists only at that degree of overidentification. The size critical values start from exact identification, $l_e = k_e$.

3. Anderson-Rubin confidence intervals.

In the model of

$$\mathbf{y} = \mathbf{X}\boldsymbol{\beta} + \mathbf{u}, \quad \mathbf{E}(\mathbf{u}\mathbf{u}') = \boldsymbol{\sigma}_{\mathbf{u}}^{\mathbf{2}}\mathbf{I}, \tag{4.17}$$

$$\mathbf{X} = \mathbf{Z}\boldsymbol{\Pi} + \mathbf{v}, \quad \mathbf{E}(\mathbf{v}\mathbf{v}') = \boldsymbol{\sigma}_{\mathbf{v}}^{\mathbf{2}}\mathbf{I}, \tag{4.18}$$

The null hypothesis $H_0 : \beta = \beta_0$. Anderson-Rubin statistic is the F statistic in the regression of $y - X\beta_0$ on Z, the F test that the coefficient vector on Z is zero. Under the model that coefficient equals $\Pi(\beta - \beta_0)$, which is zero exactly when $\beta = \beta_0$ provided Π has full column rank. Without that rank condition the test stays valid and weak-instrument robust, but it cannot distinguish β_0 from an alternative differing in a direction Π annihilates:

$$AR(\beta_0) = \frac{(y - X\beta_0)' P_Z (y - X\beta_0)/l}{(y - X\beta_0)' M_Z (y - X\beta_0)/(N - l)} \tag{4.19}$$

where l is the number of instruments. The idea of AR confidence interval is to collect all values of β_0 at which we fail to reject $H_0 : \beta = \beta_0$.

4. First-stage F test.

The familiar rule of thumb is $F > 10$, and its scope is worth knowing. It comes from the single-endogenous-regressor case with homoskedastic errors, where it corresponds roughly to a bound on the relative bias of 2SLS. The formal Stock-Yogo critical values depend on how many instruments there are and on which criterion – relative bias or size distortion – is being controlled, so one number does not carry over to every design. Under heteroskedasticity or clustering the relevant statistic is the effective first-stage F of Montiel Olea and Pflueger (2013) rather than the conventional one. And when identification is weak the dependable route is not a higher threshold but weak-instrument-robust inference – the Anderson-Rubin and CLR tests in this list, whose size does not depend on instrument strength.

5. Kleibergen's LM test.

This test is dominated by the CLR test, thus no longer the optimal test to use.

6. First-stage R^2, or partial R^2, etc., are not recommended.

4.6 When is endogeneity a problem and what can we say from an IV regression

Let's consider a model of housing price on no. of rooms and square footage.

$$P = \alpha R + \beta F + \epsilon \tag{4.20}$$

Should square footage really be there? We are not interested in, given square footage, how much money we need to pay for an additional room. We are generally interested in how much money we need to pay for an additional footage. However, if we do not include footage, then we "sort of" have endogeneity problem due to omitted variable.

Another example would be education's impact on wage. Say IQ is an omitted variable. Do we want to say something about education's impact on wage controlling on IQ, or not?

So it depends on what questions to ask. If I am a builder and ask what an extra bedroom on the already built house would bring to me, I need to control for footage. If I am generally asking what an extra bedroom would cost, then I am implying what that bedroom and associated footage would cost me. In that case, without controlling for footage is not a problem, since the question is: what an additional bedroom (AND the associated average square footage) would cost me.

In the wage and education example, if I am asking: what's the effect of education on wage? I implicitly ask what would be my pay raise if I go to college (given IQ, of course), for example. Then we are concerned that this model of

$$wage = \alpha edu + \epsilon \tag{4.21}$$

will generate biased estimate of α, because we are not controlling for IQ. But if I am an employer and interested in hiring a person, what would I have to pay extra to hire a college graduate vs. a high school graduate? Then I don't have to include that IQ variable, because implicitly I am asking: what extra I need to pay for that college graduate who comes with a higher IQ, and everything else that is associated with higher education.

In addition, what an IV regression really gives us is part of the causal inference of X (suspected to be endogenous) on y, namely the part induced by Z. For example, we are interested in the relationship between college education and wage. If we use distance to college as an instrument, then our inference is the effect of college education from those who decide to go because of proximity of the college.

This is the Local Average Treatment Effect (LATE) of Imbens and Angrist (1994): with a binary instrument and heterogeneous treatment effects, IV identifies the causal effect for *compliers* — those whose treatment status is shifted by the instrument — rather than for the full population. The formal derivation, including the monotonicity assumption and the complier/always-taker/never-taker decomposition, is in the Causal Econometrics Guide.

4.7 Summary

A regressor is endogenous when it is correlated with the error, and then OLS is biased and inconsistent. Instrumental variables use variables that are uncorrelated with the error and predict the regressor. 2SLS projects the regressors onto the instruments and regresses the outcome on the projection. The control function adds the first-stage residual to the regression and gives the same coefficients. Its coefficient on the residual is the Durbin-Wu-Hausman test of exogeneity, which is only as good as the instruments behind it.

Identification needs at least as many excluded instruments as endogenous regressors. With more, the Sargan or Hansen J test checks the overidentifying restrictions jointly. It cannot certify the instruments, because it has no power against a violation they all share. Weak instruments make IV biased toward OLS, with nonstandard distributions. The conditional likelihood-ratio test and Anderson-Rubin intervals stay valid when instruments are weak, and a first-stage F is only a rough screen. With heterogeneous effects, IV identifies a LATE, the effect for the compliers.

5 Generalized Method of Moments (GMM)

```
# Packages used in this chapter
library(MASS)
library(ivreg)
library(gmm)
library(sandwich)
```

5.1 The Method of Moments (MOM)

A population moment γ can be defined as the expectation of some continuous function g of a random variable x:

$$\gamma = \mathrm{E}[g(x)] \tag{5.1}$$

On the other hand, a sample moment is the sample version of the population moment in a particular sample:

$$\hat{\gamma} = \frac{1}{n}\sum[g(x)] \tag{5.2}$$

5.2 OLS as a moment problem

Consider the simple linear regression

$$\mathbf{y} = \mathbf{X}\boldsymbol{\beta} + \mathbf{u}, \quad \mathbf{u} \sim \mathbf{IID}(\mathbf{0}, \boldsymbol{\sigma^2}). \tag{5.3}$$

If the regressors are exogenous, uncorrelated with the error term, then

$$\mathbf{E}(\mathbf{X}'\mathbf{u}) = \mathbf{0}. \tag{5.4}$$

The MOM principle suggests that we replace the left-hand side with its sample analog $\frac{1}{n}\mathbf{X}'(\mathbf{y} - \mathbf{X}\boldsymbol{\beta})$.

Since we know that the true $\boldsymbol{\beta}$ sets the population moment equal to zero in expectation, it seems reasonable to assume that a good choice of $\hat{\boldsymbol{\beta}}$ would be one that sets the sample moment to zero. The MOM procedure suggests an estimate of $\boldsymbol{\beta}$ that solves

$$\frac{1}{n}\mathbf{X}'(\mathbf{y} - \mathbf{X}\hat{\boldsymbol{\beta}}) = \mathbf{0}. \tag{5.5}$$

The MOM estimator is

$$\hat{\boldsymbol{\beta}} = (\mathbf{X}'\mathbf{X})^{-\mathbf{1}}\mathbf{X}'\mathbf{y}, \tag{5.6}$$

which is the same as the OLS estimator.

5.3 IV as a moment problem

Consider the simple linear regression

$$\mathbf{y} = \mathbf{X}\boldsymbol{\beta} + \mathbf{u}, \quad \mathbf{u} \sim \mathbf{IID}(\mathbf{0}, \boldsymbol{\sigma}^{\mathbf{2}}). \tag{5.7}$$

If one or more of the regressors is endogenous, then

$$\mathbf{E}(\mathbf{X}'\mathbf{u}) \neq \mathbf{0}. \tag{5.8}$$

We have to find an instrumental variable $\mathbf{Z}$ which is

$$\mathbf{E}(\mathbf{Z}'\mathbf{u}) = \mathbf{0}. \tag{5.9}$$

Or,

$$\mathbf{E}(\mathbf{Z}'(\mathbf{y} - \mathbf{X}\boldsymbol{\beta})) = \mathbf{0}. \tag{5.10}$$

The sample analogy of this is

$$\frac{1}{n}\mathbf{Z}'(\mathbf{y} - \mathbf{X}\hat{\boldsymbol{\beta}}) = \mathbf{0}. \tag{5.11}$$

When the model is *exactly identified* – the number of instruments equals the number of regressors, so $\mathbf{Z}'\mathbf{X}$ is square and nonsingular – the sample moment condition has a unique solution, the simple IV estimator

$$\hat{\boldsymbol{\beta}} = (\mathbf{Z}'\mathbf{X})^{-\mathbf{1}}\mathbf{Z}'\mathbf{y}. \tag{5.12}$$

When the model is *overidentified* – more instruments than regressors – $\mathbf{Z}'\mathbf{X}$ is no longer square and we cannot set all sample moments to zero simultaneously. We instead minimize a quadratic form in the moments, which gives the two-stage least squares (2SLS) estimator

$$\hat{\boldsymbol{\beta}} = (\mathbf{X}'\mathbf{P}_{\mathbf{Z}}\mathbf{X})^{-\mathbf{1}}\mathbf{X}'\mathbf{P}_{\mathbf{Z}}\mathbf{y}, \qquad \mathbf{P}_{\mathbf{Z}} = \mathbf{Z}(\mathbf{Z}'\mathbf{Z})^{-\mathbf{1}}\mathbf{Z}', \tag{5.13}$$

the special case of the GMM estimator below with weighting matrix $(\mathbf{Z}'\mathbf{Z})^{-\mathbf{1}}$.

5.4 The Generalized Method of Moments

The expectation $\mathrm{E}(Y^r)$ for any $r = 1, 2, ...$ is called the rth (raw) moment of Y. The expectation $\mathrm{E}[(Y - \mathrm{E}(Y))^r]$ is called the rth centered moment of Y.

The mean is the first raw moment.

The variance is the second centered moment.

The third centered moment measures the skewness of the distribution.

The fourth centered moment measures the kurtosis of the distribution. Interpreted as a measure of "fatness of tails".

The standardized kurtosis is

$$k = \frac{E[(Y - E(Y))^4]}{E[(Y - E(Y))^2]^2}. \tag{5.14}$$

For a normal distribution, $k = 3$.

For a t distribution with $v > 4$ degrees of freedom, $k = 3 + 6/(v-4) > 3$, i.e., the t distribution has fatter tails than a normal distribution (the excess kurtosis $6/(v-4)$ shrinks toward 0 as v grows).

The distribution function of a random variable captures all information about the random variable. If the moment generating function exists in a neighborhood of zero, the full set of moments also determines the distribution (without such a condition this can fail: the lognormal distribution is not determined by its moments).

This matters directly for GMM: GMM only ever matches a finite set of moment conditions, and if the underlying distribution is not uniquely pinned down by its (even infinite) sequence of moments, then no amount of moment-matching – however many moment conditions we add – can recover the full data-generating distribution. This is precisely the sense in which GMM asks for less than MLE: MLE assumes we know the full distributional family and estimates its parameters (recovering the whole distribution when correctly specified), while GMM only requires correctly specified moment conditions and only ever identifies the parameters entering those moments, not the full distribution. When the distributional assumption behind MLE is correct, MLE is efficient and gives you more (the whole distribution); when it is not, GMM's weaker requirements make it more robust.

This distinction underlies the relative strengths and weaknesses of ML and GMM.

5.5 GMM

The statistical model takes the general form

$$E[m(Y_i; \theta_0)] = 0 \tag{5.15}$$

where - $Y_1, \cdots, Y_n$ are random variables from which the sample $y_1, \cdots, y_n$ is drawn, - $m(Y, \theta)$ is a function specifying the model, - θ_0 is the "true value" of the parameter.

$E[m(Y_i; \theta_0)] = 0$ are called the population moment conditions.

Two ideas behind GMM:

1. Replace the population mean $E[.]$ with the sample mean calculated from the observed sample $y_1, \cdots, y_n$.
2. Since $E[m(Y_i; \theta_0)] = 0$, choose $\hat{\theta}_{GMM}$ to make $\frac{1}{n}\sum_{i=1}^{n} m(y_i; \hat{\theta}_{GMM})$ as close to zero as possible.

Define the notation

$$\bar{m}(\theta) = \frac{1}{n}\sum_{i=1}^{n} m(y_i; \theta). \tag{5.16}$$

$\hat{\theta}_{GMM}$ is chosen to make $\bar{m}(\theta)'\bar{m}(\theta)$ as close to zero as possible.

More generally, $\hat{\theta}_{GMM}$ is chosen to minimize $\bar{m}(\theta)'W\bar{m}(\theta)$ for some weighting matrix W.

The choice of W matters only under overidentification, provided the just-identified system has a unique zero. With as many moment conditions as parameters, if $\bar{m}(\theta) = 0$ has exactly one solution,

that solution minimizes $\bar{m}(\theta)'W\bar{m}(\theta)$ for every positive-definite W, so W drops out. The linear IV system above has this property when its square sample matrix is nonsingular, which is why the just-identified IV estimator did not need a weighting matrix. Counting moments does not guarantee a unique zero in a nonlinear problem. The moment $E[Y] - \theta^2 = 0$ with $E[Y] = 1$ has two roots, $\theta = 1$ and $\theta = -1$, and a sample with $\bar{y} < 0$ has none. Nonlinear moments need their own argument for existence and identification, such as the sign restriction $\theta > 0$ here. When there are more moments than parameters, the efficient choice is $W = S^{-1}$, where S is the asymptotic variance of the sample moments, $S = \lim_{n\to\infty} \mathrm{Var}(\sqrt{n}\,\bar{m}(\theta_0))$. With that weighting,

$$\sqrt{n}(\hat{\theta}_{GMM} - \theta_0) \xrightarrow{d} N(0, (G'S^{-1}G)^{-1}), \qquad G = \mathrm{E}\left[\frac{\partial m(Y_i;\theta_0)}{\partial \theta'}\right], \tag{5.17}$$

which is the GMM counterpart of the information-matrix bound of the previous chapter. Since S depends on θ_0, in practice one starts from any consistent weighting (often the identity, or $(\mathbf{Z}'\mathbf{Z})^{-1}$ – scalar multiples of $\mathbf{W}$ leave the minimizer unchanged, so with $\bar{m}$ defined as a sample average this can equally be written $[n^{-1}\mathbf{Z}'\mathbf{Z}]^{-1}$), estimates $\hat{S}$ from the resulting residuals, and re-minimizes: two-step GMM. Stopping at $W = (\mathbf{Z}'\mathbf{Z})^{-1}$ is exactly 2SLS, which is therefore efficient GMM only under homoskedasticity.

5.5.1 An example

Let's see an example with GMM, using the same simulated data as before. We have the same situation as before, X is endogenous. We are doing GMM version of 2sls.

Here I use R's "gmm" library, which makes things easy. Its first two arguments are `g` and `x`: with the formula interface, `g` is the model formula whose residual forms the moment function, and `x` is a formula giving the instruments. The moment condition in this example is $E(\mathbf{z}'\mathbf{u}) = \mathbf{0}$, where $\mathbf{z}$ is the instrument – the variable `w` in the simulated data – and u is the residual from the structural equation. Note that $\mathbf{W}$ above denotes the *weighting matrix* in $\bar{m}'W\bar{m}$, which is a different object entirely, so the instrument is written $\mathbf{z}$ here.

```
## DGP: data$y <- data$x + data$z + data$u

set.seed(66)
nobs=10000
nDim = 3
sdxx = 1
sdww=1
sdzz=1

## here we have three variables x,z,w.
## z is the omitted variable,x and z are correlated,  w is the instrument, which
 ↪  is correlated with x, but not z.  u is independent of everything else.
crxz=.6
crzw=0
crxw=.5
```

```
covarMat = matrix( c(sdxx^2, crxz, crxw, crxz, sdzz^2, crzw,  crxw, crzw, sdww^2
 ↪  ) , nrow=nDim , ncol=nDim )
covarMat
```

```
     [,1] [,2] [,3]
[1,]  1.0  0.6  0.5
[2,]  0.6  1.0  0.0
[3,]  0.5  0.0  1.0
```

```
data  = data.frame(mvrnorm(n=nobs, mu=rep(0,nDim), Sigma=covarMat ))
names(data) <- c('x','z','w')
data$u <- rnorm(nobs,0,1)
# dgp
data$y <- data$x +  data$z + data$u

gmm.fit = gmm(y ~ x, x = ~w, data = data)
summary(gmm.fit)
```

```
Call:
gmm(g = y ~ x, x = ~w, data = data)


Method:  twoStep

Kernel:  Quadratic Spectral

Coefficients:
             Estimate     Std. Error   t value      Pr(>|t|)
(Intercept)   1.1531e-02   1.4151e-02   8.1481e-01   4.1518e-01
x             9.5129e-01   2.9971e-02   3.1740e+01  4.3442e-221

J-Test: degrees of freedom is 0
                J-test                P-value
Test E(g)=0:    1.16401567357162e-26  *******
```

```
# It returns the same estimates as the 2sls results.
tsls.model <- ivreg(y ~ x | w, data=data)
summary(tsls.model)
```

```
Call:
ivreg(formula = y ~ x | w, data = data)

Residuals:
      Min        1Q    Median        3Q       Max
```

```
-6.144530 -0.962464  0.003414  0.944728  5.340560

Coefficients:
            Estimate Std. Error t value Pr(>|t|)
(Intercept)  0.01153    0.01433   0.804    0.421
x            0.95129    0.03007  31.637   <2e-16 ***

Diagnostic tests:
                  df1  df2 statistic p-value
Weak instruments    1 9998    3074.7  <2e-16 ***
Wu-Hausman          1 9997     807.8  <2e-16 ***
Sargan              0   NA        NA      NA
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 1.431 on 9998 degrees of freedom
Multiple R-Squared: 0.5007, Adjusted R-squared: 0.5006
Wald test:  1001 on 1 and 9998 DF,  p-value: < 2.2e-16
```

Two things about that agreement are worth being explicit about, because it is easy to over-generalise from it. First, GMM and 2SLS coincide *numerically* here because the model is just-identified: one instrument (`w`) for one endogenous regressor (`x`), so $\mathbf{Z}'\mathbf{X}$ is square and the sample moments can be set to zero exactly, whatever the weighting matrix. The output says so – the J-test reports "degrees of freedom is 0", and its value ($\sim 10^{-26}$) is numerical noise rather than evidence. Second, `gmm()` is using an optimal two-step weighting with a heteroskedasticity-and-autocorrelation-consistent kernel (note `Method: twoStep` and `Kernel: Quadratic Spectral` in the output). Under *over*-identification that choice would generally **not** reproduce 2SLS, for exactly the reason given above: $W = (\mathbf{Z}'\mathbf{Z})^{-1}$ is the efficient weighting only under homoskedasticity.

In OLS case, it would be $E(Xu) = 0$.

```r
gmm.ols = gmm(y ~ x, x = ~x, data = data)
summary(gmm.ols)
```

```
Call:
gmm(g = y ~ x, x = ~x, data = data)


Method:  twoStep

Kernel:  Quadratic Spectral

Coefficients:
             Estimate     Std. Error   t value      Pr(>|t|)
(Intercept)  -0.0042841    0.0126399   -0.3389314    0.7346614
x             1.5961141    0.0128364  124.3424084    0.0000000
```

```
J-Test: degrees of freedom is 0
                J-test                  P-value
Test E(g)=0:    2.66671052451529e-28    *******
```

```
# It returns the same estimates as the OLS results.
ols <- lm(y ~ x, data=data)
summary(ols)
```

```

Call:
lm(formula = y ~ x, data = data)

Residuals:
    Min      1Q  Median      3Q     Max
-5.3506 -0.8612  0.0123  0.8504  4.9096

Coefficients:
             Estimate Std. Error t value Pr(>|t|)
(Intercept) -0.004284   0.012843  -0.334    0.739
x            1.596114   0.013079 122.035   <2e-16 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 1.284 on 9998 degrees of freedom
Multiple R-squared:  0.5983,    Adjusted R-squared:  0.5983
F-statistic: 1.489e+04 on 1 and 9998 DF,  p-value: < 2.2e-16
```

5.5.2 Overidentification, the weighting matrix, and the J test

The example above cannot show what GMM adds, because it is just-identified. One instrument for one endogenous regressor leaves nothing for the weighting matrix to do and nothing for a specification test to check. Two things have to change. We need more instruments than endogenous regressors, and we need errors that are not homoskedastic – under homoskedasticity $(\mathbf{Z}'\mathbf{Z})^{-1}$ already is the efficient weighting, so 2SLS and efficient GMM agree and the comparison is empty.

The design keeps $y = x + z + u$ with z omitted from the regression, so the composite error is $z + u$ and x is endogenous through its correlation with z. There are now two instruments. Both w_1 and w_2 correlate 0.5 with x and 0.2 with each other, and neither correlates with z, so both are valid. The error is drawn with standard deviation $\exp(0.8w_1)$, which leaves $E[u \mid w] = 0$ intact while making the variance depend on an instrument. That is exactly the case where the efficient weighting differs from $(\mathbf{Z}'\mathbf{Z})^{-1}$. The true coefficient on x is 1.

```
set.seed(66)
nobs <- 10000

# cor(x,z)=.6; cor(x,w1)=cor(x,w2)=.5; cor(z,w1)=cor(z,w2)=0; cor(w1,w2)=.2
covarMat <- matrix(c(1,  .6, .5, .5,
```

```
                     .6, 1,  0,  0,
                     .5, 0,  1,  .2,
                     .5, 0,  .2, 1), nrow = 4)
d <- data.frame(mvrnorm(n = nobs, mu = rep(0, 4), Sigma = covarMat))
names(d) <- c("x", "z", "w1", "w2")

# the error variance depends on w1, so E[u|w] = 0 still holds but 2SLS is
# no longer the efficient GMM estimator
d$u <- rnorm(nobs, 0, exp(0.8 * d$w1))
d$y <- d$x + d$z + d$u

fit_ident <- gmm(y ~ x, x = ~ w1 + w2, data = d, wmatrix = "ident")
fit_eff   <- gmm(y ~ x, x = ~ w1 + w2, data = d)
tsls      <- ivreg(y ~ x | w1 + w2, data = d)

data.frame(
  estimator = c("GMM, identity weighting", "GMM, efficient two-step",
                "2SLS, classical SE", "2SLS, robust SE"),
  b_x = round(c(coef(fit_ident)["x"], coef(fit_eff)["x"],
                coef(tsls)["x"], coef(tsls)["x"]), 4),
  se = round(c(sqrt(diag(vcov(fit_ident)))["x"], sqrt(diag(vcov(fit_eff)))["x"],
               sqrt(diag(vcov(tsls)))["x"],
               sqrt(diag(vcovHC(tsls, type = "HC0")))["x"]), 4),
  row.names = NULL)
```

```
                estimator    b_x     se
1 GMM, identity weighting 1.0152 0.0463
2 GMM, efficient two-step 0.9887 0.0416
3      2SLS, classical SE 1.0157 0.0328
4         2SLS, robust SE 1.0157 0.0459
```

```
specTest(fit_eff)
```

```
 ##  J-Test: degrees of freedom is 1  ##

                J-test   P-value
Test E(g)=0:    1.64870  0.19914
```

Three readings. The weighting matrix now matters: identity weighting gives 1.015 and the efficient two-step estimator gives 0.989, where in the just-identified case every weighting gave the same number. The efficiency gain is real but modest, and it shows up only against the *robust* 2SLS standard error, 0.042 against 0.046. And the classical 2SLS standard error, 0.033, is far too small – the earlier remark about homoskedasticity, arriving as a number instead of a caveat.

The J statistic is now a test rather than a formality. It is a joint specification test of the moment conditions, not a certificate of instrument validity. It has one degree of freedom – three moment

conditions, from the constant and the two instruments, less the two parameters – and its value of 1.65 with a p-value of 0.20 gives no reason to doubt the moment conditions.

To see it earn its keep, make w_2 invalid by correlating it with the omitted z. Nothing else changes.

```
covarMat2 <- covarMat
covarMat2[2, 4] <- covarMat2[4, 2] <- 0.4   # w2 now correlates with omitted z

set.seed(66)
d2 <- data.frame(mvrnorm(n = nobs, mu = rep(0, 4), Sigma = covarMat2))
names(d2) <- c("x", "z", "w1", "w2")
d2$u <- rnorm(nobs, 0, exp(0.8 * d2$w1))
d2$y <- d2$x + d2$z + d2$u

fit_bad <- gmm(y ~ x, x = ~ w1 + w2, data = d2)
coef(fit_bad)["x"]
```

```
       x
1.595747
```

```
specTest(fit_bad)
```

```
 ##  J-Test: degrees of freedom is 1  ##

                J-test      P-value
Test E(g)=0:    1.4345e+02  4.6853e-33
```

The coefficient moves to 1.60 and the J statistic goes to 143, with a p-value of 5×10^{-33}. This is what overidentification adds. With one instrument there is no way to notice that it is invalid, because a single moment condition can always be satisfied exactly. With two, the instruments have to agree about β, and when one of them is contaminated they do not. Two limits are worth keeping in view. The test does not say *which* instrument is at fault, and it has no power against violations the instruments share: if w_1 and w_2 were both correlated with z in the same way they would agree on the same wrong answer, and J would pass.

5.6 A few concepts of conditioning

5.6.1 Independence

If X and Y are independent then

$$f(x, y) = f(x)f(y) \tag{5.18}$$

and hence

$$f(y|x) = f(y). \tag{5.19}$$

If X and Y are independent then

$$E[g(X)h(Y)] = E[g(X)] \cdot E[h(Y)] \tag{5.20}$$

and hence

$$Cov[g(X), h(Y)] = 0. \tag{5.21}$$

i.e. all functions of X and Y are uncorrelated.

5.6.2 Law of Iterated Expectations

$$E[Y] = E[E(Y|X)]. \tag{5.22}$$

5.6.3 Dependence Concepts

X, Y independent:

$$Cov[g(X), h(Y)] = 0 \tag{5.23}$$

Y mean independent of X, $E[Y|X] = E[Y]$ (for a mean-zero disturbance u, $E[u|X] = 0$):

$$Cov[g(X), Y] = 0 \tag{5.24}$$

X, Y uncorrelated:

$$Cov[X, Y] = 0 \tag{5.25}$$

These are listed in decreasing strength, and the implications run one way only: independence ⇒ mean independence ⇒ uncorrelatedness. The middle condition is what a *model of the conditional mean* needs. Neither converse holds: X and Y can be uncorrelated with $E[Y \mid X] \neq 0$ (take $Y = X^2 - 1$ with $X \sim N(0,1)$, so $Cov[X,Y] = E[X^3] = 0$ while $E[Y \mid X] = X^2 - 1$), and mean independence permits dependence through higher moments (any pure heteroskedasticity, $Y = X\epsilon$ with ϵ independent of X and mean zero). Keeping the three apart matters because the moment conditions below need only the *weakest* of them. $E[u_i] = 0$ and $E[X_i u_i] = 0$ are statements about means and covariances, nothing more. Conditional mean zero is sufficient for them and is not necessary, which is exactly why GMM asks for less than a conditional-mean model does.

5.6.4 Regression

A regression model is a model of $E[Y_i|X_i]$. For example,

$$Y_i = \beta_0 + \beta_1 X_i + u_i \tag{5.26}$$

where $E[u_i|X_i] = 0$.

5.6.5 GMM regression

The regression model

$$Y_i = \beta_0 + \beta_1 X_i + u_i, \quad E[u_i|X_i] = 0 \tag{5.27}$$

implies the moment condition

$$E[u_i] = 0 \quad \text{and} \quad E[X_i u_i] = 0 \tag{5.28}$$

That is,

$$E[Y_i - \beta_0 - \beta_1 X_i] = 0 \tag{5.29}$$

$$E[X_i(Y_i - \beta_0 - \beta_1 X_i)] = 0 \tag{5.30}$$

The sample moment conditions are

$$\frac{1}{n}\sum_{i=1}^{n}(y_i - \hat{\beta}_0 - \hat{\beta}_1 x_i) = 0 \tag{5.31}$$

$$\frac{1}{n}\sum_{i=1}^{n} x_i(y_i - \hat{\beta}_0 - \hat{\beta}_1 x_i) = 0 \tag{5.32}$$

These are just normal equations for OLS.

A characteristic of GMM: the specification of the model generates the estimator. i.e. only $E[Y_i|X_i] = \beta_0 + \beta_1 X_i$ is assumed.

Note there are no assumptions that u_i is homoscedastic, not autocorrelated or normally distributed. These properties affect the statistical properties of the GMM estimator, not its definition.

5.7 Summary

The method of moments sets sample moments equal to their population counterparts. OLS solves the moment condition that the regressors are uncorrelated with the error, and IV solves the same condition with instruments. GMM generalizes this to more moment conditions than parameters, minimizing a weighted quadratic form in the sample moments. The efficient weight is the inverse of the moments' variance, estimated in two steps. 2SLS is the special case that is efficient only under homoskedasticity. GMM asks less than MLE. It needs correct moment conditions and makes no assumption about the full distribution.

In the just-identified example, GMM and 2SLS coincide and the weighting does not matter. With two instruments and heteroskedastic errors, two-step GMM gives 0.989 against a true 1, with a slightly smaller standard error than robust 2SLS. The J test then has content. It is 1.65 ($p = 0.20$) with valid instruments and 143 when one instrument is made invalid. It cannot say which instrument is at fault, and it has no power when all instruments share the same violation.

6 Censored, Truncated or Selected Data

6.1 Compare three different cases

A sample is truncated if some observations are systematically excluded from the sample. For example, suppose we are interested in relationship between people's income(y) and education(x). If we have observations of both y and x only for people whose income is above \$20,000 per year, then we have a truncated sample. A sample is censored if no observations have been systematically excluded but some of information contained in them has been suppressed. In other words, in a truncated sample, you don't have observations which have been truncated; while in a censored sample, you do have observations but you don't have full information. The idea of "censoring" is that some data above or below a threshold are mis-reported at the threshold. If we use the same example of people's income, if people with income lower than \$20,000 per year do not appear in the sample at all—we have no observations of either y or x for them—then we have a truncated sample. If we have observations of x for people whose income above AND below that limit, but no information on exactly how much their income is for those whose income below the limit (only thing we know is that it's lower than \$20,000 per year), then we have a censored sample.

Sample selection is a third case, and it is not censoring with an extra variable attached. In a censored sample the outcome is missing in a *known* way: the threshold is known, and the suppressed value is known to lie beyond it. That is information, and the Tobit model below uses it. In a selected sample the outcome is simply unobserved for some units, and what decides whether we observe it is a separate selection process rather than the value of y crossing a threshold. There need be no threshold in y at all.

	Truncation	Censoring	Sample selection
What we observe for a unit that is selected out	nothing; the unit is absent from the sample	x, and the fact that y lies beyond the threshold	x; y is missing
Is a threshold in y known?	yes	yes	not in general
What determines observation	the value of y	the value of y	a separate selection process, usually its own equation

The union example is sample selection. We observe education for everybody, members and non-members alike, and income only for members. Nothing tells us that a non-member's income sits above or below any particular figure; it is simply missing. Union membership is the selection equation, and it is what the estimation below has to model. So read sample selection as missing outcome data governed by a selection process.

6.2 An illustration for truncated or censored data

Consider the model

$$y_t^0 = \beta_1 + \beta_2 x_t + u_t,\ u_t \sim NID(0, \sigma^2), \tag{6.1}$$

where y_t^0 is a latent variable. We actually observe y_t, which differs from y_t^0 , because it's either truncated or censored.

Suppose that censorship or truncation occurs whenever y_t^0 is less than 0. Clearly, the larger the error term u_t, the larger is y_t^0 and thus the greater must be the probability that $y_t^0 \geq 0$. This probability also depends on x_t. So for the sample we observe, $E(y_t \mid x_t)$ is no longer $x_t\beta$: censoring or truncation changes the observed conditional mean, so the gap between y_t and $x_t\beta$ does not have conditional mean 0. The latent error u_t itself need not be endogenous. OLS of y_t on x_t in truncated or censored samples is therefore biased and inconsistent for the latent slope β. Selection on x_t alone would not do this. Normally we would like to draw inference on the population which is represented by the full sample. Shown in the figure below, ideally we would have the OLS regression line (if we had the full sample), which is very close to the "true" regression line (which is the mechanism generates the data). We would have the "OLS censored sample" line if we run OLS on the censored sample; we would have "OLS truncated sample" if we run OLS on the truncated sample. It shows that both of them are severely biased from the "true" regression line.

Here is a graph to illustrate from Davidson and MacKinnon (2004).




Figure 6.1: Effects of truncation and censoring from Davidson and MacKinnon (2004)

6.3 Truncated Models

For truncated data, a consistent estimator comes from maximum likelihood estimator (MLE). If we assume the error terms in the latent variable model has a known distribution, then MLE estimator

can be applied. The most popular choice is Gaussian.

$$\begin{aligned}\Pr(y_t^0 \geq 0) &= \Pr(\mathbf{X}_t\beta + u_t \geq 0) \\ &= 1 - \Pr(u_t/\sigma < -\mathbf{X}_t\beta/\sigma) \\ &= 1 - \Phi(-\mathbf{X}_t\beta/\sigma) = \Phi(\mathbf{X}_t\beta/\sigma)\end{aligned} \tag{6.2}$$

The density of y_t is proportional to the density of y_t^0 when $y_t^0 \geq 0$ and y_t is observed. It is 0 elsewhere. The factor of proportionality, which is needed to ensure that the density integrates to unity, is the inverse of the probability that $y_t^0 \geq 0$. The density of y_t can be written as

$$\frac{\sigma^{-1}\phi((y_t - \mathbf{X}_t\beta)/\sigma)}{\Phi(\mathbf{X}_t\beta/\sigma)} \tag{6.3}$$

This implies the log-likelihood function,

$$\ell(\mathbf{y}, \boldsymbol{\beta}, \sigma) = -\frac{n}{2}\log(2\pi) - n\log(\sigma) - \frac{1}{2\sigma^2}\sum_{t=1}^{n}(y_t - \mathbf{X}_t\beta)^2 - \sum_{t=1}^{n}\log\Phi(\mathbf{X}_t\beta/\sigma). \tag{6.4}$$

This can be estimated by MLE. In Stata, truncreg is the command to do truncated regression model.

6.4 Censored Models

The most popular censored model is Tobit model.

$$\begin{aligned}&y_t^0 = \mathbf{X}_t\beta + u_t, \quad u_t \sim NID(0, \sigma^2) \\ &y_t = y_t^0 \text{ if } y_t^0 > 0;\ y_t = 0 \text{ otherwise.}\end{aligned} \tag{6.5}$$

We see that

$$\begin{aligned}\Pr(y_t = 0) = \Pr(y_t^0 \leq 0) &= \Pr(\mathbf{X}_t\beta + u_t \leq 0) \\ &= \Pr(u_t/\sigma < -\mathbf{X}_t\beta/\sigma) \\ &= \Phi(-\mathbf{X}_t\beta/\sigma)\end{aligned} \tag{6.6}$$

The contribution to the log-likelihood function made by observations with $y_t = 0$ is

$$\ell_t(y_t, \beta, \sigma) = \log\Phi(-\mathbf{X}_t\beta/\sigma). \tag{6.7}$$

If y_t is positive, the contribution to the log-likelihood is the logarithm of the density,

$$\log(\frac{1}{\sigma}\phi((y_t - \mathbf{X}_t\beta)/\sigma)). \tag{6.8}$$

The log-likelihood function of the tobit model is

$$\sum_{y_t=0}\log\Phi(-\mathbf{X}_t\beta/\sigma) + \sum_{y_t>0}\log(\frac{1}{\sigma}\phi((y_t - \mathbf{X}_t\beta)/\sigma)) \tag{6.9}$$

This can be estimated by MLE. In Stata, tobit or intreg can be used for censoring models.

Both this likelihood and the truncated one in Equation 6.4 lean on the normal, homoskedastic latent error far more than a linear regression does. The slope β and the scale σ enter together through $\Phi(\mathbf{X}_t\beta/\sigma)$ and the Mills-ratio terms, so if u_t is heteroskedastic or not normal, the MLE of β is inconsistent. This is not like linear OLS or Poisson QMLE, where a correct conditional mean is enough and a sandwich variance repairs the standard errors: `vce(robust)` after `tobit` or `truncreg` fixes nothing about the point estimates.

6.5 Sample Selection

The sample selection models differ from censored model in that it involves a different variable (from y itself) to determine the censorship (selection).

Suppose that y_t^0 and z_t^0 are two latent variables, generated by the bivariate process

$$\begin{bmatrix} y_t^0 \\ z_t^0 \end{bmatrix} = \begin{bmatrix} \mathbf{X}_t\beta \\ \mathbf{W}_t\gamma \end{bmatrix} + \begin{bmatrix} u_t \\ v_t \end{bmatrix}, \ \begin{bmatrix} u_t \\ v_t \end{bmatrix} \sim NID(\mathbf{0}, \begin{bmatrix} \sigma^2 & \rho\sigma \\ \rho\sigma & 1 \end{bmatrix}) \tag{6.10}$$

Here ρ is the correlation between u_t and v_t; since $\text{Var}(u_t) = \sigma^2$ and $\text{Var}(v_t) = 1$, their covariance is $\rho\sigma$.

We observe y_t and z_t:

$$y_t = y_t^0 \text{ if } z_t^0 > 0;\ y_t \text{ unobservable otherwise; } z_t = 1 \text{ if } z_t^0 > 0;\ z_t = 0 \text{ otherwise.} \tag{6.11}$$

There are two types of observations, ones we observe $y_t = y_t^0$ and $z_t = 1$, along with both $\mathbf{X_t}$ and $\mathbf{W_t}$, and ones we observe only $z_t = 0$ and $\mathbf{W_t}$.

Each observation contributes to the likelihood function by

$$I(z_t = 0)\Pr(z_t = 0) + I(z_t = 1)\Pr(z_t = 1)f(y_t^0|z_t = 1), \tag{6.12}$$

Under the normality assumption,

$$u_t = \rho\sigma v_t + e_t \tag{6.13}$$

where e_t is independent of $v_t \sim N(0, 1)$, with $\text{Var}(e_t) = \sigma^2(1-\rho^2)$. A useful fact about the standard normal distribution is that

$$\text{E}(v_t|v_t > -x) = \lambda(x) = \frac{\phi(x)}{\Phi(x)} \tag{6.14}$$

and the function $\lambda(x)$ is called the inverse Mills ratio.

The log-likelihood function can be shown to be

$$\sum_{z_t=0} \log\Phi(-\mathbf{W}_t\gamma) + \sum_{z_t=1} \log(\frac{1}{\sigma}\phi((y_t - \mathbf{X}_t\beta)/\sigma)) + \sum_{z_t=1} \log\Phi(\frac{\mathbf{W}_t\gamma + \rho(y_t - \mathbf{X}_t\beta)/\sigma}{(1-\rho^2)^{1/2}}) \tag{6.15}$$

So this model can be estimated by MLE.

It is also popular to use Heckman's (1979) two-step method.

Heckman's method is based on the fact that the original latent model can be rewritten as

$$y_t = \mathbf{X}_t\beta + \rho\sigma v_t + e_t \tag{6.16}$$

Here the error term u_t is divided into two parts, one perfectly correlated with v_t, and one independent of v_t.

In the first step, an ordinary probit model is used to obtain consistent estimates $\hat{\gamma}$ of the parameters of the selection equation.

In the second step, the unobserved v_t is replaced by the selectivity regressor $\frac{\phi(\mathbf{W}_t\hat{\gamma})}{\Phi(\mathbf{W}_t\hat{\gamma})}$.

Therefore, the regression becomes

$$y_t = \mathbf{X}_t\beta + \rho\sigma\frac{\phi(\mathbf{W}_t\hat{\gamma})}{\Phi(\mathbf{W}_t\hat{\gamma})} + e_t \tag{6.17}$$

Or,

$$y_t = \mathbf{X}_t\beta + \rho\sigma\hat{\lambda}_t + e_t \tag{6.18}$$

Strictly, Equation 6.18 hides two parts of the error. Write $\lambda_t = \phi(\mathbf{W}_t\gamma)/\Phi(\mathbf{W}_t\gamma)$ for the inverse Mills ratio at the true γ. Conditional on selection, the second-step error is

$$u_t^* = e_t + \rho\sigma(v_t - \lambda_t) + \rho\sigma(\lambda_t - \hat{\lambda}_t). \tag{6.19}$$

The middle term makes the error heteroskedastic even with γ known. Among selected units, $\mathrm{Var}(v_t \mid z_t = 1, \mathbf{W}_t) = 1 - \lambda_t(\lambda_t + \mathbf{W}_t\gamma)$, so the error variance is $\sigma^2[1 - \rho^2\lambda_t(\lambda_t + \mathbf{W}_t\gamma)]$, which changes from unit to unit. The last term comes from estimating γ: $\hat{\lambda}_t$ is a generated regressor, as in the control-function discussion of the Endogeneity chapter. A robust (sandwich) variance in the second-step OLS handles the first problem but not the second. Heckman's two-step covariance matrix, or a bootstrap over both steps, handles both. Neither term biases the coefficients: $\rho\sigma$ on $\hat{\lambda}_t$ and β remain consistent.

One thing to note in Heckman model is that in many situations, $\hat{\lambda}$ is believed to be highly collinear with $\mathbf{X_t}$, if $\mathbf{W_t}$ is the same as $\mathbf{X_t}$. However, the model can still be estimated due to the nonlinearity of the model. Nevertheless, it becomes a regular practice to require $\mathbf{W_t}$ contains at least one extra variable than $\mathbf{X_t}$, which is sometimes called exclusion restriction. In many situations, $\mathbf{W_t}$ contains all $\mathbf{X_t}$ variables, and at least one more variable. The reason to include all the $\mathbf{X_t}$ variables is more mundane than it is often made to sound: whatever plausibly affects selection belongs in the selection equation, and the determinants of the outcome usually do. What identification turns on is the other direction – at least one variable in $\mathbf{W_t}$ kept *out* of $\mathbf{X_t}$, affecting selection but not the outcome. Without such an exclusion, $\hat{\lambda}$ is a nonlinear function of the same $\mathbf{X_t}$'s and identification rests entirely on that nonlinearity.

In Stata, heckman is the command to do sample selection models. It has options to do a Heckman two-step estimation or MLE estimation. MLE is usually preferred, on the understanding that its efficiency advantage holds *if* the joint normality it assumes is right. The two-step estimator is less efficient but leans less heavily on the full likelihood, which makes it easier to diagnose when the specification is in doubt.

6.6 Switching Regression (Treatment-Effects Model)

There is another situation that is similar to sample selection model: we observe y for $z = 1$ and $z = 0$. In the example of effect of education on income, we observe both union member's income and non-member's income. In this case, we have switching regression model or treatment-effect model. The treatment effect model estimates the effect of an endogenous binary treatment z_t (treatment, program participation, etc.) on a continuous, fully-observed variable, y_t, conditional on the independent variables x_t and w_t.

Suppose z_t^0 is a latent variable.

$$\begin{bmatrix} y_t \\ z_t^0 \end{bmatrix} = \begin{bmatrix} \mathbf{X}_t\beta \\ \mathbf{W}_t\gamma \end{bmatrix} + \delta \begin{bmatrix} z_t \\ 0 \end{bmatrix} + \begin{bmatrix} u_t \\ v_t \end{bmatrix}, \ \begin{bmatrix} u_t \\ v_t \end{bmatrix} \sim NID(\mathbf{0}, \begin{bmatrix} \sigma^2 & \rho\sigma \\ \rho\sigma & 1 \end{bmatrix}) \tag{6.20}$$

We observe y_t and z_t:

$$z_t = 1 \text{ if } z_t^0 > 0; \ z_t = 0 \text{ otherwise.} \tag{6.21}$$

Notice that the only difference between this model and the selection model is that we observe y_t when $z_t = 0$ and $z_t = 1$. It's not a selection, but a regime switching.

One caveat about the name. The specification above lets treatment shift only the *intercept*: there is a single β and a single δ, so the treatment effect is δ for everyone. That is the treatment-effects model. A **switching regression** in the fuller sense lets the whole coefficient vector differ by regime,

$$y_t = z_t(\mathbf{X}_t\beta_1 + u_{1t}) + (1 - z_t)(\mathbf{X}_t\beta_0 + u_{0t}), \qquad z_t^0 = \mathbf{W}_t\gamma + v_t, \tag{6.22}$$

with u_{1t} and u_{0t} each allowed to correlate with v_t (two selection parameters, ρ_1 and ρ_0, rather than one). This is the Roy model, and the treatment effect $\mathbf{X}_t(\beta_1 - \beta_0)$ then varies with $\mathbf{X_t}$ instead of being a constant. The distinction matters in practice: if the returns to education genuinely differ between union and non-union jobs, the common-slope model above will report a single δ that is some average of them and will not reveal the heterogeneity. Fitting it requires two outcome equations rather than one intercept shift.

A word on software, since the two models above are easy to conflate. Stata's etregress fits the *common-slope* specification – the endogenous treatment-effects model with a single δ – not the two-regime switching regression with separate β_1 and β_0. For the fuller Roy-type model, Stata users generally reach for the user-written movestay (endogenous switching regression), and SAS's qlim covers a range of limited-dependent and selection specifications. The default in each case is MLE.

6.7 Summary

Truncation drops units outside a threshold. Censoring keeps them but reports only that the outcome lies beyond a known threshold. Sample selection leaves the outcome missing for units chosen by a separate process. In all three, OLS on the observed data is biased and inconsistent, because the error no longer has mean zero given the regressors. Truncated regression and Tobit fix the first two by maximum likelihood under normal errors.

For selection, the Heckman model adds a selection equation whose error is correlated with the outcome error. It can be estimated by MLE, or in two steps with a probit followed by OLS that includes the inverse Mills ratio. The two-step version is less efficient but relies less on the full likelihood. It is fragile when the selection equation has no variable excluded from the outcome equation. When the outcome is observed in both regimes, the problem becomes an endogenous treatment-effects model. Its fuller form, the Roy switching regression, lets every coefficient differ by regime.

7 Discrete and Limited Dependent Variables

7.1 Binary Response Models

7.1.1 Probit and Logit

Let P_t denote the probability that $y_t = 1$ conditional on the information set Ω, which consists of exogenous and predetermined variables. A binary response model serves to model this conditional expectation. Since the values are 0 or 1, it is clear that P_t is also the expectation of y_t conditional on Ω_t:

$$P_t \equiv \Pr(y_t = 1|\Omega_t) = \mathrm{E}(y_t|\Omega_t). \tag{7.1}$$

Here X_t is a set of regressors. A linear probability model would set $P_t = X_t\beta$ directly, which requires the fitted values to stay in $[0, 1]$ – an awkward constraint, since the linear index $X_t\beta$ is unbounded. Logit and probit avoid this problem by modeling $P_t = F(X_t\beta)$, mapping the unbounded index through a CDF-valued transformation into $[0, 1]$.

We ensure that $0 \leq P_t \leq 1$ by specifying that

$$P_t \equiv \Pr(y_t = 1|\Omega_t) = F(\mathbf{X_t}\boldsymbol{\beta}). \tag{7.2}$$

$F(x)$ is a transformation function, which has the same characteristics as the CDF of a probability distribution.

Two popular choices of $F(x)$ are Gaussian (probit) and Logistic (logit).

The less familiar logistic function is

$$\Lambda(x) = \frac{e^x}{1 + e^x} \tag{7.3}$$

The logit model is most easily derived by assuming that

$$\log(\frac{P_t}{1 - P_t}) = \mathbf{X_t}\boldsymbol{\beta} \tag{7.4}$$

which says the logarithm of the odds (the ratio of the two probabilities) is equal to $\mathbf{X_t}\boldsymbol{\beta}$. Therefore,

$$P_t = \frac{\exp(\mathbf{X}_t\beta)}{1 + \exp(\mathbf{X}_t\beta)} = \Lambda(\mathbf{X}_t\beta) \tag{7.5}$$

7.1.2 MLE for binary data

The likelihood for an observation t is the probability that $y_t = 1$ if $y_t = 1$, or the probability that $y_t = 0$ if $y_t = 0$. The logarithm of the appropriate probability is then the contribution to the loglikelihood made by observation t. Therefore, if $\mathbf{y}$ is an n-vector with typical element y_t, the loglikelihood function for $\mathbf{y}$ can be written as

$$\ell(\mathbf{y}, \boldsymbol{\beta}) = \sum_{t=1}^{n} (y_t \log F(\mathbf{X_t}\boldsymbol{\beta}) + (1 - y_t) \log(1 - F(\mathbf{X_t}\boldsymbol{\beta}))) \tag{7.6}$$

For the logit and probit models, this function is globally concave with respect to β, so there are no local maxima to worry about. Concavity is not the whole story, though. For the MLE to exist and be unique and *finite* we also need $\mathbf{X}$ to have full column rank, and the data must not be separable. If some linear combination of the regressors predicts the outcome perfectly – complete or quasi-complete separation – the likelihood keeps rising along that direction and the maximum is attained at no finite β, concave or not. Software reports this as coefficients diverging or standard errors blowing up. With full rank and no separation, the first-order conditions, or likelihood equations, uniquely define the MLE estimator $\hat{\beta}$. These likelihood equations can be written as

$$\sum_{t=1}^{n} \frac{(y_t - F(\mathbf{X_t}\boldsymbol{\beta}))f(\mathbf{X_t}\boldsymbol{\beta})x_{ti}}{F(\mathbf{X_t}\boldsymbol{\beta})(1 - F(\mathbf{X_t}\boldsymbol{\beta}))} = 0, \ i = 1, \ldots, k. \tag{7.7}$$

where $f = F'$ is the density (e.g. ϕ for probit). For logit, $f = F(1 - F)$, so the numerator cancels the denominator and the score reduces to $\sum_t (y_t - F)x_{ti} = 0$.

Newton's Method can be used to find $\hat{\beta}$.

7.2 Models for More than Two Discrete Responses

7.2.1 The Ordered Probit

Ordered Probit can be easily derived from a latent variable model.

$$y_t^0 = \mathbf{X}_t\beta + u_t, \ u_t \sim NID(0, 1) \tag{7.8}$$

Suppose we observe y_t with three values.

$$\begin{cases} y_t = 0 & \text{if } y_t^0 < \gamma_1; \\ y_t = 1 & \text{if } \gamma_1 \leq y_t^0 < \gamma_2; \\ y_t = 2 & \text{if } y_t^0 \geq \gamma_2. \end{cases} \tag{7.9}$$

Therefore,

$$\begin{aligned} \Pr(y_t = 0) &= \Pr(y_t^0 < \gamma_1) = \Pr(\mathbf{X}_t\beta + u_t < \gamma_1) \\ &= \Pr(u_t < \gamma_1 - \mathbf{X}_t\beta) = \Phi(\gamma_1 - \mathbf{X}_t\beta) \end{aligned} \tag{7.10}$$

Similarly,

$$\Pr(y_t = 2) = 1 - \Phi(\gamma_2 - \mathbf{X}_t\beta) \tag{7.11}$$

$$\Pr(y_t = 1) = \Phi(\gamma_2 - \mathbf{X}_t\beta) - \Phi(\gamma_1 - \mathbf{X}_t\beta) \tag{7.12}$$

These probabilities depend solely on the value of the index function and on the two threshold parameters.

Two normalizations are needed before any of this can be estimated, and they are easy to pass over. First, the *location*: an intercept in $\mathbf{X}_t\beta$ and both thresholds cannot be identified together, since adding a constant to the index and the same constant to γ_1 and γ_2 leaves every probability unchanged. Software resolves this by dropping the intercept or by fixing $\gamma_1 = 0$. Second, the *scale*: that is what setting $\mathrm{Var}(u_t) = 1$ above does, since multiplying the index, the thresholds and u_t by any positive constant is likewise unobservable. The multinomial logit below needs a location normalization of its own, $\beta^0 = 0$: adding the same vector to every alternative's coefficients leaves all choice probabilities unchanged. Its scale is fixed by the extreme-value error distribution, much as $\mathrm{Var}(u_t) = 1$ fixes it here.

The loglikelihood function is

$$\ell(\beta, \gamma_1, \gamma_2) = \sum_{y_t=0} \log(\Phi(\gamma_1 - \mathbf{X}_t\beta)) + \sum_{y_t=2} \log(\Phi(\mathbf{X}_t\beta - \gamma_2)) + \sum_{y_t=1} \log(\Phi(\gamma_2 - \mathbf{X}_t\beta) - \Phi(\gamma_1 - \mathbf{X}_t\beta)) \tag{7.13}$$

7.2.2 The Multinomial Logit

When responses are unordered, two popular choices are multinomial logit and conditional logit. They are easy to conflate because they share the same softmax form, but they differ in what varies across the $J + 1$ alternatives $l = 0, \dots, J$: in multinomial logit the *regressors* are individual-specific and only the *coefficients* vary by alternative; in conditional logit it's the reverse.

Multinomial logit. Here the regressor vector $\mathbf{X_t}$ (individual-specific characteristics, e.g. age, income) is the same for every alternative, and it is the coefficient vector that is alternative-specific:

$$\Pr(y_t = l) = \frac{\exp(\mathbf{X}_t\beta^l)}{\sum_{j=0}^{J} \exp(\mathbf{X}_t\beta^j)} \quad \text{for } l = 0, \dots, J, \tag{7.14}$$

Here $\mathbf{X_t}$ is a row vector of length k of individual-specific covariates, and β^l is a k-vector of alternative-specific parameters. For identification one category is taken as the base with $\beta^0 = 0$ (so its numerator is 1).

Conditional logit. Here it is the regressors that vary by alternative (e.g. the price or characteristics of choice j itself), while the coefficient vector is common across alternatives:

$$\Pr(y_t = l) = \frac{\exp(\mathbf{W}_{tl}\beta)}{\sum_{j=0}^{J} \exp(\mathbf{W}_{tj}\beta)} \quad \text{for } l = 0, \dots, J, \tag{7.15}$$

Here $\mathbf{W}_{tj}$ is a row vector of length k of alternative-j-specific variables, and β is a single k-vector of parameters shared across all alternatives $j = 0, \dots, J$.

For binary and multinomial models alike, the coefficients β do not directly measure the effect of a unit change in X on the probability. That effect depends on where in the distribution the

change occurs. The average marginal effect (AME) — averaging the partial effect over the observed distribution of covariates — is the standard summary. The full treatment of AME vs. marginal effects at the mean, including fixed-effects models, is in the Blog Book.

Writing *both* choice-varying regressors $\mathbf{W}_{tl}$ *and* choice-specific coefficients β^l into the same formula, as some treatments do, conflates the two models. It is worth being precise about what that costs, because the usual complaint – that such a specification is not identified – is not true. Choice probabilities depend only on utility *differences*, so $\Pr(y_t = l)$ is governed by $\mathbf{W}_{tj}\beta^j - \mathbf{W}_{tl}\beta^l$; when the $\mathbf{W}_{tj}$ vary independently across alternatives, those differences pin down each β^j separately, with no normalization required. It is a perfectly estimable model – it is what Stata's `asclogit` fits when alternative-specific coefficients are requested.

What the ambiguous formula really costs is clarity about *which normalization is in force*. Individual-specific regressors require one alternative's coefficients to be fixed ($\beta^0 = 0$), since only differences across alternatives are identified; alternative-specific regressors entering with a common coefficient require no such restriction. One caveat on "no normalization required" above: it concerns the slope coefficients β^j. If the specification also carries alternative-specific intercepts, those are identified only up to a location shift and one of them still has to be fixed. A formula that does not say which of the two it means leaves the reader unable to tell what has been normalized, and therefore unable to interpret the coefficients.

7.2.3 Independence of Irrelevant Alternatives

Both logit forms carry a restriction that comes from the softmax itself. Take the ratio of two choice probabilities. With individual-specific regressors,

$$\frac{\Pr(y_t = j)}{\Pr(y_t = l)} = \exp(\mathbf{X}_t(\beta^j - \beta^l)), \tag{7.16}$$

and with alternative-specific regressors and a common coefficient the ratio is $\exp((\mathbf{W}_{tj} - \mathbf{W}_{tl})\beta)$. Either way everything to do with the other alternatives cancels. The relative odds of j against l do not depend on what else is on the menu, or on the attributes of any third alternative. That is **independence of irrelevant alternatives**, or IIA.

Sometimes it is harmless. Often it is not. The standard illustration is travel mode. Suppose commuters split evenly between a car and a red bus, so the odds are 1:1 and each share is 1/2. Now add a blue bus, identical to the red one in everything but colour. IIA holds the car-to-red-bus odds at 1:1, and the two buses are interchangeable, so the model predicts three shares of 1/3. What we would expect is that the blue bus draws almost entirely from the red bus, leaving the car near 1/2. The model has no way to know that the two buses are close substitutes, because the softmax gives it no way to express that.

There are two standard escapes. **Nested logit** groups alternatives into nests – the two buses in one, the car in another – and allows correlation within a nest, so IIA holds inside nests but not across them. **Mixed logit**, also called random-coefficients logit, lets β vary across individuals; shared taste parameters then induce correlation across alternatives and IIA goes away, at the cost of simulation-based estimation. Which one to reach for depends on whether the substitution pattern you are worried about can be described by a grouping.

7.3 Summary

A binary outcome's conditional mean is a probability, so probit and logit pass a linear index through a CDF to keep it between 0 and 1. Both are fitted by maximum likelihood with a globally concave log-likelihood. The estimate exists and is finite only with full-rank regressors and no separation. Ordered probit adds thresholds on a latent index, with location and scale fixed by normalization. Multinomial logit lets coefficients vary by alternative for individual characteristics. Conditional logit uses one coefficient vector for characteristics that vary by alternative. In every case the coefficients are not effects on probabilities, and the average marginal effect is the usual summary.

Both logit forms impose independence of irrelevant alternatives: the odds between two choices do not depend on the rest of the menu. The red-bus/blue-bus example shows why that can fail. Nested logit relaxes it across groups of close substitutes, and mixed logit removes it through random coefficients.

8 Count Data Models

8.1 Poisson Model

If the interested variable is a count, it is natural to model its conditional distribution as Poisson. What the regression assumes is the *distribution*, not that the counts were generated by a continuous-time Poisson process – that process is one way to arrive at the distribution, not a requirement of the model. The number of occurrences follows the Poisson distribution:

$$\Pr[Y = y] = \frac{e^{-\mu}\mu^{y}}{y!} \tag{8.1}$$

One of Poisson's properties is that it has equal variance as its mean:

$$\mathrm{E}[Y] = \mathrm{Var}[Y] = \mu \tag{8.2}$$

In a count data model such as Poisson regression model, we are modeling log of the expected counts:

$$\log(\mathrm{E}(Y)) = \mathbf{X}_{\mathbf{i}}'\boldsymbol{\beta} \tag{8.3}$$

The log-likelihood function is

$$\log L(\beta) = \sum_{i=1}^{N} -\exp(X_i'\beta) + y_i X_i'\beta - \ln y_i! \tag{8.4}$$

To maximize it, the first order condition is

$$\sum_{i=1}^{N} (y_i - \exp(X_i'\beta))X_i' = 0 \tag{8.5}$$

We can use Newton-Raphson or other optimization algorithms to find the solution for the first order condition.

8.2 QMLE Poisson

8.2.1 Model

Note that in the first-order condition above, if X_i' includes a constant, then $(y_i - \exp(X_i'\beta))$ sums to zero. If $\mathrm{E}[y_i|X_i] = \exp(X_i'\beta)$, then the summation on the left-hand side has expectation of zero. Hence the only specification needed to apply it is the conditional expectation of Y given X. Even if the data is not Poisson-distributed, the estimator is still consistent. Therefore, this estimator is called the quasi-ML (QML) Poisson estimator.

Although the QML Poisson estimator is consistent under relatively weak condition, the regular variance estimator for the coefficients are not valid anymore.

If a stronger assumption is made that the data follows a Poisson distribution, then the error term has a variance which equals to the mean of Y. The estimator $\hat{\beta}_P$ (ML estimator) follows a normal distribution asymptotically with a variance matrix

$$\mathrm{Var}_{ML}[\hat{\beta}_P] = (\sum_{i=1}^{N} \mu_i X_i X_i')^{-1} \tag{8.6}$$

On the other hand, the variance matrix for the QML Poisson estimator $\hat{\beta}_{QML}$ is

$$\mathrm{Var}_{QML}[\hat{\beta}_{QML}] = (\sum_{i=1}^{N} \mu_i X_i X_i')^{-1} (\sum_{i=1}^{N} \omega_i X_i X_i') (\sum_{i=1}^{N} \mu_i X_i X_i')^{-1} \tag{8.7}$$

where $\omega_i = \mathrm{Var}[y_i|X_i]$ is the conditional variance of y_i.

8.2.2 Implementation

Poisson estimation is implemented in almost every statistical package. However, some of them may not work if you have a continuous dependent variable.

Stata's implementation of the Poisson model: "poisson" and "xtpoisson" do take a continuous dependent variable. Standard errors reported without any "vce" specification assume the Poisson variance equals the mean and are likely too small under overdispersion. For a cross-sectional Poisson treated as QMLE, the robust (sandwich) standard errors – the "robust" option – *are* exactly the adjustment needed for a misspecified variance, so they are valid even when the equidispersion assumption fails; the conditional mean only needs to be correctly specified. In the panel fixed-effects case, serial correlation within a unit is a separate issue that requires *clustered* standard errors. Older versions of "xtpoisson" did not offer a cluster-robust option, which is why the user-written program xtpqml was written to supply the cluster-robust standard errors of Wooldridge (1999) after xtpoisson.

8.3 Fixed-effect Poisson model

We specify a panel data count model as:

$$E[y_{it}|\alpha_i, X_{it}] = \alpha_i \exp(X'_{it}\beta) = \exp(\gamma_i + X'_{it}\beta). \tag{8.8}$$

It turns out that for fixed-effect Poisson model, just like in linear regression case, there is NO incidental parameter problem. Poisson MLE is the same for conditional and unconditional likelihood (see Cameron and Trivedi 2005, 805).

What the result does not do is relax the exogeneity requirement. Consistency of $\widehat{\beta}$ needs $E[y_{it} \mid \alpha_i, X_{i1}, \ldots, X_{iT}] = \alpha_i \exp(X'_{it}\beta)$ – strict exogeneity conditional on the unit effect, which rules out feedback from past outcomes to later regressors. The absence of an incidental-parameter problem is a statement about estimating β alongside the α_i, not a licence to ignore what the conditional mean assumes.

Therefore, the coefficient estimates are the same for conditional or unconditional fixed-effect Poisson model. That is, in Stata, "xtpoisson, fe" will return the same results as "xi: poisson i.group", is the same as "xtpqml, fe". The only difference is that "xtpqml, fe" returns "correct" standard errors for QMLE Poisson. You can also calculate standard errors with the other commands, using clustered standard errors, or bootstrapped standard errors.

A conditional fixed-effects negative binomial model also exists, from J. Hausman, Hall, and Griliches (1984) and implemented as `xtnbreg, fe` in Stata. It is not the negative binomial analogue of fixed-effects Poisson. It puts the unit effect in the dispersion parameter, not in the conditional mean, so it does not condition out an arbitrary multiplicative effect on the mean, and time-invariant regressors stay identified, which a true fixed-effects model would not allow (Allison and Waterman 2002; Guimarães 2008). Fixed-effects Poisson needs only the conditional mean to be right, since it is a QMLE, which is why it is preferred.

8.4 Negative Binomial model

The Poisson model has a restrictive property that the conditional variance equals the conditional mean. Often we'll see "overdispersion", that is, the variance exceeds the mean, in real data. The negative binomial model is a generalization of poisson model which allows for overdispersion by introducing an unobserved heterogeneity term. In a negative binomial model,

$$\mathrm{E}[Y|X, \tau] = \mu\tau = e^{X'\beta} \cdot \tau \tag{8.9}$$

This extra τ term follows a $Gamma(\theta, \theta)$ distribution, with $E(\tau) = 1$ and $Var(\tau) = 1/\theta$.

Conditional on X and τ, the distribution of Y is still poisson:

$$\Pr[Y = y|X, \tau] = \frac{e^{-\mu\tau}(\mu\tau)^y}{y!} \tag{8.10}$$

Conditional on X only, the distribution of Y is negative binomial:

$$\Pr[Y = y|X] = \frac{\theta^\theta \mu^y \Gamma(\theta + y)}{\Gamma(y + 1)\Gamma(\theta)(\mu + \theta)^{\theta+y}} \tag{8.11}$$

This distribution still has conditional mean of μ, but variance is $\mu(1 + (1/\theta)\mu)$.

When $\alpha = 1/\theta$ approaches zero, the negative binomial model converges to poisson model.

8.5 Models for truncated counts

In some situations, all we observe are non-zero counts, because of the way data was collected. For example, we only observe people's visits to a park, only if they visit. To model how many times they visit, based on this kind of data set, we need to use zero-truncated count models.

$$\Pr[Y = y|y > 0, X] = \frac{\Pr(y|X)}{\Pr(y > 0|X)} = \frac{\Pr(y|X)}{1 - \exp(-\mu)} \tag{8.12}$$

where the last equality holds for the Poisson case, since $\Pr(y = 0|X) = e^{-\mu}$; for the negative binomial, $\Pr(y = 0|X) = (\theta/(\theta + \mu))^\theta$, so the truncation factor is $1 - (\theta/(\theta + \mu))^\theta$ instead. Thus we are modeling the counts given that we have a positive outcome. Both zero-truncated poisson ("ztp" in Stata) and zero-truncated negative binomial ("ztnb" in Stata) are estimated by Maximum-likelihood method.

We need to pay extra attention to the "over-dispersion" problem when dealing with zero-truncated data. When data is not truncated, coefficients estimated by poisson model is consistent, even if data is "overdispersed". However, if we have zero-truncated sample, then "over-dispersion" results in inconsistent coefficient estimation (Grogger and Carson 1991). Therefore, "over-dispersion" needs to be tested before using a truncated *Poisson* model: under truncation the mean-variance restriction stops being innocuous, and violating it breaks consistency of the coefficients rather than only the standard errors.

Overdispersion is not an argument against the truncated negative binomial, though – that model exists precisely to accommodate it. What it does require is that the *form* of the overdispersion match the model. NB2 comes from a particular gamma-mixed Poisson, so overdispersion arising some other way leaves the truncated negative binomial misspecified in its turn. A test tells you that equidispersion fails; it does not tell you that the NB2 mixing story is the right replacement.

8.6 The hurdle regression model

Although negative binomial model relaxes the assumption of equal mean and variance, some researchers prefer modeling the process of generating zeros differently from the process of generating other values. Suppose zero counts are generated by a binary process:

$$\Pr[y = 0|X] = \frac{\exp(X\gamma)}{1 + \exp(X\gamma)} = \pi \tag{8.13}$$

Positive counts are generated by a truncated count process.

In this model, zero is a "hurdle" to get past before reaching positive counts. The hurdle model is estimated by two separate equations. It is easy to estimate. However, the marginal effect is tricky to calculate, since an independent variable, if appearing in both equations, will have effect through both equations.

$$\mathrm{E}(y|X) = [\pi \times 0] + (1-\pi) \times \mathrm{E}(y|y>0, X) = (1-\pi) \times \mathrm{E}(y|y>0, X) \tag{8.14}$$

In stata, there are commands such as "hplogit", which is a hurdle model with first equation logit, and second equation truncated poisson. It is the same as two separate estimations; namely "logit" first, then "ztp" on positive counts. Mind the direction of the first step. Equation 8.13 models the probability of a zero, π. A `logit` on the indicator $1(y>0)$ models the probability of crossing the hurdle, $1-\pi$, and its coefficients are $-\gamma$. Before plugging fitted probabilities into Equation 8.14, check which outcome the first step models.

The difference between a hurdle model and a Heckman selection model is that in a Heckman model the two equations are jointly estimated and the outcome equation must account for selection bias. An exclusion restriction – a variable that affects selection but not the outcome – is strongly recommended for the selection equation. Strictly speaking, under joint normality the model is identified by the nonlinearity of the inverse Mills ratio even without an excluded variable, but identification then rests entirely on functional form and is fragile, so an exclusion restriction is recommended in practice. On the other hand, a hurdle model assumes the two processes are separate; there is no selection.

8.7 Zero-inflated count models

A zero-inflated count model also models two different processes. One is a binary process: generating zero or non-zeros. The second one is a count model. Note there is a different between zero-inflated and hurdle model: In hurdle model, the second process is a truncated count model (positive counts). In a zero-inflated model, a zero can come from either the binary process, or from the count process.

A second difference is that the estimation of zero-inflated model is a joint estimation of the two processes.

$$y_i \sim \begin{cases} 0 \text{ with probability } \psi \\ g(y_i|X_i) \text{with probability } 1-\psi \end{cases} \tag{8.15}$$

Then

$$P(Y_i = y_i|X_i, Z_i) = \begin{cases} \psi(\gamma' Z_i) + (1-\psi(\gamma' Z_i))g(0|X_i) \text{ if } y_i = 0 \\ (1-\psi(\gamma' Z_i))g(y_i|X_i) \text{ if } y_i > 0 \end{cases} \tag{8.16}$$

This says that every non-zero observation follows a count process $g(y_i|X_i)$. Every zero observation has two possible sources: from a binary process (with probability $\psi(\gamma' Z_i)$) and from a count process (with probability $1-\psi(\gamma' Z_i)$ into the count process and then with probability of $g(0|X_i)$ to be a zero). The count process can be a Poisson process ("zip" in Stata) or a Negative Binomial process ("zinb" in Stata).

8.8 An example of extra-zero count data model comparison

A companion simulation study (see the detailed write-up in this guide's companion `blog_book`, chapter "Which count data model to use") compares plain Poisson, Negative Binomial, zero-inflated Poisson (ZIP, both with a correctly and an incorrectly specified zero-process), and hurdle models across sample sizes and zero-inflation levels. The main findings: with large samples, plain Poisson QMLE is consistent even with unmodeled excess zeros, provided the *observed* conditional mean is correctly specified. That proviso is worth spelling out. Under zero inflation the observed mean is $E[Y \mid X] = (1 - \pi(X))\,\mu(X)$, so fitting $\exp(\mathbf{X}\beta)$ recovers the slopes of μ only when π does not depend on X: constant inflation shifts the intercept by $\log(1-\pi)$ and leaves the slopes alone. If π varies with X, the term $\log(1-\pi(X))$ enters the mean function and the slopes are contaminated – independence of the latent zero and count *draws* does not rescue them. Where the inflation is constant, the excess zeros only inflate the standard errors (via overdispersion), not the point estimates, so robust/sandwich standard errors should be used rather than switching models. Zero-inflated Poisson with a correctly specified zero process performs best when that specification is available; hurdle and misspecified-zero-process ZIP perform similarly to plain Poisson; log-linear regression on $\log(y+1)$ is consistently the worst performer, since it targets a different (log-scale) conditional expectation than the models above.

8.9 Interpreting Coefficients and Marginal Effects in Nonlinear Models

This section is broader than count data: it covers how to read coefficients and compute marginal effects in nonlinear models generally, including the binary discrete-choice models of the Discrete and Limited Dependent Variables chapter as well as the count models above.

8.9.1 Binary Response Models

In a linear model

$$y_i = X_i'\beta + \epsilon, \tag{8.17}$$

$$E(y_i) = X_i'\beta \tag{8.18}$$

When y_i is binary (meaning it takes two values, 1 or 0), We can also do an OLS regression on it, but a more popular way is to use a non-linear model. The most popular choices for modeling binary response are logit model and probit model. Both of them use the same idea: use a link function to map the binary variable into a continuous variable which is a linear function of the predictors.

Suppose p_i is the probability that $y_i = 1$. Then

$$p_i = E(y_i) = g^{-1}(X_i'\beta), \tag{8.19}$$

where $g^{-1}()$ is a function that maps from the linear predictor to y_i, which is often called index function or inverse link function.

This says that the expectation of y_i (equivalently, p_i) is not a linear function of x_i's, but by using the link function, we are still able to model a transformed p_i as a linear function of x_i's:

$$g(p_i) = X_i'\beta. \tag{8.20}$$

$g^{-1}()$ can be a normal CDF then we have a probit model:

$$g^{-1}(z) = \Phi(z) = \int_{-\infty}^{z} \frac{1}{\sqrt{2\pi}} \exp[-(t^2/2)]dt, \tag{8.21}$$

or it can be a logit:

$$g^{-1}(z) = \Lambda(z) = e^z/(1+e^z). \tag{8.22}$$

8.9.2 Marginal Effects

Often times we are interested in the marginal effects of the predictors. In a linear model, it's easy: the coefficient on x_1 means when x_1 increase by one unit, how much will y increase.

In a binary response model,

$$\frac{\partial p}{\partial x} = \frac{\partial g^{-1}(X_i'\beta)}{\partial x} \tag{8.23}$$

In the case of probit:

$$\frac{\partial p}{\partial x} = \frac{\partial \Phi(X_i'\beta)}{\partial x} = \phi(X_i'\beta)\beta, \tag{8.24}$$

where $\phi()$ is the density function of normal distribution.

In the case of logit:

$$\frac{\partial p}{\partial x} = \frac{\partial \Lambda(X_i'\beta)}{\partial x} = \gamma(X_i'\beta)\beta, \tag{8.25}$$

where $\gamma()$ is the density function for logistic distribution:

$$\gamma(X_i'\beta) = \Lambda(X_i'\beta)(1-\Lambda(X_i'\beta)) = p(1-p). \tag{8.26}$$

Therefore for logit, a quick approximation of the marginal effect is to plug in the sample proportion $\bar{p}$ and coefficient estimate $\hat{\beta}$:

$$\frac{\hat{\partial p}}{\partial x} \approx \bar{p}(1-\bar{p})\hat{\beta} \tag{8.27}$$

This evaluates the effect at the mean probability; it is a rough rule of thumb, not the average marginal effect $\frac{1}{n}\sum_i \hat{p}_i(1-\hat{p}_i)\hat{\beta}$, which is what software reports.

The logit model can also be formulated as

$$\log(\frac{p}{1-p}) = \mathbf{X_i'}\boldsymbol{\beta} \tag{8.28}$$

which says the logarithm of the odds (the ratio of the two probabilities) is equal to $\mathbf{X_i'}\boldsymbol{\beta}$. Therefore,

$$p = \frac{\exp(X_i'\beta)}{1+\exp(X_i'\beta)} = \Lambda(X_i'\beta) \tag{8.29}$$

Often times we have discrete valued predictors, such as gender, or other dummy variables. In that case, it makes more sense to ask what's the effect of gender being 1 vs. 0 (female vs. male). Write x_{1i} for the dummy and β_1 for its coefficient, $X_{-1,i}$ for unit i's other covariates and β_{-1} for their coefficients, with the intercept included in $X_{-1,i}$. The average discrete change sets the dummy to 1 and then to 0 for every unit, holding the other covariates at their observed values, and averages the difference over the n units:

$$\frac{1}{n}\sum_{i=1}^{n}\left[\Lambda(\beta_1 + X_{-1,i}'\beta_{-1}) - \Lambda(X_{-1,i}'\beta_{-1})\right]. \tag{8.30}$$

This is what software computes when it predicts with x_1 set to 1, then to 0, and takes the mean difference. It is not the group comparison $E(p \mid x_1 = 1) - E(p \mid x_1 = 0)$, which averages over two different covariate distributions.

8.9.3 Count Data Models

In a count data model such as Poisson regression model or Negative Binomial model, we are modeling log of the expected counts:

$$\log(\mathrm{E}(y)) = \mathbf{X_i'}\boldsymbol{\beta} \tag{8.31}$$

Therefore the marginal effect of x_1, for example, on $\mathrm{E}(y)$ is

$$\frac{\partial \log \mathrm{E}(y)}{\partial x} = \hat{\beta}, \tag{8.32}$$

which means β is the marginal effect of x on log of the expected counts. Suppose the expected counts at x is μ_x, then

$$\frac{(\partial \mu_x)/\mu_x}{\partial x} = \hat{\beta}, \tag{8.33}$$

which says that β represents the marginal effect of percentage change in μ_x with respect to x.

Another way to formulate this is to use the Incidence Rate Ratio Interpretation. Suppose the expected counts at x is μ_x, and expected counts at $x+1$ is μ_{x+1}.

$$\log(\mu_x) = x'\beta. \tag{8.34}$$

Therefore,

$$\mu_x = \exp(x'\beta), \tag{8.35}$$

and

$$\frac{\mu_{x+1}}{\mu_x} = \exp(\beta). \tag{8.36}$$

This says that the exponentiated β is the incidence rate ratio of the expected counts, if x increases by 1.

To calculate the marginal effect of x on y (or $\mathrm{E}(y)$), we need to calculate

$$\frac{\partial \mu_x}{\partial x} = \hat{\beta}\mu_x = \hat{\beta}\exp(X_i'\hat{\beta}), \tag{8.37}$$

if we evaluate μ_x at predicted value.

8.9.4 How to calculate marginal effects in Stata

Stata's margins command is a powerful command to get the predicted value or marginal effects after an estimation command. Check Stata's manual for more details.

8.10 Summary

Poisson regression models the log of the expected count. Its real strength is as a quasi-MLE: it is consistent whenever the conditional mean is right, even with overdispersion or a continuous outcome, as long as the standard errors are robust. Fixed-effects Poisson has no incidental-parameters problem, but it still needs strict exogeneity given the unit effect. Negative binomial adds gamma heterogeneity to allow overdispersion. Hurdle and zero-inflated models treat zeros as coming from a separate process. Zero-truncated models handle samples with no zeros, and there overdispersion breaks a truncated Poisson's consistency.

In all these nonlinear models a coefficient is not a marginal effect. The effect of a regressor on the expected outcome or probability depends on where it is evaluated. The average marginal effect, averaged over the observed covariates, is the standard summary. For a dummy regressor it is the average difference in predicted outcomes between its two values.

9 Panel Data Models

```
# Packages used in this chapter
library(fixest)
```

9.1 Background

Panel data are repeated observations on the same "unit" (we call that cross-sectional units) over time. Panel data models are used since it has advantages over cross-sectional models or time-series models. The major advantage it has over cross-sectional models is: **Controlling for individual heterogeneity**.

For example, Baltagi and Levin (1992) studies cigarette demand across 46 states for the years 1963-88. Consumption is modeled as a function of lagged consumption, price and income. These variables vary across states and time. However, there are other variables that may be time-invariant (z_i) or individual-invariant (w_t). Examples of z_i are religion and, in a panel of adults, completed education. For example, Utah, as a Mormon state, has very low cigarette demand due to religious reason. Generally we consider it does not change over time or change very little over time. Examples of w_t include cigarette commercials on national TV or radio. Panel data models are able to control for individual heterogeneity (or cross-sectional heterogeneity) while cross-sectional models are not. In many (or most) social science studies, there is "unobserved effects" which is embedded in the error term. In other words, we have "omitted variable" problem. Without controlling for it, the estimation results are generally biased and inconsistent.

Panel data give more informative data, more variability and less collinearity among variables. Often time series data suffers from multicollinearity, while panel data has more variability from its cross-sectional units.

The basic unobserved effects model can be written as:

$$y_{it} = \mathbf{x_{it}}\boldsymbol{\beta} + c_i + u_{it} \tag{9.1}$$

where i indexes "units" (can be people, firms, or households, etc.), and t indexes time periods.

Three kinds of regressor appear in this chapter, and it pays to name them once. $\mathbf{x}_{it}$ varies across units *and* time — price, income, and in most applications the treatment itself. z_i varies across units only, so within a unit it is fixed over time; religion and completed education above are the examples. w_t varies across time only, so in any period it is common to every unit; the national TV campaign is one. Which group a variable falls into is what decides almost everything that follows, including which coefficients fixed effects can estimate and how OLS weights the data.

Instruments do not appear until the panel turns dynamic. When they do, they are written $\mathbf{Z}$, which is a different object from the time-invariant z_i here — see the Endogeneity, GMM and Dynamic Panel Data chapters.

Comparing to regular cross-sectional models, there is an extra term c_i. c_i can be treated as a random effect or fixed effect. Traditionally it is distinguished by whether it is estimated as a random variable or a parameter. Modern econometrics distinguishes the estimators by what they assume about c_i and $\mathbf{x_{it}}$. Fixed effects permits any dependence between c_i and the regressor history; it does not require that dependence to exist. Conventional random effects imposes that c_i is uncorrelated with the regressors in every period. Both still need the remaining disturbance u_{it} to be strictly exogenous.

It is true that in some cases it is hard to justify that there is no correlation between c_i and $\mathbf{x_{it}}$, which is one reason that people in economics tend to use fixed-effect models. However, fixed-effect models have their own limitations. One of them is that removing c_i handles only omitted factors that are constant over time. Omitted factors that change over time stay in u_{it}, and they must be uncorrelated with the regressors in every period.

9.2 Random Effect Methods

A random effect model puts c_i into the error term, then estimate by FGLS (feasible GLS). Random effect model is sometimes called "Error-components model" since the overall error term is divided into an individual level error term (c_i) and individual-time level error term (u_{it}).

Under a random effect model, the variance-covariance matrix of the error term becomes:

$$\mathbf{\Omega} = \begin{bmatrix} \sigma_c^2 + \sigma_u^2 & \sigma_c^2 & \cdots & \sigma_c^2 \\ \sigma_c^2 & \sigma_c^2 + \sigma_u^2 & \cdots & \sigma_c^2 \\ \vdots & \vdots & \ddots & \vdots \\ \sigma_c^2 & \sigma_c^2 & \cdots & \sigma_c^2 + \sigma_u^2 \end{bmatrix} \tag{9.2}$$

The random effect estimator is:

$$\hat{\boldsymbol{\beta}}_{\mathbf{RE}} = (\sum_{\mathbf{i=1}}^{\mathbf{N}} \mathbf{X_i'} \hat{\mathbf{\Omega}}^{\mathbf{-1}} \mathbf{X_i})^{\mathbf{-1}} (\sum_{\mathbf{i=1}}^{\mathbf{N}} \mathbf{X_i'} \hat{\mathbf{\Omega}}^{\mathbf{-1}} \mathbf{y_i}) \tag{9.3}$$

It is a special case of GLS. Before calculating $\hat{\beta}_{RE}$, σ_c^2 and σ_u^2 need to be estimated. This is generally done by using residuals from within and between regressions (the method of Swamy and Arora (1972), the default in most software), or from pooled OLS.

9.3 Fixed Effect Methods

Random effect methods assume that c_i be orthogonal to $\mathbf{x_{it}}$. In many applications, the whole point of using panel data is to allow c_i be correlated with $\mathbf{x_{it}}$. We need fixed-effect models in those cases.

For the same model as in the equation above, the fixed-effect methods try to eliminate c_i using fixed effects transformation, or "within transformation". The FE transformation is to "de-mean" each observation by subtracting the group mean. Basically subtracting the group-mean equation below from the unit-level equation above:

$$\bar{y}_i = \bar{\mathbf{x}}_\mathbf{i}\boldsymbol{\beta} + c_i + \bar{u}_i \tag{9.4}$$

where $\bar{y}_i$, etc., means group means. For example, if we have 50 states with state level data of cigarette consumption across years, then groups means states, and we have 50 group means which are state level average cigarette consumption across years.

The reason for "demeaning" is to remove c_i from final estimation. The "demeaned" regression equation is the final regression used in estimation:

$$(y_{it} - \bar{y}_i) = (\mathbf{x}_{it} - \bar{\mathbf{x}}_i)\beta + (u_{it} - \bar{u}_i). \tag{9.5}$$

The unit-specific effect c_i cancels because it is constant over t ($c_i - \bar{c}_i = 0$). Writing the within-transformed variables as $\ddot{y}_{it} = y_{it} - \bar{y}_i$ and $\ddot{\mathbf{x}}_{it} = \mathbf{x}_{it} - \bar{\mathbf{x}}_i$, OLS on the demeaned data gives the **fixed-effects (within) estimator** (here $\mathbf{x}_{it}$ is a $1 \times k$ *row* vector, so $\ddot{\mathbf{x}}'_{it}\ddot{\mathbf{x}}_{it}$ is the $k \times k$ outer product)

$$\hat{\beta}_{FE} = \left(\sum_i \sum_t \ddot{\mathbf{x}}'_{it}\ddot{\mathbf{x}}_{it} \right)^{-1} \sum_i \sum_t \ddot{\mathbf{x}}'_{it}\ddot{y}_{it}. \tag{9.6}$$

This is consistent for β even when c_i is correlated with $\mathbf{x}_{it}$, which is the key advantage of FE over random effects. Consistency still asks for the rest of the usual package: strict exogeneity of the idiosyncratic error conditional on the unit effect, $E[u_{it} \mid c_i, \mathbf{x}_{i1}, \ldots, \mathbf{x}_{iT}] = 0$, enough within-unit variation for the demeaned cross-product to have full rank, and an asymptotic scheme – usually $N \to \infty$ with T fixed. FE allows correlation between c_i and the regressors. It does not allow correlation between u_{it} and them. The price is that any time-invariant regressor (the z_i above) is also swept out by the demeaning and cannot be estimated. The usual variance estimator is $\widehat{V}(\hat{\beta}_{FE}) = \hat{\sigma}_u^2(\sum_i \sum_t \ddot{\mathbf{x}}'_{it}\ddot{\mathbf{x}}_{it})^{-1}$ with $\hat{\sigma}_u^2$ using degrees of freedom $NT - N - k$; in practice one clusters by unit to allow serial correlation within i.

9.4 Fixed effects or random effects: the Hausman test

Random effects is more efficient (it uses both within and between variation) but is consistent only if c_i is uncorrelated with $\mathbf{x}_{it}$; fixed effects is consistent either way. The J. A. Hausman (1978) test compares the two estimators:

$$H = (\hat{\beta}_{FE} - \hat{\beta}_{RE})' \left[\hat{V}(\hat{\beta}_{FE}) - \hat{V}(\hat{\beta}_{RE})\right]^{-1} (\hat{\beta}_{FE} - \hat{\beta}_{RE}) \;\sim\; \chi^2_k \tag{9.7}$$

under the null that random effects is consistent. A large H (small p-value) favors fixed effects. In R this is `phtest(fe, re)` from the `plm` package; in Stata, `hausman fe re`.

Caveat: this classic form requires RE to be efficient. The formula above relies on $\hat{V}(\hat{\beta}_{FE}) - \hat{V}(\hat{\beta}_{RE})$ being positive semi-definite, which in turn requires the RE estimator to actually be the *efficient* estimator under the null – true only under the RE model's homoscedasticity and no-serial-correlation assumptions. If you are using robust or cluster-robust covariance matrices (the usual practice when errors are heteroscedastic or serially correlated), RE is generally no longer efficient, the covariance difference can fail to be positive definite, and the classic Hausman test statistic can be negative or otherwise ill-behaved. In that case, use a regression-based Hausman test instead: augment the RE (or pooled) regression with the within-unit means of the time-varying regressors (Mundlak's approach) or with the demeaned regressors themselves (Wooldridge's regression-based test), and test the added coefficients jointly with a standard (robust) Wald test – this is valid under heteroscedasticity/clustering and reduces to the same idea as the classic Hausman test under its assumptions.

9.5 Collapse to unit means, or keep the long format?

A practical question comes up whenever we hold a panel. Should we average each unit over time and run one regression on the N resulting rows, or keep all NT rows and cluster the standard errors by unit? Many people treat this as a trade-off and assume the long format must carry more information. It usually does not. The answer turns entirely on where the regressors live.

Take the case where every regressor is time-invariant, so the model is $y_{it} = \mathbf{z}_i\gamma + c_i + u_{it}$. With a balanced panel, pooled OLS on the NT rows and OLS on the N unit means give the same $\hat{\gamma}$. Not close — identical. Random effects returns the same number as well. Wooldridge (2010) states the reason plainly: because c_i is there, new observations within a unit carry no information about γ beyond how they move the unit average $\bar{y}_i$. In effect we only have N useful pieces of information. Averaging discards nothing, because $\mathbf{z}_i$ was already constant over t and the within variation was never available to γ in the first place.

What does differ is the inference. The coefficients coincide, but the naive pooled standard error treats the NT rows as independent, while the T rows of a unit share c_i. In the balanced error-components model, write σ_c^2 and σ_u^2 for the variances of c_i and u_{it}, and $\rho = \sigma_c^2/(\sigma_c^2 + \sigma_u^2)$ for the share of error variance due to c_i. Then the true variance of the pooled coefficient on $\mathbf{z}_i$ is $1 + (T-1)\rho$ times the naive one; at $T = 6$ and $\rho = 0.5$ that factor is 3.5. A change of degrees of freedom cannot fix that. The degrees of freedom matter as well: the unit-mean regression uses $N - k$, pooled OLS reports $NT - k$. Moulton (1990) is the classic illustration, and Donald and Lang (2007) show that t_{N-k} is the reference distribution that keeps the size right here – an exact result resting on the

distributional and design assumptions of the group-level regression, not a universal small-sample repair. With few clusters and heteroskedasticity the practical routes are a cluster-robust variance with a few-cluster correction, or a cluster bootstrap. Clustering by unit repairs the standard error when N is large, but it does not repair the degrees of freedom when N is small.

Now put a time-varying $\mathbf{x}_{it}$ into the model. The two options stop agreeing, and not only for the coefficient on $\mathbf{x}_{it}$ — every coefficient moves, the one on $\mathbf{z}_i$ included. Averaging to unit means throws away all within variation, so what survives is the between estimator. Pooled OLS keeps both sources and forces a single coefficient on them, so it returns a variance-weighted average of the within and the between relationship. The two agree only when the within and between coefficients are equal, which is what the Hausman test is looking for in practice. Strictly the null is consistency of random effects – orthogonality of c_i and the regressor history – which *implies* equal within and between coefficients rather than being the same statement. Equality could hold under an alternative, and equality in one sample is not the null at all. So this is not a new problem. It is the FE-versus-RE question in different clothes.

Situation	Collapsed vs long
All regressors time-invariant, balanced panel	Same coefficients; only the degrees of freedom differ
Any time-varying regressor	Coefficients generally differ, including those on $\mathbf{z}_i$
Unbalanced panel	Generally differ, since pooled OLS weights each unit by T_i

The word to hold on to in the last two rows is *generally*. The differences come from the design, not from a theorem: a regressor orthogonal to the rest, or T_i variation that happens to leave the weighting unchanged, can leave particular coefficients coinciding. What you lose is the right to count on it.

The reconciliation is Mundlak's (1978) device, the same one used above for the regression-based Hausman test. Add the unit means $\bar{\mathbf{x}}_i$ to the long regression alongside $\mathbf{x}_{it}$, and the within and between variation carry separate coefficients: $\mathbf{x}_{it}$ picks up the within estimate, $\bar{\mathbf{x}}_i$ picks up the *difference* between the between and within estimates, so the two coefficients sum to what the collapsed regression reports. That second coefficient is the Hausman contrast. With a single time-varying regressor its t-statistic *is* the Hausman test; with several, the test is the joint Wald or F test on all the mean coefficients together, not any one t. Exact numerical agreement with the collapsed regression also depends on balance and on the weighting, so read it as the same comparison rather than an identity to check to the last digit. We keep the within variation and recover the between answer in one regression.

The useful way to remember it: collapse freely when every regressor is constant within the group, and never when one is not.

9.6 What the long format adds

The section above is mostly a warning. The other side is worth seeing directly, so we simulate panels where we know the truth. We use `fixest` here; in Stata the same regressions are `regress`, `xtreg,`

`be` and `xtreg, fe`.

9.6.1 The equivalence, and where it fails

Take 1000 units over 6 periods, one time-invariant regressor z_i, and a true $\gamma = 2$.

```
set.seed(42)
N <- 1000; T <- 6

z   <- rnorm(N)
bal <- data.frame(id = rep(1:N, each = T), t = rep(1:T, N),
                  z  = rep(z, each = T), c = rep(rnorm(N), each = T))
bal$y <- 2 * bal$z + bal$c + rnorm(N * T)
agg   <- aggregate(cbind(y, z) ~ id, data = bal, FUN = mean)

print(c(long      = coef(feols(y ~ z, bal))[["z"]],
        collapsed = coef(feols(y ~ z, agg))[["z"]]), digits = 12)
```

```
         long     collapsed
1.99351677619 1.99351677619
```

The two agree to every digit printed. Now break the balance by dropping the later periods for units with high z_i, so T_i varies, and varies with the regressor.

```
unb  <- bal[!(bal$z > 0.5 & bal$t > 2), ]
agg2 <- aggregate(cbind(y, z) ~ id, data = unb, FUN = mean)

print(c(long      = coef(feols(y ~ z, unb))[["z"]],
        collapsed = coef(feols(y ~ z, agg2))[["z"]]), digits = 12)
```

```
         long     collapsed
2.00682070731 2.01481052322
```

They no longer agree. Pooled OLS weights each unit by T_i; the collapsed regression counts each unit once. Neither is wrong — they answer different questions, and we should know which one we asked. That T_i is a special case of a general rule, set out in the next section.

9.6.2 What collapsing costs

Now the case that matters. Let x_{it} vary over time and make c_i correlated with $\bar{x}_i$, which is the situation fixed effects exists for. The truth is $\beta = 1$.

```
mu <- rnorm(N)
d  <- data.frame(id = rep(1:N, each = T),
                 x  = rep(mu, each = T) + rnorm(N * T),
                 c  = rep(mu + rnorm(N), each = T))
d$y    <- 1 * d$x + d$c + rnorm(N * T)
d$xbar <- ave(d$x, d$id)
dagg   <- aggregate(cbind(y, x) ~ id, data = d, FUN = mean)

between <- feols(y ~ x,        data = dagg)
pooled  <- feols(y ~ x,        data = d, cluster = ~id)
within  <- feols(y ~ x | id,   data = d)
mundlak <- feols(y ~ x + xbar, data = d, cluster = ~id)

pull <- function(m, v) c(coef(m)[[v]], se(m)[[v]])
tab <- rbind(`between (collapsed)` = pull(between, "x"),
             `pooled OLS`          = pull(pooled,  "x"),
             `within / FE`         = pull(within,  "x"),
             `Mundlak: x`          = pull(mundlak, "x"),
             `Mundlak: xbar`       = pull(mundlak, "xbar"))
colnames(tab) <- c("estimate", "std. error")
round(tab, 3)
```

```
                   estimate std. error
between (collapsed)    1.845      0.035
pooled OLS             1.484      0.022
within / FE            0.995      0.014
Mundlak: x             0.995      0.014
Mundlak: xbar          0.850      0.036
```

Read the table down. The collapsed regression returns about 1.85, badly biased, because it is the between estimator and $\bar{x}_i$ carries c_i with it. Pooled OLS returns about 1.48 — closer, since some within variation is mixed in, but still inconsistent. The within estimator returns about 0.99, and only the long format can compute it: once we collapse there is nothing left to demean. Mundlak delivers both at once, exactly as described above — 0.99 on x_{it}, 0.85 on $\bar{x}_i$, and $0.99 + 0.85$ recovers the between estimate.

Mundlak's device is more than a test: it is an estimator in its own right. Wooldridge's correlated random effects (CRE) framework generalizes the idea, allowing unit-level heterogeneity to depend on the time-averages of the regressors while retaining the RE structure. The full CRE treatment, including extensions to nonlinear models, is in the Blog Book.

So the argument for the long format is not precision. It is that demeaning, the Hausman comparison, unit-specific slopes, lagged dependent variables and anything else built on within variation stop existing once the panel is collapsed. Dynamic models are the extreme case, and they get their own chapter, Dynamic Panel Data.

The rule behind that T_i, and behind the variance weighting in the between-versus-within comparison, is not specific to panels. It is set out in Interpreting OLS: Effect Weights and its companion, Outcome Weights.

9.7 Summary

Panel data let us control for a unit effect that is fixed over time. Random effects puts it in the error and estimates by feasible GLS. That is efficient, but it is consistent only if the unit effect is uncorrelated with the regressors. Fixed effects removes the unit effect by demeaning within units. It is consistent when the unit effect is correlated with the regressors, but it still needs strict exogeneity of the remaining error, and it cannot estimate time-invariant regressors. The Hausman test compares the two. Its regression form, via the Mundlak device, stays valid with robust standard errors.

Collapsing to unit means loses nothing when every regressor is constant within units in a balanced panel, apart from the degrees of freedom, which should be $N - k$. With a time-varying regressor, collapsing leaves only the between estimator. In the simulation it returns about 1.85 against a true 1, pooled OLS about 1.48, and fixed effects about 0.99. The Mundlak regression recovers the within and between estimates in one fit.

10 Survival Models

```
# Packages used in this chapter
library(ggfortify)
library(survival)
```

10.1 Survival function

Survival analysis is also called time to event analysis or event history analysis. We use survival models to analyze data that have the following characteristics:

1. the dependent variable is the waiting time till the occurrence of some event;
2. some observations are censored; that is, some units are not in the data set anymore for some reason before an event happens. For some units we observe the event and the exact waiting time; for others it has not occurred, and all we know is that the waiting time exceeds the observation time. However, we believe if they stayed in the study, sooner or later an event would happen.
3. we are trying to explain (predict) the occurrence of an event with some predictors (explanatory variables).

Censoring is what makes this its own subject, and it carries an assumption that belongs before any estimator. Censoring must be **independent**, or noninformative: being censored may not carry information about when the event would have happened. What that requires is not the same for every estimator, and the difference matters.

Kaplan-Meier estimates a *marginal* survival curve and adjusts for nothing, so it needs censoring independent of the event time marginally, in the population the curve describes. Independence conditional on covariates does not rescue it. If sicker patients are censored earlier, the unadjusted curve is biased even when censoring is independent within every covariate stratum.

The Cox partial likelihood conditions on the covariates in the model, so there the weaker conditional version is enough: given those covariates, units censored at a given time must be neither more nor less likely to experience the event than comparable units still under observation.

When censoring depends on covariates and a marginal curve is the target, name the estimand and make the adjustment explicit. Stratify, standardize over the covariate distribution, or weight by the inverse probability of remaining uncensored.

This is an identification assumption, not a technical footnote. It fails whenever units leave for reasons tied to the outcome – patients withdrawing because they are deteriorating, firms vanishing from a sample shortly before they fail, workers leaving a spell study just as re-employment arrives.

Nothing in the data announces the failure and no sample size repairs it. The usual response is to bring the relevant variables into the model, which weakens what the conditional estimators have to assume, and then to argue explicitly why the censoring that remains is unrelated to the event process.

Let T denote the time variable, a nonnegative, continuous random variable with PDF $f(t)$ and CDF $F(t)$, where t is a realization of T. The survival function is defined by

$$S(t) = Pr(T > t) = 1 - F(t) \tag{10.1}$$

The graph of $S(t)$ against t is known as the survival curve. If there is no censoring, then the survival function is easy to estimate by the ratio of the number of units surviving beyond t to the total number of units in the sample. The most common method to estimate the survival function with a survival data set containing censored observations is the Kaplan-Meier method. The basic idea is to use the product of a series of conditional probabilities.

First sort the event times from the smallest to the largest:

$$t_{(1)} \leq t_{(2)} \leq ... \leq t_{(r)} \tag{10.2}$$

Then the survival curve is estimated by:

$$\hat{S}(t) = \prod_{j|t_{(j)} \leq t} (1 - \frac{d_j}{r_j}), \tag{10.3}$$

where r_j is the number of units at risk at $t_{(j)}$ and d_j is the number of "failure" or events at $t_{(j)}$. Note that units censored at $t_{(j)}$ are included in r_j.

The variance of this estimator can be estimated by:

$$\hat{var}(S(t)) = (\hat{S}(t))^2 \sum_{j|t_{(j)} \leq t} \frac{d_j}{r_j(r_j - d_j)}. \tag{10.4}$$

10.2 Hazard function

We are often interested in which periods have the highest or lowest change of "death" or "failure" or other events. That is, we may be interested in the probability that a state ends between t and $t + \Delta t$ conditional on having reached t in the first place.

The probability is

$$\Pr(t < T \leq t + \Delta t | T \geq t) = \frac{F(t + \Delta t) - F(t)}{S(t)} \tag{10.5}$$

The hazard function is defined by

$$h(t) = \lim_{\Delta t \to 0} \frac{\Pr(t < T \leq t + \Delta t | T \geq t)}{\Delta t} = \frac{f(t)}{S(t)} = \frac{f(t)}{1 - F(t)} \tag{10.6}$$

One of the simplest functional forms is the exponential distribution

$$f(t, \theta) = \theta e^{-\theta t} \tag{10.7}$$

Therefore, the hazard function is

$$h(t) = \theta \tag{10.8}$$

A much more flexible functional form is Weibull, with CDF, survival, and density

$$F(t, \theta, \alpha) = 1 - \exp(-(\theta t)^{\alpha}), \quad S(t) = \exp(-(\theta t)^{\alpha}), \quad f(t) = \alpha \theta^{\alpha} t^{\alpha - 1} \exp(-(\theta t)^{\alpha}). \tag{10.9}$$

Hazard function can be shown to be

$$h(t) = \alpha \theta^{\alpha} t^{\alpha - 1} \tag{10.10}$$

However, Weibull distribution does not allow for the possibility that the hazard may first increase then decrease over time. Log-normal distribution does allow this.




Figure 10.1: Various hazard functions from Davidson and MacKinnon (2004)

10.3 Log-likelihood function

Suppose we have n units with a survival function $S(t)$, density $f(t)$, and hazard $h(t)$. If for unit i, the event happens at t_i, it's contribution to the likelihood function is the density at that point:

$$L_i = f(t_i) = S(t_i) h(t_i) \tag{10.11}$$

If no event at t_i, all we know is that the lifetime exceeds t_i. The probability is

$$L_i = S(t_i), \tag{10.12}$$

which is the contribution to the likelihood function from a censored observation.

Therefore, if U denotes the set of uncensored observations, the log-likelihood function for the entire sample can be written as

$$\ell(t, \theta) = \sum_{i \in U} \log h(t_i | \mathbf{X}_i, \theta) + \sum_{i=1}^{n} \log S(t_i | \mathbf{X}_i, \theta) \tag{10.13}$$

10.4 Proportional Hazard Models

10.4.1 Model and likelihood

One class of models that is quite widely used is the proportional hazard models (based on Cox (1972)).

$$h(t|\mathbf{x}_i) = h_0(t) \exp(\mathbf{x}_i' \beta), \tag{10.14}$$

In this model $h_0(t)$ is called baseline hazard rate; it describes the risk for units with $\mathbf{x_i} = \mathbf{0}$. $\exp(\mathbf{x}_i'\beta)$ represents the relative risk, a proportionate increase or decrease in risk, due to $\mathbf{x_i}$. The interpretation of β is that $\exp(\beta_i)$ gives the relative risk change associated with an increase of one unit in x_i, all other explanatory variables remaining constant.

β is estimated by maximizing a partial likelihood function. It is partial likelihood function since the baseline hazard is factored out. Suppose one event happens at time t_j. Conditional on this event the probability that case i dies (meaning this event happened to case i, not other cases in the risk set) is

$$L_i = \frac{h_0(t_j) \exp(x_i' \beta)}{\sum_l I(T_l \geq t_j) h_0(t_j) \exp(x_l' \beta)} = \frac{\exp(x_i' \beta)}{\sum_l I(T_l \geq t_j) \exp(x_l' \beta)} \tag{10.15}$$

We can see here that Cox's model is for continuous time survival data. It is assumed there is no ties at time t_j (in continuous time is not possible for two events to happen at the same time). In reality, we observe ties of event time, because in many situations, we observe grouped data. For example, we observe case m and n have events at the same time j. In that case, in the denominator, whether we include case m in the calculation of the likelihood of case n is ambiguous. In that case, special treatment is needed. The standard approximation here is Breslow's (1974):

$$L_{m \in D(t_j)} \simeq \frac{\prod_{m \in D(t_j)} \exp(x_m' \beta)}{[\sum_{l \in R(t_j)} \exp(x_l' \beta)]^{d_j}} \tag{10.16}$$

where $D(t_j)$ is the set of cases that die at time t_j and d_j denotes the number of cases that die at time t_j. This approximation works well with small number of ties relative to total number of cases at risk. Peto's (1972) approximation is a close relative of it. Efron's (1977) is a finer adjustment that apportions the risk set among the tied cases, and it is the more accurate of the two when ties

are common. The defaults differ by software: R's `coxph` uses Efron, Stata's `stcox` uses Breslow, so the same data can produce slightly different coefficients in the two without either being wrong.

One important feature (disadvantage?) for the proportional hazard model is that for all units, the effect is the same at all time t. To see this:

$$\frac{h_1(t)}{h_2(t)} = \frac{h_0(t)\exp(\mathbf{x}_1'\beta)}{h_0(t)\exp(\mathbf{x}_2'\beta)} = \frac{\exp(\mathbf{x}_1'\beta)}{\exp(\mathbf{x}_2'\beta)}, \tag{10.17}$$

which does not depend on t.

One advantage of Cox's model is that it is considered as a semi-parametric model since it does not have to specify the distribution of the survival time. The estimation process relies only on the order in which events occur, not on the spacing between them. The exact times still matter, because they are what determines who sits in each risk set. What drops out is the baseline hazard, and with it any need to model how far apart the events are. Parameter estimates are derived assuming continuous survival times.

10.4.2 Interpretation

In the equation above, if we would like to calculate the partial effect of x_j on h, we have:

$$\partial h(t|\mathbf{x}_i)/\partial x_j = h_0(t)\exp(\mathbf{x}_i'\beta)\beta_j = \beta_j h(t|\mathbf{x}_i), \tag{10.18}$$


If we do it the other way, plug in $x_j + 1$ will generate

$$h_0(t)\exp(\mathbf{x}_i'\beta + \beta_j) = \exp(\beta_j)h(t|\mathbf{x}_i) \tag{10.19}$$

Therefore, a one unit change in x_j multiplies the hazard by $\exp(\beta_j)$ (a proportionate change of $\exp(\beta_j) - 1$).

10.4.3 Example in R

```
library(ggfortify)
library(survival)
fit <- survfit(Surv(time, status) ~ sex, data = lung)
autoplot(fit)
```



Censored observations are denoted by crosses. This is the so-called Kaplan-Meier curve.

Now let's do a Cox model example.

```
fit2 <- coxph(Surv(time, status) ~ sex + age, data = lung, method="breslow")
summary(fit2)
```

```
Call:
coxph(formula = Surv(time, status) ~ sex + age, data = lung,
    method = "breslow")

  n= 228, number of events= 165

         coef exp(coef)  se(coef)      z Pr(>|z|)
sex -0.512565  0.598957  0.167462 -3.061  0.00221 **
age  0.017013  1.017158  0.009222  1.845  0.06506 .
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

    exp(coef) exp(-coef) lower .95 upper .95
sex     0.599     1.6696    0.4314    0.8316
age     1.017     0.9831    0.9989    1.0357

Concordance= 0.603  (se = 0.025 )
Likelihood ratio test= 14.08  on 2 df,   p=9e-04
Wald test            = 13.44  on 2 df,   p=0.001
Score (logrank) test = 13.69  on 2 df,   p=0.001
```

Examples of parametric survival models: exponential, Weibull, or log-logistic:

```
exp_fit <-  survreg(Surv(time, status) ~ sex + age, data = lung,
 ↪  dist="exponential")
summary(exp_fit)
```

```
Call:
survreg(formula = Surv(time, status) ~ sex + age, data = lung,
    dist = "exponential")
              Value Std. Error     z      p
(Intercept)  6.35967    0.63547 10.01 <2e-16
sex          0.48093    0.16709  2.88  0.004
age         -0.01562    0.00911 -1.72  0.086

Scale fixed at 1

Exponential distribution
Loglik(model)= -1156.1   Loglik(intercept only)= -1162.3
    Chisq= 12.48 on 2 degrees of freedom, p= 0.002
Number of Newton-Raphson Iterations: 4
n= 228
```

```
weibull <-  survreg(Surv(time, status) ~ sex + age, data = lung, dist="weibull")
summary(weibull)
```

```
Call:
survreg(formula = Surv(time, status) ~ sex + age, data = lung,
    dist = "weibull")
              Value Std. Error     z       p
(Intercept)  6.27485    0.48137 13.04 < 2e-16
sex          0.38209    0.12748  3.00  0.0027
age         -0.01226    0.00696 -1.76  0.0781
Log(scale)  -0.28230    0.06188 -4.56 5.1e-06

Scale= 0.754

Weibull distribution
Loglik(model)= -1147.1   Loglik(intercept only)= -1153.9
    Chisq= 13.59 on 2 degrees of freedom, p= 0.0011
Number of Newton-Raphson Iterations: 5
n= 228
```

```
loglog <-  survreg(Surv(time, status) ~ sex + age, data = lung,
 ↪  dist="loglogistic")
summary(loglog)
```

```
Call:
survreg(formula = Surv(time, status) ~ sex + age, data = lung,
    dist = "loglogistic")
               Value Std. Error     z       p
(Intercept)  5.92232    0.53269 11.12 < 2e-16
sex          0.47751    0.14036  3.40 0.00067
age         -0.01401    0.00771 -1.82 0.06946
Log(scale)  -0.56991    0.06543 -8.71 < 2e-16

Scale= 0.566

Log logistic distribution
Loglik(model)= -1152.9   Loglik(intercept only)= -1160.9
    Chisq= 16.07 on 2 degrees of freedom, p= 0.00032
Number of Newton-Raphson Iterations: 4
n= 228
```

One caveat when reading these next to the Cox fit above. `survreg` estimates *accelerated failure time* models, parameterized on log survival time, whereas `coxph` estimates proportional hazards. The coefficients therefore carry *opposite* signs for the same underlying effect: a covariate that raises the hazard shortens expected survival time. A sign flip between the two tables is the expected result of the differing parameterizations, not a contradiction.

Expect the reversal of direction; do not expect mirror-image magnitudes. Exponential and Weibull are the families that are both PH and AFT, and there the two parameterizations are related exactly through the shape parameter, so the AFT coefficient scaled by the shape recovers the PH coefficient up to sign. Log-logistic, the third fit above, is an AFT family that is *not* proportional hazards at all, so it has no PH coefficient to be the mirror of. Setting a log-logistic fit beside a Cox fit compares two different models of the same data. Agreement in direction is then substantive evidence that the covariate acts the same way under both, not an identity delivered by the parameterization. If either model is misspecified, nothing forces even the signs to match.

10.5 Discrete-time Survival Models

Cox (1972) is the *continuous-time* proportional hazards model; the discrete-time analogue models the odds of an event at time t_j given survival up to that point. This proportional-odds specification for the discrete hazard,

$$\frac{h(t_j|\mathbf{x}_i)}{1-h(t_j|\mathbf{x}_i)} = \frac{h_0(t_j)}{1-h_0(t_j)}\exp(\mathbf{x}_i'\beta), \tag{10.20}$$

is a discrete-time *proportional-odds* hazard model rather than a direct discretization of Cox's continuous-time proportional hazards model. (The complementary log-log link is the discrete-time form that corresponds more directly to a grouped continuous-time proportional hazards model.) If we take logs on both sides, we have

$$\text{logit}(h(t_j|\mathbf{x}_i)) = \alpha_j + \mathbf{x}_i'\beta, \tag{10.21}$$

This looks very similar to a logit model. In fact, we can fit a discrete-time proportional hazard model by running a logistic regression on a set of pseudo observations generated by "filling in" the observations between the start time to the event time or time at the end of the study period (censored). The outcome of this process is 0 unless there is an event, then it turns 1. The likelihood then takes the same algebraic form as a sequence of Bernoulli trials, and that is what makes the logistic-regression trick work. The reason is a factorization, not independence. A person contributes the probability of surviving each period, and of failing in the last one if an event is observed; the chain rule writes that as a product of one-period *conditional* probabilities, each equal to h_{ij} or $1 - h_{ij}$ for individual i at time t_j. Rows from the same person are not free of one another. They share covariates and history, and a later row exists only because the earlier ones were survived. The distinction leaves the point estimates alone, which is why the trick is standard, but it governs the covariance: cluster on the person, or fit a frailty model, when unobserved heterogeneity leaves a person's periods dependent given the model (Singer and Willett 1993 has more details).

10.5.1 Do it in Stata

To do a discrete-time survival model in Stata, your data set needs to be set up as by unit-time. For example, observations by company month. The event (or failure) variable needs to be set to zero until the event happens. If no event until the end of the study period, then it is censored. A data set by unit-time can be analyzed by logit model, optionally with clustered standard error. To set up a structure like this, user-written programs called *dthaz* and *prsnperd* (means person-period) can be used.

To do a continuous-time survival model in Stata, your data can be either set up as discrete-time case (in which case you can use a time-varying explanatory variable) or a single-record-per-person data set. Stata has built-in command *stset* to set up a survival analysis structure. Then you can run *stcox* for Cox's proportional hazard model.

10.6 Summary

Survival analysis models the time until an event when some spells are censored. Censoring must be noninformative. Kaplan-Meier needs that marginally, while the Cox model needs it only given its covariates. Nothing in the data can check this assumption. Kaplan-Meier estimates the survival curve as a product of conditional survival probabilities. The hazard function gives the instantaneous risk of the event among units still at risk. Parametric families such as the exponential, Weibull and log-normal impose shapes on it, and each censored unit contributes its survival probability to the likelihood.

The Cox model writes the hazard as a baseline times $\exp(x'\beta)$. It estimates β by partial likelihood without specifying the baseline, and $\exp(\beta_j)$ is the proportional change in the hazard per unit of x_j. Parametric `survreg` fits are accelerated failure time models, so their coefficients carry the opposite sign. Discrete-time hazards can be fitted as a logit on person-period data, clustered by person.

11 Dynamic Panel Data

11.1 When Is It a Problem

Model setup:

$$y_{i,t} = \gamma y_{i,t-1} + X'_{i,t}\beta + C_i + \epsilon_{i,t} \tag{11.1}$$

Pooled regression is biased and inconsistent. The problem is:

$$Cov(y_{i,t-1}, (C_i + \epsilon_{i,t})) \approx \frac{\sigma_c^2}{(1-\gamma)} \tag{11.2}$$

Random effects model is biased and inconsistent, for the same reason that $y_{i,t-1}$ is necessarily correlated with C_i, which is part of the composite error term in the random effect model.

Fixed effect model setup:

$$y_{i,t} - \bar{y}_i = (X_{i,t} - \bar{X}_i)'\beta + \gamma(y_{i,t-1} - \overline{y_{i,-1}}) + (\epsilon_{i,t} - \bar{\epsilon}_i) \tag{11.3}$$

where $\overline{y_{i,-1}}$ is the unit mean of the *lagged* dependent variable. This is not exactly $\bar{y}_i$ (the mean of current $y_{i,t}$). Over the usable periods the two are $(y_{i2}+\cdots+y_{iT})/(T-1)$ and $(y_{i1}+\cdots+y_{i,T-1})/(T-1)$, so their difference is exactly

$$\frac{y_{iT} - y_{i1}}{T-1}, \tag{11.4}$$

the endpoints and nothing else. Balance does not make that vanish – only T growing does, under stability. The distinction is asymptotically negligible and finite-sample real.

Nickell (1981) shows that (see also Anderson and Hsiao 1981)

$$Cov((y_{i,t-1} - \overline{y_{i,-1}}), (\epsilon_{i,t} - \bar{\epsilon}_i)) \approx -\frac{\sigma_\epsilon^2}{T(1-\gamma)^2}[\frac{(T-1) - T\gamma + \gamma^T}{T}] \tag{11.5}$$

which indicates that the demeaned lagged y is correlated with the demeaned error term. The correlation may be large if T is small, but when T is big, then the correlation goes down to near zero. When T is big, we don't have a problem with fixed effect model.

11.2 How Big Is the Bias

When T is small, which is typically the case in many microeconomic settings, then we say the fixed effect model with a lagged dependent variable is inconsistent. But how bad is it?

The limit of $\hat{\gamma} - \gamma$ is approximately $-\frac{(1+\gamma)}{T-1}$. When $T = 10$ and $\gamma = .5$, the bias is about -0.167 which is $1/3$ of the true value.

But a simulation study (Judson and Owen 1999) shows that the bias can be as big as 20% of the true coefficient even when $T = 30$.

11.3 Anderson and Hsiao estimator

Anderson and Hsiao (1981) suggested looking at the first difference estimator:

$$y_{i,t} - y_{i,t-1} = (X_{i,t} - X_{i,t-1})'\beta + \gamma(y_{i,t-1} - y_{i,t-2}) + (\epsilon_{i,t} - \epsilon_{i,t-1}) \tag{11.6}$$

or

$$\Delta y_{i,t} = \Delta X_{i,t}\beta + \gamma \Delta y_{i,t-1} + \Delta \epsilon_{i,t} \tag{11.7}$$

This does not solve the endogeneity problem, since $\Delta y_{i,t-1}$ is still correlated with the error term. AH's idea is to instrument $\Delta y_{i,t-1}$ with the past level $y_{i,t-2}$, or past difference $y_{i,t-2} - y_{i,t-3}$. This estimator is consistent, since neither of these instruments is correlated with $\Delta\epsilon_{i,t}$ – provided the idiosyncratic error is serially uncorrelated. That proviso carries the argument. If $\epsilon_{i,t}$ follows a moving average of order q, the levels $y_{i,t-2-q}$ and older remain valid, so we can reach further back. If instead $\epsilon_{i,t}$ is AR(1), $\epsilon_{i,t} = \rho\epsilon_{i,t-1} + \nu_{i,t}$ with $|\rho| < 1$, no finite lag works. For a stationary series with $y_{i,t} = \gamma y_{i,t-1} + \epsilon_{i,t}$, writing σ_ϵ^2 for the variance of $\epsilon_{i,t}$, the covariance between the level $s \geq 1$ periods back and the differenced error is

$$\mathrm{Cov}(y_{i,t-s}, \Delta\epsilon_{i,t}) = \frac{(\rho - 1)\rho^{s-1}\sigma_\epsilon^2}{1 - \gamma\rho}. \tag{11.8}$$

To derive it, write $y_{i,t-s} = \sum_{k\geq 0} \gamma^k \epsilon_{i,t-s-k}$ and use $\mathrm{Cov}(\epsilon_{i,t-j}, \epsilon_{i,t}) = \rho^j\sigma_\epsilon^2$. The covariance shrinks with s but is never zero. Older lags are weaker violations, not valid instruments. Under AR errors we need an explicit model of the error process, different moment restrictions, or external instruments.

One implication of the assumption is testable, which is the point of the Arellano-Bond test for serial correlation in the differenced residuals. First-order correlation is expected by construction, since $\Delta\epsilon_{i,t}$ and $\Delta\epsilon_{i,t-1}$ share $\epsilon_{i,t-1}$; it is the *second*-order test that carries information. Rejecting AR(2) says $\epsilon_{i,t}$ is serially correlated, so $y_{i,t-2}$ is not a valid instrument. If the correlation is a short moving average, deeper lags restore validity; if it is autoregressive, they do not.

The other regressors need classifying too, because that is what decides how they are instrumented. Strictly exogenous X can serve as its own instrument. Predetermined X – correlated with past shocks but not with current or future ones – is instrumented by its own lags from $t-1$ backwards. Endogenous X is treated like the lagged dependent variable, with lags from $t-2$ backwards.

Misclassifying a regressor here is one of the commoner ways a difference-GMM specification goes quietly invalid.

11.4 Arellano-Bond estimator

Arellano and Bond (1991) expanded the idea by using additional lags of the dependent variable as instruments. For example, both $y_{i,t-2}$ and $y_{i,t-3}$ can be used as instruments. In fact, as t increases, the number of instruments available also increases. In period 3 only $y_{i,1}$ is available. In period 4 $y_{i,1}$ and $y_{i,2}$ are available. In period 5 $y_{i,1}$ and $y_{i,2}$ and $y_{i,3}$ are available, and so on. In other words, we'll have an instrument matrix with one row for each time period that we are instrumenting:

$$Z_i = \begin{bmatrix} y_{i,1} & 0 & 0 & 0 & 0 & 0 & \cdots & 0 & 0 & 0 \\ 0 & y_{i,1} & y_{i,2} & 0 & 0 & 0 & \cdots & 0 & 0 & 0 \\ 0 & 0 & 0 & y_{i,1} & y_{i,2} & y_{i,3} & \cdots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots & \cdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & 0 & \cdots & 0 & y_{i,1} & \cdots & y_{i,T-2} \end{bmatrix} \tag{11.9}$$

This is the so called difference GMM estimator.

11.5 Blundell and Bond estimator

AB model has a problem: when the instruments are weak, the estimator is not good. That will happen when y follows random walk or near random walk. In that case the past levels won't be a good predictor of the future changes.

Blundell and Bond (1998) add a second set of moment conditions in *levels*. The basic idea is that lagged differences $\Delta y_{i,t-1}$ are uncorrelated with the composite error $C_i + \epsilon_{i,t}$ under an additional initial-condition/stationarity restriction, so they can instrument the level equation.

This is implemented by so-called *system GMM*, which stacks two equations and estimates them jointly:

- the **differenced equation** ($\Delta y_{i,t}$), instrumented by lagged *levels* $y_{i,t-2}, y_{i,t-3}, \ldots$ (the Arellano–Bond moments); and
- the **level equation** ($y_{i,t}$ in levels, keeping C_i), instrumented by lagged *differences* $\Delta y_{i,t-1}$.

The level-equation moments are valid only under the extra assumption that the initial deviations from the long-run mean are uncorrelated with C_i (mean stationarity of the process). System GMM is more efficient than difference GMM when y is persistent (near a random walk), and the level equation lets time-invariant regressors enter, which differencing removes altogether. That they appear is not the same as their coefficients being identified. A time-invariant regressor has no within variation, so its own lagged differences cannot instrument it. Identification comes from the level moments, and it requires the regressor to be uncorrelated with the unit effect C_i, or an external instrument for it. For a time-invariant regressor correlated with C_i and lacking such an instrument, system GMM reports a coefficient that is not identified. A practical warning: the instrument count grows with T^2, so *instrument proliferation* can overfit the endogenous regressors and weaken the Hansen J test; lags should be collapsed or capped in practice.

11.6 Diagnostics in practice

The warning about instrument proliferation is easy to state and easy to ignore. It is worth seeing what it does to real estimates, and what the standard diagnostics look like when they are working.

The data are Arellano and Bond's own: an unbalanced panel of 140 UK firms observed between 1976 and 1984, shipped with `plm` as `EmplUK`. The specification is theirs too – log employment on its own first two lags, current and lagged log wages, three lags of log capital, current and lagged log output, with firm and year effects removed and two-step weighting. Only the instrument set changes across the three fits: every available lag of log employment from $t-2$ back, the same capped at three lags, and then the system-GMM version that adds the level-equation moments.

```
library(plm)
data("EmplUK", package = "plm")

dif_full <- pgmm(log(emp) ~ lag(log(emp), 1:2) + lag(log(wage), 0:1) +
                   lag(log(capital), 0:2) + lag(log(output), 0:1) |
                   lag(log(emp), 2:99),
                 data = EmplUK, effect = "twoways", model = "twosteps")

dif_cap  <- pgmm(log(emp) ~ lag(log(emp), 1:2) + lag(log(wage), 0:1) +
                   lag(log(capital), 0:2) + lag(log(output), 0:1) |
                   lag(log(emp), 2:4),
                 data = EmplUK, effect = "twoways", model = "twosteps")

sys      <- pgmm(log(emp) ~ lag(log(emp), 1:2) + lag(log(wage), 0:1) +
                   lag(log(capital), 0:2) + lag(log(output), 0:1) |
                   lag(log(emp), 2:99),
                 data = EmplUK, effect = "twoways", model = "twosteps",
                 transformation = "ld")

diag_row <- function(fit, label) {
  co <- summary(fit, robust = TRUE)$coefficients
  sg <- sargan(fit)
  data.frame(specification = label,
             moments  = ncol(fit$W[[1]]),
             lag1     = round(co[1, 1], 3),
             se       = round(co[1, 2], 3),
             lag2     = round(co[2, 1], 3),
             max_root = round(max(Mod(polyroot(c(-co[2, 1], -co[1, 1], 1)))), 3),
             sargan_p = round(sg$p.value, 3),
             AR1_p    = round(mtest(fit, order = 1)$p.value, 3),
             AR2_p    = round(mtest(fit, order = 2)$p.value, 3))
}

rbind(diag_row(dif_full, "difference GMM, lags 2+"),
      diag_row(dif_cap,  "difference GMM, lags 2-4"),
      diag_row(sys,      "system GMM"))
```

```
                 specification moments  lag1    se   lag2 max_root sargan_p
chisq   difference GMM, lags 2+      40 0.645 0.194 -0.059    0.534    0.162
chisq1 difference GMM, lags 2-4      30 0.437 0.359 -0.081    0.285    0.166
chisq2               system GMM      55 1.114 0.051 -0.099    1.017    0.121
       AR1_p AR2_p
chisq  0.002 0.596
chisq1 0.065 0.848
chisq2 0.028 0.689
```

Read the AR columns first, because they are the ones that can invalidate everything else. First-order serial correlation in the differenced residuals is expected: $\Delta\epsilon_{i,t}$ and $\Delta\epsilon_{i,t-1}$ both contain $\epsilon_{i,t-1}$, so a small AR(1) p-value is the normal state of affairs and says nothing against the specification. The AR(2) test is the informative one, and none of the three fits shows evidence that the instrumenting lag sits too close.

The Sargan column passes in all three cases, which is exactly where the instrument-count warning earns its keep. The full difference-GMM fit imposes 40 moment conditions on 15 parameters across 140 firms; the system-GMM fit imposes 55. A Sargan or Hansen test with many moments and few groups has little power, so a comfortable p-value is weak reassurance rather than strong evidence. The test is being asked to detect a violation using an instrument set flexible enough to fit almost anything.

The coefficients show what that costs. Capping the instruments at three lags cuts the moment conditions from 40 to 30, moves the coefficient on lagged log employment from about 0.65 to about 0.44, and nearly doubles its standard error. That is the honest trade: fewer, better-behaved instruments give less precision. System GMM goes the other way and returns a first-lag coefficient above 1 with a standard error near 0.05, which should stop the reader rather than reassure them. With two lags, the first coefficient alone does not decide whether the process is stable. Write γ_1 and γ_2 for the coefficients on the first and second lags of log employment. The process is stable when both roots z of $z^2 - \gamma_1 z - \gamma_2 = 0$ have modulus below one, and `max_root` reports the larger modulus. A first-lag coefficient of 1.1 with a second-lag coefficient of -0.3, for example, gives roots 0.6 and 0.5, which is stable. For system GMM the coefficients are 1.114 and -0.099 and the larger root is 1.017, so the point estimate implies an explosive employment process. The tight interval around it comes from 55 moment conditions on 140 firms. Every diagnostic in the table passes. The estimate is still not credible.

So what to report: the AR(2) test, the Sargan or Hansen statistic with its degrees of freedom, the instrument count next to the number of groups, and the estimate under at least one narrower instrument set. The last of those does most of the work, because it is the one thing in the list that a flexible instrument set cannot make look good.

11.7 Summary

A lagged dependent variable is correlated with the unit effect, so pooled OLS and random effects are inconsistent. Fixed effects is inconsistent too when T is small, because the demeaned lag is correlated with the demeaned error. This is the Nickell bias, roughly $-(1+\gamma)/(T-1)$. Anderson-Hsiao

first-differences the model and instruments the lagged difference with earlier levels. Arellano-Bond difference GMM uses all available lags as instruments. Blundell-Bond system GMM adds moments in levels, which helps when the series is persistent but needs an initial-condition restriction.

On Arellano and Bond's UK firm data, every diagnostic passes for all three instrument sets. Capping the instruments at three lags moves the employment coefficient from about 0.65 to 0.44 and nearly doubles its standard error. System GMM returns a first-lag coefficient above 1 with a small standard error. With both lags, the larger root of the lag polynomial is 1.017, so the point estimate implies an explosive process, which is not credible. Report the AR(2) test, the Sargan or Hansen statistic with its degrees of freedom, and the instrument count against the number of groups. Also report the estimate under a narrower instrument set.

12 Missing Data

Data set like:

y	x	z	w	q
1	4	2	?	?
4	?	1	2	?
2	?	?	?	?
?	2	1	?	?
?	?	?	?	?

12.1 Different cases under different assumptions

1. Missing completely at Random (MCAR): the occurrence of missing data is not related to the missing value, the values of any other variables, or the pattern of missingness in other variables. Too good to be true situation.
2. Missing at Random (MAR): the probability that a value is missing depends only on data we observe. Write R for the missingness indicator and split the data into an observed and a missing part. MAR is the restriction

   $$R \perp Y_{mis} \mid Y_{obs},$$

   that given what we see, whether a value is missing does not depend on the value we do not see. Two things follow. It is a restriction on the mechanism, so it is untestable from the observed data: the data we have are equally consistent with MAR and with MNAR alternatives. And it is not the statement that the missingness can be modelled. A model for missingness can always be fitted, and fitting one is no evidence that the restriction holds.
3. Missing Not at Random (MNAR): the occurrence of missing values is systematically related to the unobserved value itself, even after conditioning on observed data. This is the hardest case: there is no model-free fix, but it is not hopeless. It can be addressed with selection models, pattern-mixture models, or sensitivity analysis, all of which rely on untestable assumptions about the missingness mechanism.

The MAR case is the one we are interested in.

For MCAR: you lose efficiency if you simply drop the data. For MAR: if you don't model it, you generally suffer bias as well as efficiency.

"Generally" is doing work in that sentence, because the consequence depends on the analysis and not only on the mechanism. Complete-case regression of y on X stays consistent for β when missingness

depends only on X – which is MAR, not MCAR – because dropping rows on the basis of X leaves the conditional distribution of y given X untouched. What breaks consistency is missingness that depends on y itself, or on something unobserved, after conditioning on X. So MAR is not automatically fatal and MCAR is not the only safe case. Which it is depends on what is missing and on what you are trying to estimate.

If we model missingness, for MCAR, you gain efficiency. For MAR, you correct bias and gain efficiency.

12.2 Old methods

1. Listwise deletion

2. Mean imputation.

Problems: * understates variability in the imputed variable * does not recover associations between variables.

So standard errors are in general under-estimated.

3. Regression-based imputation

Use a model such as $x = \alpha_0 + \alpha_1 z + \alpha_2 y$. But that introduce extra noise that are not accounted for (similar to generated variable problem).

4. Interpolation of panel data

That is, to use the observation from last period or a linear interpolation for the same unit.

12.3 Modern methods

- Account for uncertainty in imputed variable
- Use a model to predict missing observation.
- Instead of picking one, pick many
- uncertainty is represented by VCV matrix of the coefficients used to predict missing values.

12.3.1 Basic ideas

Imputation model: $x = \alpha_0 + \alpha_1 z + \alpha_2 y$

Main model: $y = \beta_0 + \beta_1 x + \beta_2 z$

1. Pick M values of α out of the asymptotic distribution, the multivariate normal $N(\hat{\alpha}, \hat{\Sigma})$, using the estimate $\hat{\alpha}$ and its VCV $\hat{\Sigma}$ as the mean and VCV of the distribution.
2. For each of the M draws, predict the missing values *and add a random residual* drawn from the imputation model's estimated error distribution, creating M data sets. Both draws are needed: the draw of α carries parameter uncertainty, and the residual draw carries the irreducible scatter around the prediction. Dropping the residual draw gives what is called "improper" imputation, and reintroduces precisely the understated-variability problem that sank mean imputation and single regression imputation above.

A fully proper procedure draws the residual variance too. Step 1 as written holds $\hat{\Sigma}$, and with it $\hat{\sigma}^2$, fixed at its estimate, so it propagates uncertainty about α but not about σ^2. The Bayesian version draws $\sigma^{2(m)}$ from its posterior – for the normal linear model a scaled inverse-χ^2 – then draws $\alpha^{(m)}$ from $N(\hat{\alpha}, \sigma^{2(m)}(\mathbf{X}'\mathbf{X})^{-1})$, and finally the residual from $N(0, \sigma^{2(m)})$. In large samples the difference is slight. In small ones, fixing σ^2 understates the between-imputation variance.

3. calculate M new estimates and combine them by averaging, $\tilde{\beta} = \frac{1}{M}\sum_{m=1}^{M} \tilde{\beta}_m$, using each of the imputed data sets. then calculate the variance of $\tilde{\beta}$:

$V_\beta = W + (1 + 1/M)B$

where $W = \frac{1}{M}\sum_{m=1}^{M} s_m^2$, $B = \frac{1}{M-1}\sum_{m=1}^{M}(\tilde{\beta}_m - \tilde{\beta})^2$, in other words, the within-imputation and between-imputation variation. The standard error is $\sqrt{V_\beta}$. These formulas (Rubin's rules, Rubin 1987) are written here for a single scalar coefficient β; for the full coefficient vector, W is the average of the within-imputation covariance matrices, B is the between-imputation covariance matrix $\frac{1}{M-1}\sum_m(\tilde{\beta}_m - \tilde{\beta})(\tilde{\beta}_m - \tilde{\beta})'$, and $V_\beta = W + (1 + 1/M)B$ is the total covariance matrix.

12.3.2 MI through Chained Equations (MICE) (by Buuren)

The standard reference is Buuren (2018).

1. Drop an observation only when it carries no usable information: no observed outcome, no auxiliary variable, no weight or design information. A row holding any of these still contributes, to the imputation model if not to the analysis, and discarding it wastes information and can induce selection.
2. Fill in the missing data with random draws from the observed values.
3. Move through the columns and perform single-variable imputation using some method.
4. Replace the original replacements with the fitted replacements. Repeat 3 for a large number of times, or with a convergence criteria.
5. Do 1-4 m times to create m imputed data sets.

Many ways to implement MICE.

- Regression (linear, logit, multinomial), get $\hat{w}$ or $f(\hat{w})$ sample.
- The default is Predictive Mean Matching (PMM)

  a. create predicted value

  b. pick a small set of donor cases that have the closest predicted values (the number of donors is software-specific; e.g., three or five).

  c. randomly choose one of those donors' observed values to impute.

12.3.3 Bayesian Data Augmentation

- MICE iterates a sequence of single-variable conditional models, so it has the form of a Markov chain. The chain is not guaranteed to have the stationary distribution we want. The conditionals are specified one at a time, and separately specified conditionals need not be *compatible*: there may be no joint distribution that has all of them as its conditionals. When they are incompatible the iteration can still settle down, and where it settles can depend on the order in which the variables are visited. Run the convergence diagnostics – trace plots of imputed means and variances across iterations and chains – but read them for what they are. They check mixing. They do not validate the imputation model, and no diagnostic of the observed data can, because MAR is untestable.
- Build a missing data model into a Bayesian model, treating missing values as another parameter to estimate by drawing out of its posterior distribution.

$$p(\beta, y_{mis} \mid y_{obs}) \propto p(y_{obs}, y_{mis} \mid \beta)\, p(\beta)$$

The complete-data model appears once, applied to the observed and missing components together, and the prior is over β alone. If the missingness mechanism is modelled, it enters as its own factor with its own parameters rather than being folded into the prior.

- We integrate out the missing values by sampling from the total distribution, then averaging out the beta distributions over the space of missing data points.
- Ignorability is what allows the missingness mechanism to be left out of that expression, and it asks for more than MAR. It needs MAR *and* distinct parameters – those of the missingness mechanism separate from β – and, in the Bayesian version, priors that factor accordingly. MAR on its own is not sufficient.

12.3.4 FIML

If there are no missing data, the likelihood function is

$L(\mu, \Sigma) = \prod_{i=1} f(y_i|\mu, \Sigma)$

If there are missing values, each unit contributes the density of the part of y_i that was actually observed:

$$L(\mu, \Sigma) = \prod_i f(y_{i,obs}\,;\, \mu_{i,obs},\, \Sigma_{i,obs,obs})$$

where $y_{i,obs}$ is unit i's observed subvector, and $\mu_{i,obs}$ and $\Sigma_{i,obs,obs}$ are the matching subvector of μ and submatrix of Σ. The parameters are the same μ and Σ for everyone; what varies by unit is which rows and columns get selected. Written this way the mechanism is visible – each contribution is the marginal density obtained by integrating out that unit's missing components, and no imputation happens anywhere.

FIML leaves the missingness mechanism out of the likelihood, and it needs the same ignorability conditions as the Bayesian version above to do so: MAR, together with parameters of the mechanism distinct from (μ, Σ). MAR on its own is not sufficient here either.

12.3.5 mi in stata

```
log using mi_10_model2.log, replace

clear
set more off

* set matsize is obsolete since Stata 16 (matrix sizes are now dynamic)
* and can be safely omitted on modern Stata; kept here only because the
* original script targeted an older Stata version.
set matsize 4000

use "CROSSED_ContestUserActivity FEB 2014 temp.dta", clear
*sample 10

mi set flong
*local dummies "x1 x2"
local continuous "CulturalDist_KS5D_ctr tight_ctr targ_tight_ctr
 ↪  user_country_opennessValue_ctr targetcntropennes2010_ctr"

mi register imputed `continuous'
mi impute chained  (regress) `continuous' = has_won has_submitted gender_code
 ↪  expert_cat2 expert_cat3 expert_cat4 submissions_count log_cash_000
 ↪  ave_numConcurrentContest, add(10) rseed(1)
```

```
mi estimate, cmdok: heckprob has_won c.CulturalDist_KS5D_ctr##c.tight_ctr
 ↪  targ_tight_ctr gender_code expert_cat2 expert_cat3 expert_cat4
 ↪  submissions_count, select (has_submitted =
 ↪  c.CulturalDist_KS5D_ctr##c.tight_ctr log_cash_000 gender_code expert_cat2
 ↪  expert_cat3 expert_cat4 submissions_count ave_numConcurrentContest) vce
 ↪  (cluster user_id) difficult nonrtol
```

This is a sample code. In stata, you have to do "mi set", then "mi register" to register variables you need to impute. Then "mi impute chained" if you have multiple variables to impute. Each imputed variable is then predicted from the other imputed variables in the chain and from whatever complete variables are listed after `=`. Registering a variable as regular does not put it in the imputation model; only listing it after `=` does. The command above lists the outcomes and the complete analysis covariates there, so the imputations respect the relationships the analysis estimates. Then do "mi estimate".

12.4 Summary

What missing data cost depends on why the data are missing and on what we estimate. Under MCAR, dropping incomplete rows loses only efficiency. Under MAR, missingness depends only on observed data. That cannot be tested, and ignoring it generally biases the estimates, although complete-case regression stays consistent when missingness depends only on the regressors. Under MNAR no model-free fix exists. Mean imputation and single regression imputation understate the variability of the imputed values and so understate standard errors.

Multiple imputation draws the imputation model's parameters and a residual for each of M imputed data sets, then combines the estimates with Rubin's rules, adding the within- and between-imputation variance. MICE imputes one variable at a time in a chain. Predictive mean matching is the usual default. Bayesian data augmentation and FIML handle the missing values inside the model. These last two need ignorability, which is MAR plus parameters of the missingness mechanism that are separate from the model's.

13 Interpreting OLS: Effect Weights

```
# Setup used throughout this chapter: packages, panel size, table helpers
library(fixest)
N <- 1000; T <- 6
vtab <- function(x, d = 3) knitr::kable(data.frame(value = round(x, d)),
 ↪  col.names = "value")
mtab <- function(x, d = 3) knitr::kable(round(x, d))
```

An OLS regression returns one number. If every observation has the same effect, that number is the effect. If effects differ, OLS still returns one number, and it is generally not the plain average effect. It is a weighted average, and the specification chooses the weights.

The main point of this chapter is one sentence:

> OLS gives more weight to comparisons where the regressor varies more, once the controls have been partialled out.

That one rule produces the overlap weighting of a saturated cross-sectional regression and the within-variation weighting of fixed effects. The chapter derives it, evaluates the weights in each setting, and asks when they change the answer.

Two companion notes work through related applications in causal inference: Weights in OLS in causal inference and OLS cannot be ATE.

13.1 One formula

Let V be the regressor of interest and X the controls, including an intercept, and let the effect differ across observations:

$$y_i = \beta_i V_i + X_i'\gamma + e_i \tag{13.1}$$

Write $\bar{\beta} = n^{-1}\sum_i \beta_i$ for the plain average effect. Regress V on X, call the fitted value $\hat{V}_i$ and the residual $\tilde{V}_i = V_i - \hat{V}_i$. By the Frisch–Waugh–Lovell theorem,

$$\hat{\beta}_{OLS} = \frac{\sum_{i=1}^{n} \tilde{V}_i y_i}{\sum_{i=1}^{n} \tilde{V}_i^2} \tag{13.2}$$

Be careful what $\tilde{V}_i$ is. A single observation has no variation of its own; variation belongs to a collection. What it has is a *deviation* — how far its regressor sits from what the rest of the model predicts for it. With only an intercept, that is distance from the grand mean. With unit fixed effects, it is distance from the unit's own mean. Which mean gets subtracted is the most consequential detail in this section, and we come back to it.

Substitute Equation 13.1 into Equation 13.2:

$$\hat{\beta}_{OLS} = \frac{\sum_{i=1}^{n} w_i \beta_i}{\sum_{i=1}^{n} w_i} + \frac{\sum_{i=1}^{n} \tilde{V}_i e_i}{\sum_{i=1}^{n} \tilde{V}_i^2}, \qquad w_i = \tilde{V}_i V_i \tag{13.3}$$

The control term drops out because $\tilde{V}$ is orthogonal to X by construction. The error term does not. It vanishes in conditional expectation when $E[e_i \mid V, X, \beta] = 0$, and in probability under the usual orthogonality and large-sample conditions. That is when the weighted-average reading is the right one. The outcome-weight identity in the companion chapter carries no such remainder — it is exact in every sample.

The denominator is a sum of squares. Split $V_i = \hat{V}_i + \tilde{V}_i$; the cross term $\sum_i \tilde{V}_i \hat{V}_i$ vanishes by OLS orthogonality, so

$$\sum_{i=1}^{n} \tilde{V}_i V_i = \sum_{i=1}^{n} \tilde{V}_i^2 \tag{13.4}$$

and $\sum_i w_i \geq 0$ always. A single w_i is under no such constraint — it is negative whenever $\tilde{V}_i$ and V_i have opposite signs. Non-negativity of the total says nothing about the terms.

The same split works weight by weight: $w_i = \tilde{V}_i^2 + \tilde{V}_i \hat{V}_i$. The first piece is never negative; it is the identifying variation. The second sums to zero over the sample, so it only moves weight between observations, and it is where negative weights come from. It also drops out of the weighted average whenever $\hat{V}_i \beta_i$ lies in the column space of X, because $\tilde{V}$ is orthogonal to that space. Two cases matter below. With saturated strata as controls and an effect that is constant within a stratum, $\hat{V}_i \beta_i$ is a function of the stratum. With unit fixed effects and a slope β_i that is constant over time within a unit, $\hat{V}_{it} = \bar{V}_i$, and $\sum_t \tilde{V}_{it} \bar{V}_i \beta_i = \bar{V}_i \beta_i \sum_t \tilde{V}_{it} = 0$. In both, only the non-negative $\tilde{V}_i^2$ remain.

The individual weights also depend on where V has its zero. Replace V_i by $V_i + c$. The controls carry an intercept, so $\tilde{V}_i$ does not move and neither does the total $\sum_i \tilde{V}_i^2$; each weight becomes $\tilde{V}_i V_i + c\,\tilde{V}_i$. So a reported share of negative weights is partly a choice of origin rather than a fact about the design, and it should be quoted with the scale of V stated.

Centering is not a free normalization either. Under Equation 13.1, writing $V_i = (V_i - c) + c$ turns $\beta_i V_i$ into $\beta_i (V_i - c) + \beta_i c$, and that second piece is a *heterogeneous* intercept which the common $X_i' \gamma$ cannot absorb unless β_i is constant. Centering therefore changes the structural parameterization. It earns its place when the reference level means something, or when the specification allows heterogeneous intercepts.

When the remainder vanishes,

$$E[\hat{\beta}_{OLS} \mid V, X, \beta] = \frac{\sum_{i=1}^{n} w_i \beta_i}{\sum_{i=1}^{n} w_i} \tag{13.5}$$

This is the chapter in one equation. The controls fix $\tilde{V}_i$, $\tilde{V}_i$ fixes the weights, and the weights fix which average of the β_i OLS estimates. Three things follow. Changing the controls changes the weights, so it changes the estimand and not merely the precision. Individual weights can be negative. And if $\beta_i = \beta$ for everyone the weights stop mattering, since every weighted average of a constant is that constant.

Equation 13.2 reads a second way, as a signed weighting of the observed outcomes rather than of the unobserved effects. That reading opens the companion chapter Interpreting OLS: Outcome Weights.

13.1.1 Where the identifying variation comes from

Take the simplest case, y on a single x with an intercept. The only control is the constant, so $\tilde{V}_i = x_i - \bar{x} \equiv d_i$.


Two quantities have to be kept apart here. The identifying variation is $\tilde{V}_i^2 = d_i^2$, the squared distance from the mean of x, which is what the denominator of Equation 13.2 adds up. The effect weight is $w_i = \tilde{V}_i V_i = d_i x_i$. Substituting $x_i = d_i + \bar{x}$,

$$w_i = d_i^2 + \bar{x}\, d_i \tag{13.6}$$

The two agree in total, since $\sum_i \bar{x} d_i = 0$ — that is Equation 13.4 — and differ observation by observation by $\bar{x} d_i$. The difference is not cosmetic: in the hundred-point sample below every d_i^2 is positive while 49% of the effect weights are negative. Centring the regressor, so that $\bar{x} = 0$, is what collapses the two into one.


The figures draw the identifying variation d_i^2 as the thickness of the ray from the sample centroid $(\bar{x}, \bar{y})$ to each point, with each point's share of the total printed beside it. Squaring concentrates it: a few points far from $\bar{x}$ can dominate the fit while most points near it contribute almost nothing.

```
show <- function(x, y, main = "") {
  xb <- mean(x); yb <- mean(y); d <- x - xb; s <- (y - yb) / d; w <- d^2
  plot(x, y, pch = 16, cex = 1.5, xlim = c(-0.5, 11.5), ylim = c(0, 9),
       xlab = "x", ylab = "y", main = main)
  abline(v = xb, lty = 3, col = "grey55")
  segments(xb, yb, x, y, lwd = 0.6 + 9 * w / sum(w), col = "#9E431899")
  abline(lm(y ~ x), lwd = 2, col = "#0E6473")
  points(x, y, pch = 16, cex = 1.5)
  points(xb, yb, pch = 21, bg = "white", col = "#0E6473", lwd = 2.5, cex = 1.7)
  text(x, y, labels = sprintf("%.1f%%", 100 * w / sum(w)),
       pos = ifelse(y >= yb, 3, 1), cex = .85, col = "#9E4318")
}
show(c(1, 3), c(2, 8), "Two Points")
```




With two points the weights are equal, 50% each.

```
show(c(1, 3, 4), c(2, 8, 5), "Three Points")
```



Add a third point at $x = 4$ and $\bar{x}$ moves to 2.67. The point at $x = 3$ now sits almost on top of $\bar{x}$, so it takes only 2.4% of the weight.

```
show(c(1, 3, 4, 9), c(2, 8, 5, 6), "Four Points")
```



Add an extreme point at $x = 9$ and $\bar{x}$ moves to 4.25. That one observation now carries 65% of the total weight, while $x = 4$ falls to 0.18% because it has landed near the new centroid. The weights are relative to the whole sample: add data anywhere and the weights everywhere change.


The concentration does not wash out in larger samples. We draw $n = 100$ points, $x_i \sim N(5, 2^2)$ and $y_i = 2x_i + \varepsilon_i$ with $\varepsilon_i \sim N(0, 3^2)$, and compare the weight held by the ten points nearest $\bar{x}$ against the ten furthest.

```
set.seed(3)
n <- 100
x <- rnorm(n, 5, 2)
y <- 2 * x + rnorm(n, 0, 3)

xb <- mean(x); yb <- mean(y); w <- (x - xb)^2
o <- order(w, decreasing = TRUE)

plot(x, y, pch = 16, cex = 1, col = "grey40", xlab = "x", ylab = "y", main = "N =
 ↪  100 Points")
abline(v = xb, lty = 3, col = "grey55")
segments(xb, yb, x, y, lwd = 0.3 + 10 * w / max(w), col = "#9E431870")
```

```
abline(lm(y ~ x), lwd = 2.5, col = "#0E6473")
points(xb, yb, pch = 21, bg = "white", col = "#0E6473", lwd = 2.5, cex = 1.8)
```




```
vtab(c(`Weight 10 Nearest Mean (%)`      = 100 * sum(w[o[91:100]]) / sum(w),
       `Weight 10 Furthest (%)`          = 100 * sum(w[o[1:10]]) / sum(w),
       `Effect Weights $d_ix_i < 0$ (%)` = 100 * mean((x - xb) * x < 0),
       `OLS Estimate`                    = coef(lm(y ~ x))[["x"]]), 3)
```

	value
Weight 10 Nearest Mean (%)	0.138
Weight 10 Furthest (%)	35.360
Effect Weights $d_i x_i < 0$ (%)	49.000
OLS Estimate	1.865

The 10% of points furthest from $\bar{x}$ hold 35% of the identifying variation; the 10% closest hold 0.14%. An observation sitting exactly at $\bar{x}$ contributes nothing. The last row gives the share of effect weights that are negative: 49%, in a sample where no d_i^2 can be.

13.2 Binary treatment: OLS weights overlap

Now evaluate $w_i = \tilde{V}_i V_i$ for a binary treatment with saturated controls. Two readings of the same weights follow: Angrist sorts them by covariate stratum, Słoczyński sorts them by treatment status.

Let $D \in \{0,1\}$ be a binary treatment and X discrete strata $x \in \{1,\dots,K\}$, with the controls saturated — one indicator per stratum, so no functional form is imposed within cells. Discreteness is essential: saturated controls require a finite number of cells, so continuous X cannot be saturated. With continuous controls the linear projection $\hat{D}(X)$ is not the cell proportion — it is a linear approximation that can fall outside $[0,1]$ — and the stratum-by-stratum decomposition below does not go through. With continuous X one could bin it, but the result then holds for the chosen bins, not for X itself.

Residualising D_i on a full set of stratum indicators gives $\tilde{D}_i = D_i - \hat{D}(x)$, where $\hat{D}(x) = n_x^{-1} \sum_{i \in x} D_i$ is the sample treatment fraction in cell x. Saturation makes the fitted value equal that cell mean exactly; its population counterpart is the propensity $p(x) = E[D \mid X = x]$. Sum the realised effect weights $w_i = \tilde{D}_i D_i$ within a stratum:

$$w_x = \sum_{i \in x} \tilde{D}_i D_i = \sum_{i \in x} D_i^2 - \hat{D}(x) \sum_{i \in x} D_i = n_x \hat{D}(x) - n_x \hat{D}(x)^2 = n_x \hat{D}(x)(1 - \hat{D}(x)) \tag{13.7}$$

Nothing new has been added: it is $w_i = \tilde{V}_i V_i$ from Equation 13.3 with a binary V, where $D_i^2 = D_i$ makes the sum collapse. The maximum at $\hat{D}(x) = 1/2$ is just the fact that a binary variable has most variance when it is a coin flip.

A single observation with a binary regressor has no slope of its own: forming a difference takes one treated and one control unit. The smallest unit that carries an effect here is the stratum, so what gets weighted is the stratum difference in means.

A word on the labels before going further. $\hat{\delta}_x$ is a difference in *observed* means within a stratum. Calling the aggregate an ATT or an ATE needs the usual identification conditions: conditional exchangeability, so that treatment is as good as randomly assigned within a stratum, and overlap, so that each stratum holds both treated and control units. Saturation delivers the algebra, not the interpretation. Without those conditions the formulas below still hold exactly, but they describe weighted averages of observed contrasts rather than causal effects. The weighting is what OLS does; whether the thing being weighted is an effect is settled outside the regression.

Let $\hat{\delta}_x = \bar{y}_{1x} - \bar{y}_{0x}$ be the stratum difference-in-means. The decomposition follows exactly, with no remainder, from Equation 13.2. Within stratum x, a treated unit has $\tilde{D}_i = 1 - \hat{D}(x)$ and a control has $\tilde{D}_i = -\hat{D}(x)$, so the numerator of Equation 13.2 is

$$\sum_{i \in x} \tilde{D}_i y_i = (1 - \hat{D}(x))\, n_x \hat{D}(x)\, \bar{y}_{1x} - \hat{D}(x)\, n_x (1 - \hat{D}(x))\, \bar{y}_{0x} = w_x\, \hat{\delta}_x,$$

and by Equation 13.4 the denominator is $\sum_x w_x$. Hence

$$\hat{\beta}_{OLS} = \frac{\sum_{x=1}^{K} n_x \hat{D}(x)(1 - \hat{D}(x))\, \hat{\delta}_x}{\sum_{x=1}^{K} n_x \hat{D}(x)(1 - \hat{D}(x))} \tag{13.8}$$

so strata with propensity near $\hat{D}(x) = 0.5$ get the most weight per observation. A stratum that contains only treated or only control units has $\hat{D}(x) = 1$ or 0, so $w_x = 0$: it drops out entirely, because there is no within-stratum comparison to make. A linear model with continuous controls would extrapolate into those strata instead of dropping them.

13.3 Smaller groups get larger weights

Angrist sorts by covariate stratum. Słoczyński sorts by treatment status instead, collapsing everything into two numbers — the effect on the treated and the effect on the untreated. Aggregating the stratum effects $\hat{\delta}_x$ of Equation 13.8 with each stratum's count of treated or untreated units gives

$$\text{ATT} = \frac{\sum_x n_x \hat{D}(x)\,\hat{\delta}_x}{\sum_x n_x \hat{D}(x)}, \qquad \text{ATU} = \frac{\sum_x n_x (1-\hat{D}(x))\,\hat{\delta}_x}{\sum_x n_x (1-\hat{D}(x))} \tag{13.9}$$

and the ATE combines them with their natural shares, $\rho = P(D=1)$. To see this, write $\text{ATE} = E[\tau_i]$ and condition on treatment status:

$$E[\tau_i] = P(D=1)\,E[\tau_i \mid D=1] + P(D=0)\,E[\tau_i \mid D=0] = \rho\,\text{ATT} + (1-\rho)\,\text{ATU}.$$

In the finite-sample, stratum-weighted version the same decomposition holds because $\sum_x n_x \hat{D}(x) = n_1$ and $\sum_x n_x(1-\hat{D}(x)) = n_0$, so

$$\text{ATE} = \rho\,\text{ATT} + (1-\rho)\,\text{ATU} \tag{13.10}$$

A reader may wonder how a control-group average can get any weight at all, since the observation-level weights of Equation 13.3 give every control $w_i = \tilde{D}_i D_i = 0$. A control's own effect never reaches its outcome, so it cannot enter. The ATU gets weight through the strata instead. Under conditional exchangeability the treated units of stratum x have the same average effect as its controls, $\tau(x)$, so the weighted average of treated effects in Equation 13.3 equals $\sum_x w_x \tau(x) / \sum_x w_x$, an average over strata. Słoczyński's result regroups those stratum weights into the shares that define the ATT and the ATU.

The OLS weights of Equation 13.8 go as the propensity *variance* $\hat{D}(x)(1-\hat{D}(x))$; the group weights of Equation 13.9 go as the propensity itself or its complement. The relation between the two is Słoczyński's result. Write $p = \hat{D}(X)$ for the cell propensity. Because the design is saturated — the controls are a full set of stratum dummies — residualising D on the strata leaves exactly $\tilde{D} = D - p$, so the coefficient is $E[\tilde{D}\,Y]/E[p(1-p)]$. Since p is constant within a stratum, Y enters only through the group means $m_j(p) = E[Y \mid p, D=j]$, and, using $\tilde{D} = 1-p$ for treated and $-p$ for untreated units,

$$\beta_{OLS} = \frac{\rho\,E[(1-p)\,m_1(p) \mid D=1] - (1-\rho)\,E[p\,m_0(p) \mid D=0]}{E[p(1-p)]} \tag{13.11}$$

Substitute the within-group linear projections $m_j(p) = \alpha_j + \gamma_j p + \varepsilon_j(p)$, where $\varepsilon_j(p)$ is the projection residual within group $D = j$. The residual terms $\rho\, E[(1-p)\,\varepsilon_1 \mid D=1]$ and $(1-\rho)\, E[p\,\varepsilon_0 \mid D=0]$ vanish identically: a projection residual is orthogonal to 1 and to p within its group, and so to any linear function of p. Linearity of $m_j(p)$ is not needed here; it is needed in the next step, to read the projection contrasts as the ATT and the ATU. What remains is

$$\beta_{OLS} = (\alpha_1 - \alpha_0) + (\gamma_1 - \gamma_0)\,\frac{E[p^2(1-p)]}{E[p(1-p)]} \tag{13.12}$$

The last ratio is moment algebra — an identity in the first three moments of p, exact for any distribution. With $\mu_j = E[p \mid D=j]$, $V_j = \mathrm{Var}(p \mid D=j)$, and w_1, w_0 the weights defined in Equation 13.14 below,

$$\frac{E[p^2(1-p)]}{E[p(1-p)]} = w_1\mu_1 + w_0\mu_0 \tag{13.13}$$

holds identically in the distribution of p. Under linearity $\mathrm{ATT} = (\alpha_1 - \alpha_0) + (\gamma_1 - \gamma_0)\mu_1$ and $\mathrm{ATU} = (\alpha_1 - \alpha_0) + (\gamma_1 - \gamma_0)\mu_0$, so Equation 13.11 through Equation 13.13 combine into the main result,

$$\hat{\beta}_{OLS} = w_1\,\mathrm{ATT} + w_0\,\mathrm{ATU}, \qquad w_1 = \frac{(1-\rho)V_0}{\rho V_1 + (1-\rho)V_0}, \quad w_0 = 1 - w_1 \tag{13.14}$$

where V_j is the variance of the propensity score within group j. This is exact for the effects of the within-group linear projections; for ATT and ATU it is exact when the group means are linear in p and an approximation otherwise.

Two things about Equation 13.14 matter. First, the weight w_1 on ATT falls as the treated share ρ rises: the larger the treated group, the smaller its say. That is the whole of "smaller groups get more weight," and it does not depend on the propensity variances. Differentiating Equation 13.14, $\partial w_1/\partial\rho = -V_0V_1/(\rho V_1 + (1-\rho)V_0)^2 < 0$ for any $V_0, V_1 > 0$. The variances set the level of the weight, not the direction it moves. That derivative holds V_0 and V_1 fixed, so it is a statement about the formula rather than about changing the assignment process: a policy that treats more units usually changes the within-group propensity variances as well, and then both channels move together. Second, when the propensity has equal variance in the two groups, $V_1 = V_0$, the weights collapse to the reversed shares,

$$\hat{\beta}_{OLS} \approx (1-\rho)\,\mathrm{ATT} + \rho\,\mathrm{ATU} \tag{13.15}$$

the same two quantities as the ATE in Equation 13.10, with the weights **swapped**. At $\rho = 0.1$ the coefficient puts 0.9 on the effect for the treated tenth; the ATE puts 0.1 there.

One caveat. Equation 13.15 is exact under equal propensity variance across groups — not "no covariates," though no covariates is one trivial way to get there. Słoczyński calls the condition unlikely to hold exactly and offers ρ as a rule of thumb, cleanest near $\rho = 0\%$, 50%, or 100%.

Simulation — Angrist and Słoczyński weights. We draw $n = 400{,}000$ units in five covariate strata ($X \in \{1, \ldots, 5\}$), each with its own treatment effect $\tau_x = (0, 1, 2, 3, 4)$, so the plain ATE is 2.

The outcome is $y = \tau_x D + X + \varepsilon$. We run the simulation twice with different propensity schedules: once with the treated share near half ($\rho \approx 0.5$, propensities ranging from 0.08 to 0.93 across strata) and once with few treated ($\rho \approx 0.09$, propensities from 0.02 to 0.20). The point is to show that the OLS coefficient moves with the treated share even though the stratum effects do not change.

```
set.seed(21)
n <- 400000
X <- sample(1:5, n, TRUE)

compare <- function(px) {
  d  <- rbinom(n, 1, px[X]); ta <- c(0, 1, 2, 3, 4)[X]
  yy <- ta * d + X + rnorm(n)
  dt <- resid(lm(d ~ factor(X)))
  nx <- as.vector(table(X)); dh <- as.vector(tapply(d, X, mean))
  DIM <- as.vector(tapply(yy[d == 1], X[d == 1], mean) - tapply(yy[d == 0], X[d
 ↪  == 0], mean))
  rho <- mean(d); ATT <- weighted.mean(ta, d); ATU <- weighted.mean(ta, 1 - d)
  c(`$\\rho$`                      = rho,
    `$\\max\\lvert\\text{weight diff}\\rvert$`         =
     ↪  max(abs(as.vector(tapply(dt * d, X, sum)) - nx*dh*(1-dh))),
    `OLS Saturated`              = unname(coef(lm(yy ~ d + factor(X)))[["d"]]),
    `Angrist Weighted`           = sum(nx*dh*(1-dh) * DIM) / sum(nx*dh*(1-dh)),
    `Sloczynski Approx`          = (1 - rho) * ATT + rho * ATU,
    `ATE`                        = mean(ta))
}
res <- cbind(`$\\rho \\approx 0.5$`  = compare(c(.08, .20, .50, .75, .93)),
             `$\\rho \\approx 0.09$` = compare(c(.02, .04, .08, .12, .20)),
             `$\\rho \\approx 0.91$` = compare(c(.80, .88, .92, .96, .98)))
mtab(res, 4)
```

	$\rho \approx 0.5$	$\rho \approx 0.09$	$\rho \approx 0.91$
ρ	0.4913	0.0909	0.9069
max|weight diff|	0.0000	0.0000	0.0000
OLS Saturated	2.0121	2.8740	1.1262
Angrist Weighted	2.0121	2.8740	1.1262
Sloczynski Approx	2.0293	2.8577	1.1403
ATE	1.9978	1.9978	1.9978

The two weight vectors — the stratum sums $\sum \tilde{D}_i D_i$ and Angrist's $n_x \hat{D}(x)(1-\hat{D}(x))$ — are the same, below 10^{-9} (the row for the largest absolute weight difference, 0 to four decimals), and the Angrist-weighted average reproduces the OLS coefficient exactly. With $\rho \approx 0.5$ the coefficient is 2.012 against an ATE of 1.998: close, because near-equal shares are where the *grouping by treatment status* does least — the reversed-share gap is $(1-2\rho)(\text{ATT} - \text{ATU})$, identically zero at $\rho = 0.5$. That is not a general immunity. The stratum weighting of Equation 13.8 survives it, and at any ρ the coefficient still departs from the ATE when the effects line up with $\hat{D}(x)(1-\hat{D}(x))$. Push the treated share to $\rho \approx 0.09$ and the coefficient jumps to 2.874; push it the other way to $\rho \approx 0.91$

and it falls. The two extremes are symmetric: at $\rho = 0.09$ OLS overweights ATT, at $\rho = 0.91$ it overweights ATU. The effects did not change. The propensity schedule did, and with it both the treated share and the distribution of $\hat{D}(x)(1-\hat{D}(x))$ across strata, which is what the Angrist weights use.

Why does OLS overweight the treated when they are few? The FWL residual is $\tilde{D}_i = D_i - \hat{D}(x)$. When ρ is small, $\hat{D}(x)$ is small in most strata: a treated unit has $\tilde{D}_i = 1 - \hat{D}(x) \approx 1$, a control unit has $\tilde{D}_i = -\hat{D}(x) \approx 0$. Treated units are the unusual ones — they sit far from the stratum mean of D — so they carry most of the leverage, the same way a point far from $\bar{x}$ dominates the simple regression. At $\rho = 0.09$ the reversed-share approximation puts weight $1 - \rho = 0.91$ on ATT and only 0.09 on ATU, while the ATE puts 0.09 on ATT and 0.91 on ATU. In observational data where the treated group is small — a common setting — the OLS coefficient can be far from the population average effect even under unconfoundedness, purely because of the weighting.

The reversed-share Słoczyński approximation $(1-\rho)\,\text{ATT} + \rho\,\text{ATU}$ gives 2.029 against the exact 2.012, and 2.858 against 2.874 — within about a percent, but the exact weights of Equation 13.14 are what to report.

13.3.1 The three, side by side

Decomposition	Sorted by	Smallest effect unit	Exact when	Reference
Observation rule	nothing	the unit effect β_i	always	Equation 13.3
Angrist (1998)	covariate stratum x	stratum difference $\hat{\delta}_x$	saturated controls	Equation 13.8
Słoczyński (2022)	treatment status D	group averages ATT, ATU	group means linear in p; reversed shares also need $V_1 = V_0$	Equation 13.15

Read down the last column. The observation rule always holds and says little. Angrist's needs saturation and gives an interpretable weight. Słoczyński's needs more and gives the sharpest statement of the three: the coefficient and the ATE are the same two numbers with the weights reversed. Underneath all of them, Shinkre and Hazlett (2024) show these weights are exactly the bias from fitting a constant effect when effects vary, and argue for estimators that tolerate heterogeneity rather than for diagnosing the weights after the fact.

13.4 Fixed effects: OLS weights movers

In a panel the same rule applies with a different mean subtracted. Unit fixed effects residualise against unit dummies, so $\tilde{V}_{it} = V_{it} - \bar{V}_i$, and grouping the observation weights by unit gives that unit's total weight,

$$W_i = \sum_{t=1}^{T} (V_{it} - \bar{V}_i) V_{it} = \sum_{t=1}^{T} (V_{it} - \bar{V}_i)^2 \tag{13.16}$$

its own within sum of squares. So

$$E[\hat{\beta}_{FE} \mid V, \beta] = \frac{\sum_i W_i \beta_i}{\sum_i W_i} \tag{13.17}$$

A unit whose treatment never changes has $W_i = 0$ and contributes nothing, whatever its effect. Demeaning removes time-invariant confounding; it does not make every unit equally influential.

Pooled OLS drops the dummies, so its deviation is from the grand mean and both channels are in play. Conditioning on the unit and applying the laws of total variance and total covariance splits $\mathrm{Cov}(x, y)$ and $\mathrm{Var}(x)$ each into a within and a between piece, and the pooled slope comes out as a convex combination of the two channel slopes,

$$\operatorname{plim} \hat{\beta}_{OLS} = \frac{V_{within}\, \beta_{within} + V_{between}\, \beta_{between}}{V_{within} + V_{between}} \tag{13.18}$$

with $V_{within} = E[\mathrm{Var}(x \mid G)]$, $V_{between} = \mathrm{Var}(E[x \mid G])$, and each β the corresponding covariance over variance. Fixed effects removes $V_{between}$ by demeaning and isolates β_{within}. With two equal groups $V_{between} = \frac{1}{4}\Delta\mu^2$, so pulling the group means apart inflates the between weight without touching a single within slope. The Panel Data Models chapter develops the decomposition itself.

13.4.1 Within-unit weighting: an extreme case

A fixed-effects estimate is identified by units whose x changes over time. Units whose x barely moves get near-zero weight regardless of their true effect. So the FE coefficient is a within-variation-weighted average over the units whose x moves, not the population average. When the movers and the stayers have different effects, the two diverge.

The simulation below constructs the extreme version. Half the units barely move and have a true slope of zero; the other half move a lot and have a true slope of two. The average slope across all units is one. The DGP draws $N = 1000$ units for $T = 6$ periods, with $y_{it} = b_i x_{it} + \alpha_i + \varepsilon_{it}$, $\alpha_i \sim N(0, 1)$, $\varepsilon_{it} \sim N(0, 1)$. Both groups have $E[x_{it}] = 0$, so they differ only in how much x moves within a unit, not in its level.

Group	Units	sd(x_{it})	True slope b_i
Steady	1–500	0.2	0
Movers	501–1000	3	2

```
set.seed(1)
half <- N / 2
g <- data.frame(id   = rep(1:N, each = T),
                move = rep(c(rep(0.2, half), rep(3, half)), each = T),
                b    = rep(c(rep(0,   half), rep(2, half)), each = T))
g$x <- rnorm(N * T, 0, g$move)
g$y <- g$b * g$x + rep(rnorm(N), each = T) + rnorm(N * T)
```

```
wg <- tapply(resid(feols(x ~ 1 | id, g))^2, g$id, sum)
mtab(cbind(value = c(
  `average of the two true slopes`      = mean(c(0, 2)),
  `FE estimate`                         = coef(feols(y ~ x | id, g))[["x"]],
  `Pooled OLS`                          = unname(coef(lm(y ~ x, g))[["x"]]),
  `weight held by the steady half (%)`  = 100 * sum(wg[1:half]) / sum(wg))), 3)
```

	value
average of the two true slopes	1.000
FE estimate	1.984
Pooled OLS	1.983
weight held by the steady half (%)	0.473

Both estimates are near 2, not the simple average of 1. The steady units hold under half of one percent of the FE weight: their six observations sit almost on top of their own unit mean, and squaring those tiny deviations puts them nowhere in the sum. Pooled OLS agrees because both groups have $E[x_{it}] = 0$ — there is no between-group difference in x levels, so $V_{between} = 0$ and pooled OLS collapses to β_{within}. Both estimators weight by variance, and the movers contribute $\sigma^2 = 9$ against the steady group's 0.04, so neither comes close to the simple average. The two estimators diverge only when the group means of x differ, creating a between channel with its own slope. Shift the steady group's mean x away from zero and the two part company.

13.4.2 Pulling the groups apart

Simulation — pulling the groups apart. The DGP is the same as the within-unit weighting simulation: 500 steady units ($b_i = 0$, $\mathrm{sd}(x) = 0.2$) and 500 movers ($b_i = 2$, $\mathrm{sd}(x) = 3$). The one change is that the steady group's mean x is shifted from 0 to 10 or 100, while the movers stay centred at zero. This opens a gap $\Delta\mu$ between the two group centroids without changing any within-unit slope. The between-slope $\beta_{between} = \Delta\bar{y}/\Delta\mu$ is near zero, because the steady group's slope is zero and the movers' mean y is also near zero. So as $\Delta\mu$ grows, pooled OLS is dragged toward zero while FE, which removes the between channel by demeaning, stays at $\beta_{within} \approx 2$.

```
b_within <- (0 * 0.2^2 + 2 * 3^2) / (0.2^2 + 3^2)
V_within <- 0.5 * (0.2^2 + 3^2)
predicted <- function(shift) V_within * b_within / (V_within + 0.25 * shift^2)

park <- function(shift) {
  set.seed(1)
  half <- N / 2
  move <- c(rep(0.2, half), rep(3, half))
  b    <- c(rep(0,   half), rep(2, half))
  x    <- rnorm(N * T, 0, rep(move, each = T)) + rep(c(rep(shift, half), rep(0,
 ↪  half)), each = T)
  y    <- rep(b, each = T) * x + rep(rnorm(N), each = T) + rnorm(N * T)
```

```
  d    <- data.frame(id = rep(1:N, each = T), x = x, y = y)
  c(`FE estimate`       = coef(feols(y ~ x | id, d))[["x"]],
    `OLS Predicted`     = predicted(shift),
    `OLS Actual`        = coef(lm(y ~ x, d))[["x"]])
}

mtab(t(sapply(c(`shift 0` = 0, `shift 10` = 10, `shift 100` = 100), park)), 3)
```

	FE estimate	OLS Predicted	OLS Actual
shift 0	1.984	1.991	1.983
shift 10	1.984	0.305	0.311
shift 100	1.984	0.004	0.004

Fixed effects stays flat near the theoretical $\beta_{within} = 1.991$ wherever the steady group sits. Plain OLS falls from 1.983 to 0.004 on identical true slopes: as $\Delta\mu$ grows, the between weight $\frac{1}{4}\Delta\mu^2$ swamps the fixed within weight 4.52, and the coefficient is dragged toward $\beta_{between} = 0$ like $9/(4.52 + \Delta\mu^2/4)$, the numerator $\frac{1}{2}(\beta_1\sigma_1^2 + \beta_2\sigma_2^2) = 9$.

The figure below draws this directly. Two groups of 200 points each share the same within-group slopes: movers have $y = 2x+\varepsilon$ with $x \sim N(0, 3^2)$, the steady group has $y = \varepsilon$ with $x \sim N(\text{shift}, 0.2^2)$. Across the three panels the shift increases from 0 to 10 to 100, opening a gap $\Delta\mu$ between the group centroids without changing any within-group slope. The dashed line connects the two centroids (the between-slope), and the solid line is the pooled OLS fit. As the centroids separate, the between channel dominates and the pooled fit rotates from $\beta_{within} \approx 2$ toward $\beta_{between} \approx 0$.

```
set.seed(5)
m <- 200
draw <- function(shift, main) {
  xw <- rnorm(m, 0, 3);       yw <- 2 * xw + rnorm(m)
  xs <- rnorm(m, shift, 0.2); ys <- rnorm(m)
  x <- c(xw, xs); y <- c(yw, ys)
  plot(x, y, pch = 16, cex = .5, col = c(rep("#9E431855", m), rep("#0E6473", m)),
       xlab = "x", ylab = "y", main = main)
  abline(v = mean(x), lty = 3, col = "grey50")
  segments(mean(xw), mean(yw), mean(xs), mean(ys), lty = 2, lwd = 2, col =
   ↪  "grey30")
  points(c(mean(xw), mean(xs)), c(mean(yw), mean(ys)), pch = 21, bg = "white",
   ↪  cex = 1.4)
  abline(lm(y ~ x), lwd = 2, col = "#0E6473")
  dmu <- mean(xs) - mean(xw)
  Vb  <- 0.25 * dmu^2
  Vw  <- 0.5 * (var(xs) + var(xw))
  c(`Within Slope (movers)` = unname(coef(lm(yw ~ xw))[2]),
    `Within Slope (steady)` = unname(coef(lm(ys ~ xs))[2]),
    `Between Slope`       = unname((mean(ys) - mean(yw)) / dmu),
```

```
    `Between Weight (%)` = 100 * Vb / (Vb + Vw),
    `OLS Estimate`       = unname(coef(lm(y ~ x))[2]))
}
op <- par(mfrow = c(1, 3), mar = c(4, 3.5, 2.5, 0.5))
a <- draw(0,   expression(Delta*mu == 0))
b <- draw(10,  expression(Delta*mu == 10))
c_ <- draw(100, expression(Delta*mu == 100))
```




Figure 13.1: Pooled OLS as a within/between blend (Equation 13.18). Points are the two groups, open circles their centroids. The dashed line is the between-slope $\beta_{between} = \Delta\bar{y}/\Delta\mu$ through the centroids; the solid line is the pooled fit $\hat{\beta}_{OLS}$. The within slopes are identical across panels; widening $\Delta\mu$ (left to right) inflates $V_{between}$ and rotates $\hat{\beta}_{OLS}$ toward $\beta_{between}$.

```
par(op)
```

```
mtab(cbind(`$\\Delta\\mu = 0$` = a, `$\\Delta\\mu = 10$` = b, `$\\Delta\\mu =
 ↪  100$` = c_), 3)
```

	$\Delta\mu = 0$	$\Delta\mu = 10$	$\Delta\mu = 100$
Within Slope (movers)	2.019	2.009	2.002
Within Slope (steady)	0.007	-0.288	-0.136
Between Slope	0.882	-0.007	-0.006
Between Weight (%)	0.026	85.113	99.833
OLS Estimate	2.010	0.291	-0.002

Read across the three columns. The within slopes barely change — about 2 for the movers, indistinguishable from zero for the steady strip. What changes is the between weight: 0.03% when the centres coincide, 85% at $\Delta\mu = 10$, and 99.8% at $\Delta\mu = 100$. So the OLS fit falls from 2.01 to 0.29 to -0.002 on identical true slopes.

The between slope reported in the first column is meaningless — and harmlessly so: with the centres nearly on top of each other $\Delta\mu \approx 0$, an unstable ratio is multiplied by a vanishing weight and contributes nothing.

The strip has influence in the second and third panels despite barely moving. Influence is distance from the mean of x, and the strip sits $\Delta\mu$ units from it; what it contributes is not its slope but its position — it anchors one end of the dashed line.

A group dummy removes that channel. Demeaning subtracts each cloud's *own* mean, sliding the shifted strip back to the centre, so every panel collapses onto the first — centroids coinciding. Run OLS on that and the solid fit swings back up to the within slope. That is all fixed effects is, and why its slope matches the leftmost panel's regardless of how far the strip had been pushed.

So "the weight is how much x moves" is the demeaned case only. It is exact once unit effects are removed, and false in general, because with the halves at different levels of x part of the fitted line is a comparison *between* them rather than a weighting of within-unit slopes.

13.5 When does the weighting change the answer?

The variance-weighted average equals the plain average whenever effects are unrelated to the weights. The weighting tilts the estimate only when units whose x moves more have systematically different slopes — that is, when b_i covaries with w_i.

The gap is exactly a covariance. For any weights w_i,

$$\frac{\sum_i w_i \beta_i}{\sum_i w_i} - \bar{\beta} = \frac{\mathrm{Cov}_n(w_i, \beta_i)}{\bar{w}} \tag{13.19}$$

If high-effect units carry more weight the coefficient exceeds the plain average; if they carry less it falls below; if weights and effects are unrelated the gap is zero. Nothing about the estimator is broken in any of the three cases.

The following two designs illustrate the distinction. Both draw $N = 1000$ units, $T = 6$ periods, with $\mathrm{sd}_i(x) \sim U(0.2, 3)$ and $y_{it} = b_i x_{it} + \alpha_i + \varepsilon_{it}$. Both share the same x. What differs is how the true slopes b_i relate to regressor volatility:

- **(A) Unrelated:** $b_i \sim N(1, 1)$, independent of $\mathrm{sd}_i(x)$.
- **(B) Aligned:** $b_i = 1 + 0.8(\mathrm{sd}_i - 1.6)$, so the slope rises with volatility.

Both designs average to $\bar{b} \approx 1$. The question is whether the FE estimate — a variance-weighted average of the b_i — differs from that plain average.

```
set.seed(7)
sdX <- runif(N, 0.2, 3)

panel <- function(b) {
  h   <- data.frame(id = rep(1:N, each = T),
                    x  = rnorm(N * T, 0, rep(sdX, each = T)))
  h$y <- rep(b, each = T) * h$x + rep(rnorm(N), each = T) + rnorm(N * T)
```

```
  h
}
compareAB <- function(h, b) {
  w <- tapply(resid(feols(x ~ 1 | id, h))^2, h$id, sum)
  c(`FE estimate`            = coef(feols(y ~ x | id, h))[["x"]],
    `true slopes, plain avg` = mean(b),
    `true slopes, weighted`  = sum(w * b) / sum(w))
}

b1 <- rnorm(N, 1, 1)
b2 <- 1 + 0.8 * (sdX - 1.6)
h1 <- panel(b1); h2 <- panel(b2)

mtab(cbind(`(A) unrelated` = compareAB(h1, b1),
           `(B) aligned`   = compareAB(h2, b2)), 4)
```

	(A) unrelated	(B) aligned
FE estimate	0.9813	1.5338
true slopes, plain avg	0.9716	1.0140
true slopes, weighted	0.9816	1.5422

In (A) all three numbers agree to within sampling noise: the weighting is there but costs nothing, because the slopes are independent of the weights. In (B) the FE estimate is 1.53 while the plain average of the true slopes is 1.01, and the weighted average, 1.54, confirms what the estimator was tracking all along — it overweights the high-slope units precisely because they are the high-variance units.

FE still removes confounding from the unit effects α_i — that is its job, and it does it in both designs. What FE does not do is change the within-variance weighting. Even after demeaning, OLS leans on units whose x moves more, and in design (B) those units happen to have higher slopes. The gap between the FE estimate and the plain average is not a failure of FE; it is what OLS does by construction, and it matters only when b_i covaries with $\mathrm{Var}_i(x)$. Without knowing the true slopes, the problem is undetectable from the data alone — the FE estimate looks well-behaved, standard errors are fine, it is just consistently estimating a different parameter than the simple average.

Reweighting to recover the simple average. One response is weighted least squares. The FE weight on unit i is proportional to its within-variance $\hat{\sigma}_i^2 = \sum_t (x_{it} - \bar{x}_i)^2$. In this model, running WLS-FE with weight $1/\hat{\sigma}_i^2$ gives each unit equal influence, so the estimate targets the simple average of slopes rather than the variance-weighted average. This is the logic behind Gibbons, Suárez Serrato, and Urbancic (2018), who formalize the reweighting to recover what they call the "unit-average treatment effect," and it is the continuous analog of Słoczyński's reweighting in the binary case.

The trade-off is efficiency. The high-variance units are the ones whose slope is best estimated — their x moves a lot, so there is real information there. Downweighting them inflates the standard error. We are giving up precision to change which parameter we estimate.

The weights are also not always available. A unit whose x never moves has $\hat{\sigma}_i^2 = 0$, so its weight is undefined, and a unit that barely moves gets an explosive one. A panel with true stayers forces a choice — drop them or bound the weights — and either choice changes the population the estimate refers to. Inverse-within-variation weighting swaps a variation-weighted target for a unit-weighted one; whether that is the right swap depends on the estimand, not on the weighting being more neutral.

The table below adds the reweighted FE estimate to both designs.

```
compareAB2 <- function(h, b) {
  wvar <- tapply(resid(feols(x ~ 1 | id, h))^2, h$id, sum)
  inv_w <- 1 / tapply(resid(feols(x ~ 1 | id, h))^2, h$id, mean)
  h$rw <- rep(inv_w, each = T)
  c(`FE estimate`            = coef(feols(y ~ x | id, h))[["x"]],
    `Reweighted FE`          = coef(feols(y ~ x | id, h, weights = ~rw))[["x"]],
    `true slopes, plain avg` = mean(b),
    `true slopes, weighted`  = sum(wvar * b) / sum(wvar))
}

mtab(cbind(`(A) unrelated` = compareAB2(h1, b1),
           `(B) aligned`   = compareAB2(h2, b2)), 4)
```

	(A) unrelated	(B) aligned
FE estimate	0.9813	1.5338
Reweighted FE	0.9781	1.0240
true slopes, plain avg	0.9716	1.0140
true slopes, weighted	0.9816	1.5422

In (A) the reweighted estimate agrees with the standard FE estimate — reweighting changes nothing when effects are independent of variance. In (B) the reweighted estimate falls from 1.53 back to near 1.01, recovering the simple average. The price is a larger standard error, because we have deliberately downweighted the most informative units.

Standard errors do not reveal any of this. An estimate can be precise, and unbiased for its own weighted target, while answering a different question from the one that was asked. The issue is the estimand, not estimator failure.

13.6 Start with the estimand

Before interpreting an OLS coefficient, ask which average effect is the substantive target: the average across people, the average among the treated, the average across firms, the effect where there is overlap. Then compare that target with the weights the specification implies.

If the two differ, the response depends on the application. Report the weighted estimand for what it is, model the heterogeneity, or reweight toward the population of interest. Reweighting is not free

— observations with little residual variation carry little information, so giving them more influence costs precision, as the reweighted fixed-effects estimate above shows.

The lesson is not to avoid OLS. It is to stop calling the coefficient "the average effect" without checking which average the regression built.

13.7 Summary

1. **One rule.** OLS weights each observation by $w_i = \tilde{V}_i V_i$, the regressor times what is left of it after the controls (Equation 13.3). The weights sum to $\sum_i \tilde{V}_i^2 \geq 0$; individual weights need not be positive. Two conditions travel with that reading: the remainder in Equation 13.3 has to vanish, in conditional expectation or in probability, and the individual weights shift with the origin of V.
2. **Cross-section.** With binary treatment and saturated strata the weight on a stratum is $n_x \hat{D}(x)(1-\hat{D}(x))$ (Equation 13.8): overlap gets the weight, and a stratum with no comparison gets none. Sorted by treatment status instead, the same weights put $1-\rho$ on the ATT and ρ on the ATU (Equation 13.15) — the ATE's own shares, reversed. Both qualifications from the body survive into the summary: the ATT/ATU reading is exact when the group means are linear in the propensity score, and the reversed shares need equal propensity variance across the two groups.
3. **Panel.** Under fixed effects a unit's weight is its own within sum of squares (Equation 13.16): movers are the estimate, stayers are not. Drop the dummies and group-mean differences re-enter through $V_{between}\beta_{between}$ (Equation 13.18).
4. **When it matters.** The weighting always happens. It moves the answer only when the effects line up with the weights, and then by exactly $\text{Cov}_n(w_i, \beta_i)/\bar{w}$ (Equation 13.19).

Everything here is the *effect* reading — weights on the unobservable β_i, answering what $\hat{\beta}$ estimates. The same FWL identity also weights the *observed* outcomes, a signed and fully computable combination that answers which rows drive the number and reaches, unchanged, into estimators far beyond OLS. That is the companion chapter, Interpreting OLS: Outcome Weights.

14 Interpreting OLS: Outcome Weights

```
# Setup used throughout this chapter: colours and table helpers
teal <- "#0E6473"; rust <- "#9E4318"
vtab <- function(x, d = 3) knitr::kable(data.frame(value = round(x, d)),
 ↪  col.names = "value")
mtab <- function(x, d = 3) knitr::kable(round(x, d))
```

The companion chapter Interpreting OLS: Effect Weights treats the OLS coefficient as a weighted average of unobserved effects and asks whose effect $\hat{\beta}$ estimates. We can read the same Frisch–Waugh–Lovell identity (Frisch and Waugh 1933; Lovell 1963) as a weighting of observed outcomes. This second reading asks which rows move $\hat{\beta}$ and in which direction. Both sets of weights can be calculated in OLS, but only here is the thing being weighted also observed: every contribution $\omega_i y_i$ is in the data. The outcome representation also extends beyond OLS.

14.1 The other reading of FWL

Residualise the regressor of interest V on the controls X, write $\tilde{V}_i$ for the residual, and recall the residual-on-residual form of the coefficient, Equation 13.2:

$$\hat{\beta} = \frac{\sum_{i=1}^{n} \tilde{V}_i y_i}{\sum_{i=1}^{n} \tilde{V}_i^2}. \tag{14.1}$$

The previous chapter groups $\tilde{V}_i$ with V_i and writes $\hat{\beta}$ as a weighting $w_i = \tilde{V}_i V_i$ of effects (Equation 13.3). If we instead group each residual with its outcome y_i, the same expression becomes a linear combination of the observed outcomes:

$$\hat{\beta} = \sum_{i=1}^{n} \omega_i y_i, \qquad \omega_i = \frac{\tilde{V}_i}{\sum_{j=1}^{n} \tilde{V}_j^2}. \tag{14.2}$$

Equation 14.2 is the same algebra rearranged, but a different object from the effect weights.

	Weights on outcomes ω_i	Weights on effects w_i
Weights what	the observed y_i	that observation's own effect β_i

	Weights on outcomes ω_i	Weights on effects w_i
Sign	signed, sums to zero	can be signed; non-negative for binary D with saturated controls
Computable	weight and weighted outcome	weight yes; weighted effect no, because β_i is unseen
Holds when	always, it is algebra	as a causal average when the error remainder vanishes
Answers	which rows drive $\hat{\beta}$, any wrong sign	is this the effect I meant, whose effect

Two constraints pin ω_i down, both from Equation 14.2. An intercept among the controls makes $\sum_i \tilde{V}_i = 0$, so the weights **sum to zero**. And $\sum_i \tilde{V}_i V_i = \sum_i \tilde{V}_i^2$ makes the weighted regressor **sum to one**, $\sum_i \omega_i V_i = 1$.

Use this reading when overlap is in doubt and a row may enter with the wrong sign. For binary treatment, the effect weight $w_i = (D_i - \hat{D}_i)D_i$ can flag a treated row with $\hat{D}_i > 1$, but every control has effect weight zero; the outcome weights expose both sides of the realised comparison. Keep the effect reading for questions about the estimand.

14.2 What the outcome weights look like

Take the simple regression of y on x with an intercept, so $\tilde{V}_i = x_i - \bar{x} = d_i$ and

$$\omega_i = \frac{d_i}{\sum_j d_j^2}. \tag{14.3}$$

Unlike the leverage weights d_i^2 on the observed deviation slopes in the companion chapter — non-negative, growing with the *square* of distance — this is **linear in d_i, and so signed**: a point left of $\bar{x}$ enters $\hat{\beta}$ with a negative coefficient on its outcome, a point right of it positive. The structural effect weight is instead $d_i x_i$, which need not equal d_i^2. The outcome weight is not a slope and does not involve $\bar{y}$; it is simply a signed weight on the observed y_i.

Simulation — simple regression. To see it, we simulate $n = 100$ points — regressor $x_i \sim N(5, 2^2)$ and outcome $y_i = 2x_i + \varepsilon_i$ with $\varepsilon_i \sim N(0, 3^2)$ — then size each point by $|\omega_i|$ and colour it by sign.

```
set.seed(3)
n <- 100; x <- rnorm(n, 5, 2); y <- 2 * x + rnorm(n, 0, 3)
xb <- mean(x); d <- x - xb; om <- d / sum(d^2); col <- ifelse(om >= 0, teal,
 ↪  rust)
R <- d > 0; L <- d < 0
xR <- sum(abs(d[R]) * x[R]) / sum(abs(d[R])); yR <- sum(abs(d[R]) * y[R]) /
 ↪  sum(abs(d[R]))
xL <- sum(abs(d[L]) * x[L]) / sum(abs(d[L])); yL <- sum(abs(d[L]) * y[L]) /
 ↪  sum(abs(d[L]))
```

```
plot(x, y, pch = 16, cex = 0.35 + 2 * abs(om) / max(abs(om)),
     col = paste0(col, "99"), xlab = "x", ylab = "y", main = "N = 100 Points")
abline(v = xb, lty = 3, col = "grey55")
abline(lm(y ~ x), lwd = 1.3, lty = 2, col = "grey55")
segments(xL, yL, xR, yR, lwd = 2.6, col = "grey15")
points(c(xL, xR), c(yL, yR), pch = 21, bg = c(rust, teal), col = "white", cex =
 ↪  2.3, lwd = 2)
```




Figure 14.1: Outcome weights in a 100-point sample. Each small point's **size scales with its weight** $|\omega_i|$, coloured by sign — teal ($\omega_i > 0$) pushes $\hat{\beta}$ up, rust ($\omega_i < 0$) pulls it down; the dotted line is the fulcrum $x = \bar{x}$. The two large dots are the $|d|$-weighted centroids of the points left and right of $\bar{x}$; the solid segment joining them has slope $\hat{\beta}$, and the dashed line is the OLS fit (same slope, shifted through the grand mean).

```
vtab(c(`$\\sum_i \\omega_i$`   = sum(om),      `$\\sum_i \\omega_i x_i$` = sum(om
 ↪  * x),
       `positive mass` = sum(om[om > 0]), `negative mass` = sum(om[om < 0]),
       `OLS`           = coef(lm(y ~ x))[[2]], `$\\sum_i \\omega_i y_i$` = sum(om
        ↪  * y)), 4)
```

	value
$\sum_i \omega_i$	0.0000
$\sum_i \omega_i x_i$	1.0000
positive mass	0.2486
negative mass	-0.2486
OLS	1.8648
$\sum_i \omega_i y_i$	1.8648

The printed sums confirm Equation 14.2: the weights split at $\bar{x}$ into a positive mass ($\approx +0.25$) and an equal negative one (≈ -0.25), and $\sum_i \omega_i x_i = 1$.

But the figure shows more. Split the points at $\bar{x}$, weight each side by $|d_i|$, and take its centroid — the two large dots. **The segment joining them has slope $\hat{\beta}$:** the OLS slope is the rise over run between the low-x and high-x weighted centres.

The dashed fit shares that slope, shifted through the grand mean, so the two coincide only when the sides balance. This collapse to two points falls out of the slope formula.

14.2.1 Why it collapses to two points

Write $d_i = x_i - \bar{x}$. Because $\sum_i d_i = 0$, the positive and negative sides carry equal total weight $W = \sum_{i:d_i>0} d_i$. An inner product with d is therefore a between-group contrast: for any variable v,

$$d \cdot v = W(\bar{v}_R - \bar{v}_L), \tag{14.4}$$

where $\bar{v}_R$ and $\bar{v}_L$ are the $|d_i|$-weighted means on each side of $\bar{x}$. The slope $\hat{\beta} = d \cdot y \,/\, d \cdot x$ is the ratio of two such contrasts; W cancels:

$$\hat{\beta} = \frac{\bar{y}_R - \bar{y}_L}{\bar{x}_R - \bar{x}_L}. \tag{14.5}$$

That is the rise over run between the two $|d|$-weighted centroids — the two dots in the figure.

14.3 When the sign flips

In simple regression the sign of ω_i is just the side of the mean: half the weights are negative, structurally, and a low-x point *should* count against a high-x one. With controls $\tilde{V}_i$ becomes a *residual*, and a row can flip to the sign its raw position denies. Outcome weights reveal that realised extrapolation for treated and control rows alike; binary effect weights omit the control rows because $D_i = 0$ makes their effect weight zero.

With a binary treatment D and covariate x, the weight $\omega_i \propto \tilde{V}_i = D_i - \hat{D}_i$, with $\hat{D}$ the fitted linear projection of D on $(1, x)$. Regressed on D alone, every treated unit is positive and every control negative — the *structural* pattern, producing $\hat{\beta} = \bar{y}_1 - \bar{y}_0$. Add x and the fitted $\hat{D}_i$ can leave $[0, 1]$, often where treatment and control overlap poorly. A treated unit with $\hat{D}_i > 1$ gets $\tilde{V}_i < 0$, so its outcome is subtracted from $\hat{\beta}$; a control with $\hat{D}_i < 0$ gets $\tilde{V}_i > 0$, so its outcome is added. These are **wrong-sign outcome weights**, evidence that the fitted linear comparison is extrapolating.

Two qualifications keep this diagnostic in its lane. First, out-of-range the other way is not a sign reversal: a treated unit with $\hat{D}_i < 0$ remains positive and a control with $\hat{D}_i > 1$ remains negative. Second, $\hat{D}_i$ is a sample fitted value. Leaving $[0, 1]$ is evidence of realised linear extrapolation, not proof that the population propensity is out of range or that common support literally fails; sampling error and functional-form approximation can also produce it. Conversely, in-range fits do not establish overlap or the conditional mean-independence assumption needed for a design-based causal interpretation. With a continuous regressor there is no bounded range to leave, so leverage rather than sign carries the diagnostic.

A wrong-sign outcome weight means the fitted regression contrast reaches beyond its conventional treated-versus-control comparison, and $\hat{\beta}$ can leave the convex hull of the observed outcomes. Whether it matters depends on the data — a sensitivity check is one trim away.

Simulation — thin overlap on a curved baseline. The DGP: 150 controls near $x = 2.5$ and 150 treated near $x = 4.5$ (both sd 1.1), with $y_i = D_i + 0.8\,(x_i - 3.5)^2 + 0.3\,x_i + \varepsilon_i$, $\varepsilon_i \sim N(0, 0.5^2)$ — a constant true effect of 1 sitting on a convex baseline.

```
set.seed(7)
n <- 150
x <- c(rnorm(n, 2.5, 1.1), rnorm(n, 4.5, 1.1))
d <- c(rep(0, n), rep(1, n))
y <- 1.0 * d + 0.8 * (x - 3.5)^2 + 0.3 * x + rnorm(2 * n, 0, 0.5)
df <- data.frame(y, d, x)
X <- cbind(1, x); dh <- as.vector(X %*% solve(crossprod(X)) %*% t(X) %*% d)
flag <- (d == 1 & dh > 1) | (d == 0 & dh < 0)
pch <- ifelse(d == 1, 17, 16); col <- ifelse(d == 1, teal, rust)
plot(x, y, type = "n", xlab = "x", ylab = "y")
xs <- seq(min(x), max(x), length = 200)
mc <- function(z) 0.8 * (z - 3.5)^2 + 0.3 * z              # true control mean
 ↪  E[y | x, D = 0]
lines(xs, mc(xs),     col = rust, lwd = 2.4)               # control mean
lines(xs, mc(xs) + 1, col = teal, lwd = 2.4, lty = 3)      # treated mean =
 ↪  control + true effect (1)
points(x, y, pch = pch, col = paste0(col, "88"), cex = 0.85)               # all
 ↪  points, by group
points(x[flag], y[flag], pch = 1, col = "grey15", cex = 1.7, lwd = 1.4)    # ring
 ↪  the flagged units
arrows(3.5, mc(3.5), 3.5, mc(3.5) + 1, code = 3, length = 0.05, angle = 90, lwd =
 ↪  2.4, col = "grey10")
segments(3.5, mc(3.5) + 0.5, 2.05, 13.3, col = "grey55", lwd = 1)
text(2.0, 14, "gap = true effect = 1", pos = 4, cex = 0.82, col = "grey10")
legend("bottomright", bty = "n", cex = 0.82,
```

```
       legend = c("control", "treated", "flagged (trimmed)", "control mean",
        ↪  "treated mean"),
       pch = c(16, 17, 1, NA, NA), lty = c(NA, NA, NA, 1, 3),
       col = c(rust, teal, "grey15", rust, teal))
```




Figure 14.2: Thin overlap on a convex baseline. The two curves are the true means — control (rust, solid) and treated (teal, dotted), a constant 1 apart (arrow). Controls sit low on the curve and treated high; the groups share only the middle. Ringed points have wrong-sign outcome weights: treated units with $\hat{D}_i > 1$ or controls with $\hat{D}_i < 0$. The straight $y \sim D + x$ reports $\hat{\beta} = 0.69$; dropping the flagged units, or reweighting to the overlap, restores ≈ 1.

```
w_ato <- WeightIt::weightit(d ~ x, data = df, method = "glm", estimand = "ATO")
```

```
vtab(c(`flagged units`      = sum(flag),
       `$\\hat\\beta$ (full)`        = coef(lm(y ~ d + x))[["d"]],
       `$\\hat\\beta$ (trimmed)`     = coef(lm(y ~ d + x, data = df[!flag,
        ↪  ]))[["d"]],
       `$\\hat\\beta$ (overlap-wtd)` = coef(lm(y ~ d + x, weights =
        ↪  w_ato$weights))[["d"]]), 3)
```

	value
flagged units	37.000
$\hat{\beta}$ (full)	0.688
$\hat{\beta}$ (trimmed)	0.991
$\hat{\beta}$ (overlap-wtd)	0.979

Thirty-seven units carry the wrong sign, all in the non-overlapping tails. Trim them and $\hat{\beta}$ jumps from 0.69 to 0.99; the overlap weights (`"ATO"`, Li, Morgan, and Zaslavsky 2018) give 0.98. Either fix confirms the estimate leaned on the extrapolation, read off the regression itself with no separate model. Had the outcome been linear, the same trim would barely move $\hat{\beta}$: the wrong-sign weights are still there, but they are harmless. The check tells you which case you are in.

This is the implied-weights diagnostic of Chattopadhyay and Zubizarreta (2023); their `lmw` package computes the weights and the balance, extrapolation, and influence diagnostics built on them, and `OutcomeWeights` (Knaus 2024) carries it to machine-learning estimators. The mainstream overlap check takes the other route, through the propensity score (`WeightIt` for estimation, `cobalt` for balance, and Crump et al. 2009 for trimming); both routes reach the same conclusion.

14.4 Out of OLS: the same weights beyond the linear model

The outcome reading has one advantage the effect reading lacks: it survives leaving OLS. Any estimator that can be written $\hat{\tau} = \sum_i \omega_i Y_i = \omega' Y$ — a linear combination of the observed outcomes — *has* outcome weights, whatever machinery produced them.

Knaus (2024) makes this the organising idea, and his `OutcomeWeights` package computes ω for AIPW, double machine learning, and the causal forest.

14.4.1 The weights, derived

The OLS outcome weights (Equation 14.2) followed from one fact: in the FWL numerator $\sum_i \tilde{V}_i y_i$, the outcome y_i appears once, linearly, so $\hat{\beta} = \sum_i \omega_i y_i$ with $\omega_i = \tilde{V}_i / \sum_j \tilde{V}_j^2$. The same fact holds whenever the outcome model is a *smoother* — a map $\hat{m} = SY$ that is linear in the training outcomes. Substituting SY for $\hat{m}$ keeps Y linear in the numerator, and reading off its coefficient gives ω. We carry this out for PLR and AIPW.

PLR. Take the partially-linear estimator (the R-learner behind the causal forest). It is FWL with the projections done by a flexible learner: residualise treatment and outcome on the covariates, then regress one residual on the other,

$$\hat{\tau} = \frac{\sum_i (D_i - \hat{e}_i)(Y_i - \hat{m}_i)}{\sum_i (D_i - \hat{e}_i)^2}, \tag{14.6}$$

with $\hat{e} = \hat{E}[D \mid X]$ and $\hat{m} = \hat{E}[Y \mid X]$ the fitted nuisances.

The outcome nuisance is a *smoother*: its fitted values are a linear map of the training outcomes, $\hat{m} = SY$ — true for OLS, ridge, kernels, and honest forests alike. The word *honest* carries weight there. If the learner's structure is itself chosen using the outcomes — splits grown on the same Y being predicted, a lasso whose active set is outcome-selected, boosting — then S depends on Y and the map stops being linear in it. What follows then holds *conditional on the fitted learner*, which is the weaker claim: the weights describe the estimate you got rather than a globally linear estimator. Honesty and cross-fitting narrow the problem without removing it. They stop an observation from training the learner that predicts it, which is what the orthogonality argument needs, but each fold's structure is still grown on the outcomes in the other folds, so S remains outcome-dependent overall. The representation below is exact *conditional on the fitted learners*; it is a globally linear reading of the estimator only for learners whose structure does not depend on their training outcomes. So the outcome residual is $Y - \hat{m} = (\mathbf{I} - S)Y$. Writing $\tilde{D} = D - \hat{e}$ for the treatment residual and substituting,

$$\hat{\tau} = \frac{\tilde{D}'(\mathbf{I} - S)\,Y}{\tilde{D}'\tilde{D}}. \tag{14.7}$$

Now Y appears once, linearly, in the numerator. Read off its coefficient and the weights are done:

$$\hat{\tau} = \omega' Y, \qquad \omega' = \frac{\tilde{D}'(\mathbf{I} - S)}{\tilde{D}'\tilde{D}}. \tag{14.8}$$

That is Equation 14.2 — the OLS weights — with the projection annihilator M_X replaced by the smoother annihilator $\mathbf{I} - S$. OLS is the case of a linear learner, $S = P_X$: there S is a projection, $\mathbf{I} - S = M_X$, and Equation 14.8 is Equation 14.2 exactly. Component by component, the weight on observation i is

$$\omega_i = \frac{\tilde{D}_i - \sum_j S_{ji}\,\tilde{D}_j}{\sum_k \tilde{D}_k^2}\,: \tag{14.9}$$

the treatment residual of unit i, minus how much other units' treatment residuals leak to i through the outcome smoother, normalised by the total treatment-residual variance.

The outcome weights tell which rows drive the estimate; the *effect* weights tell whose effect it estimates. Write the outcome as $Y_i = \tau(X_i)D_i + g(X_i) + \varepsilon_i$, with $g(X_i) = E[Y_i(0) \mid X_i]$ the untreated baseline and $E[\varepsilon_i \mid D_i, X_i] = 0$. The PLR nuisance $\hat{m}$ does not estimate g. It estimates the marginal mean $m(X_i) = E[Y_i \mid X_i] = g(X_i) + e(X_i)\tau(X_i)$, so with consistent nuisances $Y_i - m(X_i) = \tau(X_i)\tilde{D}_i + \varepsilon_i$, and the numerator reduces in conditional expectation to $\sum_i \tilde{D}_i^2 \tau(X_i)$. The effect weight on $\tau(X_i)$ is $\tilde{D}_i^2$. Equivalently it is $\tilde{D}_i D_i = (D_i - e_i)D_i$, since $\tilde{D}_i^2 = \tilde{D}_i D_i - \tilde{D}_i e_i$ and the last term has conditional mean zero. Since D is binary ($D_i^2 = D_i$),

$$E[\tilde{D}_i D_i \mid X_i] = E[D_i \mid X_i] - e_i\,E[D_i \mid X_i] = e_i - e_i^2 = e_i(1 - e_i). \tag{14.10}$$

The denominator $\tilde{D}'\tilde{D} = \sum_i \tilde{D}_i^2$ has the same expectation, $\sum_i e_i(1 - e_i)$, since $\mathrm{Var}(D_i \mid X_i) = e_i(1 - e_i)$. Both numerator and denominator weight by propensity variance, and the ratio is the variance-weighted average treatment effect,

$$\tau_{\text{VWATE}} = \frac{\sum_i e(X_i)(1-e(X_i))\,\tau(X_i)}{\sum_i e(X_i)(1-e(X_i))}, \tag{14.11}$$

not the plain ATE $\bar{\tau} = n^{-1}\sum_i \tau_i$. The two coincide when effects are constant, or when treatment is randomly assigned so that $e(1-e)$ is the same everywhere. Neither condition is necessary. All that is required is that $\tau(X)$ be uncorrelated with the weight $e(X)(1-e(X))$: heterogeneity that happens not to line up with propensity variance leaves the two equal. The weighting is the same $e(1-e)$ that appeared in the previous chapter's Equation 13.8, where Angrist (1998) showed OLS with saturated strata weights each stratum by $n_x\hat{D}(x)(1-\hat{D}(x))$. That result required discrete strata and saturated controls; Equation 14.11 is the same propensity-variance weighting with continuous covariates and a flexible learner. The individual $\tau(X_i)$ are never observed or estimated — they are the structural effects in $Y_i = \tau(X_i)D_i + g(X_i) + \varepsilon_i$. PLR returns one number, $\hat{\tau}$, that converges to their $e(1-e)$-weighted average; to estimate $\tau(X_i)$ itself one needs a CATE estimator (a causal forest, say).

Three moves did it: write the estimator, substitute the smoother $\hat{m} = SY$, read off the coefficient on Y. They work for any estimator built from smoothers; only the operator on Y changes. AIPW is worth carrying through, because its operator is more than $\mathbf{I} - S$.

AIPW. With outcome smoothers $\hat{\mu}_1 = S_1 Y$, $\hat{\mu}_0 = S_0 Y$ and propensity $\hat{e}$, the score of unit i is

$$\psi_i = \hat{\mu}_{1i} - \hat{\mu}_{0i} + \frac{D_i}{\hat{e}_i}(Y_i - \hat{\mu}_{1i}) - \frac{1-D_i}{1-\hat{e}_i}(Y_i - \hat{\mu}_{0i}), \qquad \hat{\tau} = \frac{1}{n}\sum_i \psi_i. \tag{14.12}$$

Each $\hat{\mu}_{ji}$ is $(S_j Y)_i$, so Y again enters only linearly; collecting its coefficient gives $\hat{\tau} = \omega' Y$ with

$$T = S_1 - S_0 + \operatorname{diag}\!\left(\tfrac{D}{\hat{e}}\right)(\mathbf{I} - S_1) - \operatorname{diag}\!\left(\tfrac{1-D}{1-\hat{e}}\right)(\mathbf{I} - S_0), \qquad \omega' = \frac{\mathbf{1}'T}{n}. \tag{14.13}$$

That is, ω_j is the j-th column-sum of T divided by n: observation j's outcome enters the ATE through its direct IPW term and through its indirect contribution via the two outcome smoothers S_1, S_0 to every other unit's augmentation correction.

Comparing the two:

	PLR (Equation 14.8)	AIPW
Matrix form	$\omega' = \tilde{D}'(\mathbf{I} - S) \,/\, \tilde{D}'\tilde{D}$	$\omega' = \mathbf{1}'T/n$
Per unit	$\omega_i = (\tilde{D}_i - \sum_j S_{ji}\tilde{D}_j) \,/\, \sum_k \tilde{D}_k^2$	$\omega_j = \frac{1}{n}\sum_i T_{ij}$
Smoother	one S (outcome on X)	two: S_1 (treated), S_0 (control)
Normalisation	treatment-residual variance	sample size n

Both follow the same recipe — substitute the smoother $\hat{m} = SY$, read off Y's coefficient — and differ only in the operator acting on Y.

Simulation — heterogeneous effects (DML). Under a heterogeneous effect the two estimators diverge, and the weights show why. We simulate $n = 2000$ units: five standard-normal covariates X, selection $W_i \sim \text{Bernoulli}(\Lambda(-1 + 1.2\,X_{i1}))$ with Λ the logistic function (treatment is scarce and tied

to X_1), and outcome $Y_i = \tau_i W_i + X_{i1} + 0.5\,X_{i2} + \varepsilon_i$ with $\varepsilon_i \sim N(0,1)$ and an effect $\tau_i = 1 + X_{i1}$ that varies across units — a $W \times X_1$ interaction. The average effect is $E[\tau] = 1$, but units with larger X_1 have larger effects. `dml_with_smoother` returns a partially-linear (PLR) and an AIPW estimator; `get_outcome_weights` extracts the weight matrix ω, one row per estimator.

```
library(OutcomeWeights)
set.seed(1)
n <- 2000; p <- 5
X <- matrix(rnorm(n * p), n, p)
etrue <- plogis(-1 + 1.2 * X[, 1])                          # asymmetric selection
 ↪  on X1
W <- rbinom(n, 1, etrue)
tau <- 1 + X[, 1]                                           # heterogeneous effect:
 ↪  W x X1
Y <- tau * W + X[, 1] + 0.5 * X[, 2] + rnorm(n)
dml <- dml_with_smoother(Y, W, X, estimators = c("PLR", "AIPW_ATE"))
om  <- get_outcome_weights(dml)                             # om$omega: 2 x n,
 ↪  om$treat = W
invisible(capture.output(s <- summary(dml)))
W1 <- om$treat == 1
tab <- t(sapply(rownames(om$omega), function(nm) {
  w <- om$omega[nm, ]
  c(estimate = s[nm, 1], `$\\boldsymbol\\omega'\\boldsymbol Y$` = sum(w * Y),
    treated = sum(w[W1]), control = sum(w[!W1]))
}))
ov <- etrue * (1 - etrue)                                   # e(1-e): PLR's per-unit
 ↪  weight
# ATE = plain mean of tau; overlap-ATE = e(1-e)-weighted mean of tau (PLR's
 ↪  target)
cat(sprintf("true ATE = %.2f   true overlap-ATE = %.2f\n", mean(tau), sum(ov *
 ↪  tau) / sum(ov)))
```

```
true ATE = 0.99   true overlap-ATE = 1.31
```

```
mtab(tab, 3)
```

	estimate	$\omega'Y$	treated	control
PLR	1.287	1.287	0.997	-0.997
AIPW-ATE	1.074	1.074	1.000	-1.000

Two things stand out. First, the estimators now *disagree* — PLR reports 1.29, AIPW 1.07 — and neither is wrong; they target different populations. PLR targets the VWATE (Equation 14.11), not the ATE. Here treatment selection loads on X_1 through $e = \Lambda(-1 + 1.2\,X_1)$, and $\tau = 1 + X_1$ also rises in X_1, so the units PLR overweights — those with e near $\frac{1}{2}$, around $X_1 \approx 0.8$ — carry

above-average effects, pushing PLR's estimate above the true ATE. AIPW instead reweights by $1/e$ and $1/(1-e)$ to undo selection, returning the plain average, the ATE (≈ 1). Same outcomes Y, weighted two ways, two population-averages of τ — and the weights are what make the difference visible.

Second, the weights still reproduce each estimate exactly and sum to zero overall. They part on the other constraint from Equation 14.2, $\sum_i \omega_i V_i = 1$: for a binary treatment that constraint is the treated weights summing to $+1$ (and, with the zero total, controls to -1). AIPW keeps it exactly; the partially-linear estimator sums to ± 0.997. That shortfall has a direct reading. If the treated weights sum to a and the controls to $-a$, then $\hat{\tau} = a\,(\bar{y}^{\omega}_{\text{treated}} - \bar{y}^{\omega}_{\text{control}})$, so the estimate is the weighted contrast scaled by a: at $a = 0.997$ the two differ by 0.3%. Whether that is an understatement depends on treating the normalized contrast as the target, which nothing here establishes — better read as a normalization discrepancy. The two estimators reach their sums by different routes. For PLR the treated weights sum to $\omega' D = \tilde{D}'(\mathbf{I} - S)D / \tilde{D}'\tilde{D}$, by Equation 14.8. This is one whenever the outcome smoother applied to D reproduces the propensity fit, $SD = \hat{e}$, because then $(\mathbf{I} - S)D = \tilde{D}$; idempotence does not matter. OLS is that case, since both residualizations use P_X. Here the outcome and the treatment are fit by separate forests, so $SD \neq \hat{e}$, and PLR's sum sits slightly off. AIPW needs less. Its treated weights are $D_i/(n\hat{e}_i)$ plus the weight each treated outcome receives through the treated-arm fit $\hat{\mu}_1$. That smoother is fit on treated units only, so each of its rows has entries only in the treated columns. If every row sums to one over those columns — the fit reproduces constants — the second part sums to $1 - \frac{1}{n}\sum_j D_j/\hat{e}_j$ and cancels the first part's excess, so the treated weights sum to exactly 1, idempotent or not. The same holds for the controls. An outcome learner that does not reproduce constants moves the AIPW sums off ± 1 as well.

That divergence shows what the representation is for. In Knaus's (2024) framing ω does three things: it puts OLS, IPW, AIPW, DML, and the causal forest on **one footing** ($\omega' Y$); it **characterises the target population** each reports — the PLR-versus-AIPW gap above; and it **checks covariate balance** via `cobalt`, flagging when weights break a property usually assumed — the sum-to-± 1 within treated and control groups that PLR violates here.

The formulas take a few lines to check against the package:

```
n  <- length(Y)
S  <- dml$NuPa.hat$smoothers$S[1, , ]
S1 <- dml$NuPa.hat$smoothers$S.d1[1, , ]
S0 <- dml$NuPa.hat$smoothers$S.d0[1, , ]
e  <- dml$NuPa.hat$predictions$D.hat[, 1]
Wtil   <- W - e
om_plr <- as.numeric((Wtil - crossprod(S, Wtil)) / sum(Wtil^2))
Imat    <- diag(n)
Tm      <- S1 - S0 + (W / e) * (Imat - S1) - ((1 - W) / (1 - e)) * (Imat - S0)
om_aipw <- colSums(Tm) / n
vtab(c(`PLR: largest gap vs package` = max(abs(om_plr  - om$omega["PLR", ])),
       `AIPW: largest gap vs package` = max(abs(om_aipw - om$omega["AIPW-ATE",
        ↪  ])),
       `PLR: $\\boldsymbol\\omega'\\boldsymbol Y$`                = sum(om_plr
        ↪  * Y),
       `AIPW: $\\boldsymbol\\omega'\\boldsymbol Y$`                = sum(om_aipw
        ↪  * Y)), 4)
```

	value
PLR: largest gap vs package	0.0000
AIPW: largest gap vs package	0.0000
PLR: $\omega'Y$	1.2873
AIPW: $\omega'Y$	1.0738

14.4.2 Two uses: balance and extrapolation

Having ω, two checks run directly on it. Does the estimator's implied weighting balance the covariates? Hand ω to `cobalt` and read the largest standardised mean difference (SMD), the between-group difference in a covariate's weighted mean divided by its standard deviation. And is it extrapolating? Count the wrong-sign weights — a control with $\omega_i > 0$ or a treated unit with $\omega_i < 0$, a sign that the fitted comparison extrapolates in this sample.

```
library(cobalt)
uses <- t(sapply(rownames(om$omega), function(nm) {
  w <- om$omega[nm, ]
  c(`$\\max\\lvert\\mathrm{SMD}\\rvert$`         =
   ↪  max(abs(bal.tab(as.data.frame(X), treat = W, weights =
   ↪  w)$Balance$Diff.Adj)),
    `wrong-sign units` = sum((om$treat == 1 & w < 0) | (om$treat == 0 & w > 0)))
}))
mtab(uses, 3)
```

	max|SMD|	wrong-sign units
PLR	0.025	21
AIPW-ATE	0.091	0

Raw, the groups are badly imbalanced — the largest covariate SMD is 1.11, since treatment was tied to X_1. Both weightings pull it down to near zero (0.03 for PLR, 0.09 for AIPW): each balances the covariates, which for a causal forest you could not otherwise check.

They differ on extrapolation. PLR carries 21 wrong-sign controls — it leans slightly off common support — while AIPW carries none. Neither is disqualifying here; the point is that balance and extrapolation are now *checkable*, on estimators that never exposed a weighting.

14.5 The DiD setting

When the regressor is a treatment indicator, every unit in a group gets the same weight, so $\hat{\beta}$ reduces to a contrast of group means. Regressed on the treatment alone it is the difference in means

between treated and control. Add a time period and its interaction with treatment and it becomes a difference-in-differences across four group-by-period cells. We take these in turn, then bring in covariates.

Simulation — treatment alone. Regress y on the treatment D alone — here 30 treated drawn $y \sim N(5,1)$ and 50 controls $y \sim N(2,1)$. The residual $\tilde{V}_i = D_i - \bar{D}$ is constant within each group, so the weight is uniform within a group and $\hat{\beta}$ reduces to the two group means: $\hat{\beta} = \bar{y}_1 - \bar{y}_0$, the difference in means. Every treated unit then carries positive weight, every control negative — the structural pattern of the previous section, with the wrong-sign off-diagonal empty until covariates enter.

```
set.seed(7)
n1 <- 30; n0 <- 50
d <- c(rep(1, n1), rep(0, n0)); y <- c(rnorm(n1, 5), rnorm(n0, 2))
dev <- d - mean(d); om <- dev / sum(dev^2); col <- ifelse(om >= 0, teal, rust)
y1 <- mean(y[d == 1]); y0 <- mean(y[d == 0])
plot(jitter(d, 0.4), y, pch = 16, col = paste0(col, "55"), cex = 0.9,
     xlim = c(-0.3, 1.3), xaxt = "n", xlab = "", ylab = "y", main = "Regression
      ↪  on treatment")
axis(1, at = c(0, 1), labels = c("control", "treated"))
segments(0, y0, 1, y1, lwd = 2.6, col = "grey15")
points(c(0, 1), c(y0, y1), pch = 21, bg = c(rust, teal), col = "white", cex =
 ↪  2.5, lwd = 2)
```




Figure 14.3: Regression on the treatment alone. Points are the two groups (rust control, teal treated); the large dots are the group means, and the segment joining them has slope $\hat{\beta} = \bar{y}_1 - \bar{y}_0$, the difference in means.

```
vtab(c(`$\\hat\\beta$ (lm)` = coef(lm(y ~ d))[["d"]], `$\\bar y_1 - \\bar y_0$`
 ↪  = y1 - y0,
        `off-diagonal weights` = sum((d == 1 & om < 0) | (d == 0 & om > 0))), 3)
```

	value
$\hat{\beta}$ (lm)	3.318
$\bar{y}_1 - \bar{y}_0$	3.318
off-diagonal weights	0.000

The estimate equals $\bar{y}_1 - \bar{y}_0$, and no weight carries the wrong sign — the off-diagonal count is zero.

Simulation — 2×2 difference-in-differences. Add a period and interact it with the group, and the design has four cells. Each collapses to its mean, and the interaction coefficient is their double contrast,

$$\widehat{\text{DiD}} = (\bar{y}_{T,\text{post}} - \bar{y}_{T,\text{pre}}) - (\bar{y}_{C,\text{post}} - \bar{y}_{C,\text{pre}}). \tag{14.14}$$

Now there are four cell means, and with them four slopes — the two group trends over time and the two cross-sectional gaps. The DiD is the difference between either parallel pair, trend minus trend

or gap minus gap. We simulate 200 units in each group×period cell from $y = 3 + 2g + p + 2\,(g \cdot p) + \varepsilon$ with $\varepsilon \sim N(0, 1)$, so the true DiD (the interaction) is 2:

```
set.seed(3)
g <- rep(c(1, 1, 0, 0), each = 200); p <- rep(c(0, 1, 0, 1), each = 200)
y <- 3 + 2 * g + 1 * p + 2 * (g * p) + rnorm(800)
m <- tapply(y, list(g, p), mean)
Tpre <- m["1", "0"]; Tpost <- m["1", "1"]; Cpre <- m["0", "0"]; Cpost <- m["0",
 ↪  "1"]
cf <- Tpre + (Cpost - Cpre)
plot(NA, xlim = c(-0.15, 1.25), ylim = range(y), xaxt = "n", xlab = "", ylab =
 ↪  "y",
     main = "Difference-in-differences")
axis(1, at = c(0, 1), labels = c("pre", "post"))
segments(0, Tpre, 1, Tpost, lwd = 2.4, col = teal)
segments(0, Cpre, 1, Cpost, lwd = 2.4, col = rust)
segments(0, Tpre, 1, cf, lty = 2, lwd = 1.6, col = "grey55")
segments(1, cf, 1, Tpost, lwd = 4, col = "grey15")
points(c(0, 1, 0, 1), c(Tpre, Tpost, Cpre, Cpost),
       pch = 21, bg = c(teal, teal, rust, rust), col = "white", cex = 2.3, lwd =
        ↪  2)
text(1.02, (cf + Tpost) / 2, "DiD", pos = 4, cex = 0.9, col = "grey15")
```




Figure 14.4: The 2×2 difference-in-differences: four cell means (dots), the treated (teal) and control (rust) trends, the parallel-trends counterfactual (dashed), and the DiD (black bar) as the gap between the treated post-mean and that counterfactual.

```
vtab(c(`treated trend` = Tpost - Tpre, `control trend` = Cpost - Cpre,
       `pre gap` = Tpre - Cpre, `post gap` = Tpost - Cpost,
       `DiD (interaction)` = coef(lm(y ~ g * p))[["g:p"]]), 3)
```

	value
treated trend	3.029
control trend	0.897
pre gap	2.019
post gap	4.151
DiD (interaction)	2.132

The estimated interaction recovers the true DiD of 2, and the four cell-mean weights carry the structural signs $(+, -, -, +)$ as long as every cell is populated.

The clean 2×2 has trivial weights — four cell means, each ± 1. Real difference-in-differences carries covariates: parallel trends holds only *conditional* on X, and the comparison has to be modelled. That model is a weighting of outcomes too.

The doubly-robust DiD of Sant'Anna and Zhao (2020) is AIPW read on the outcome *change* $\Delta Y = Y_1 - Y_0$: a propensity for treatment, a control-group regression of ΔY on X, combined. By

the recipe of the last section its estimate is $\widehat{\text{ATT}} = \omega'\Delta Y$, with $\omega' = a'(\mathbf{I} - S)$ — the control-trend smoother S and the group weights $a = w_{\text{treat}} - w_{\text{cont}}$.

Simulation — doubly-robust DiD. Two periods and $n = 2000$, with covariates $x_1, x_2 \sim N(0,1)$. Treatment is selected on x_1 with thin overlap, $\Pr(D = 1) = \Lambda(-0.5 + 2x_1)$. The outcomes are $y_{i0} = 1 + x_{1i} + 0.5\,x_{2i} + \varepsilon_{i0}$ and $y_{i1} = y_{i0} + (0.5 + 0.4\,x_{1i}) + D_i + \varepsilon_{i1}$, with both errors $N(0,1)$. The control trend $0.5 + 0.4x_1$ depends on x_1, so trends are parallel only given X, and the true ATT is 1. We fit DR-DiD with `DRDID`, rebuild its weights by hand, and check.

```
library(DRDID)
set.seed(2)
n  <- 2000
x1 <- rnorm(n); x2 <- rnorm(n); Xcov <- cbind(1, x1, x2)
D  <- rbinom(n, 1, plogis(-0.5 + 2 * x1))        # selection into treatment on x1
 ↪  (thin overlap)
y0 <- 1 + x1 + 0.5 * x2 + rnorm(n)               # pre-period outcome
y1 <- y0 + (0.5 + 0.4 * x1) + 1 * D + rnorm(n)   # control trend depends on x1;
 ↪  true ATT = 1
dY <- y1 - y0
att <- drdid_panel(y1 = y1, y0 = y0, D = D, covariates = Xcov, boot = FALSE)$ATT

# rebuild the DR-DiD weights: omega = (I - S)' a, acting on the outcome change dY
e    <- pmin(as.vector(glm(D ~ x1 + x2, family = binomial)$fitted), 1 - 1e-6)
keep <- rep(TRUE, n); keep[D == 0] <- e[D == 0] < 0.995     # DRDID's control
 ↪  trimming
Xc   <- Xcov[D == 0, ]
S    <- matrix(0, n, n); S[, D == 0] <- Xcov %*% solve(crossprod(Xc)) %*% t(Xc)
w.t  <- keep * D
w.c  <- keep * e * (1 - D) / (1 - e)
a    <- w.t / sum(w.t) - w.c / sum(w.c)          # group weights, treated minus
 ↪  reweighted control
omega <- as.vector(a - crossprod(S, a))

Xk <- cbind(x1, x2); sds <- apply(Xk, 2, sd)
vtab(c(`DRDID ATT`                = att,
       `$\\boldsymbol\\omega'\\Delta\\boldsymbol Y$`               = sum(omega
        ↪  * dY),
       `wrong-sign weights`       = sum((D == 1 & omega < 0) | (D == 0 & omega >
        ↪  0)),
       `covariate imbalance, raw` = max(abs((colMeans(Xk[D == 1, ]) -
        ↪  colMeans(Xk[D == 0, ])) / sds)),
       `covariate imbalance, preliminary $\\boldsymbol a$` = max(abs(as.vector(a
        ↪  %*% Xk) / sds)),
       `covariate imbalance, augmented $\\boldsymbol\\omega$` =
        ↪  max(abs(as.vector(omega %*% Xk) / sds))), 3)
```

	value
DRDID ATT	0.980
$\omega'\Delta Y$	0.980
wrong-sign weights	151.000
covariate imbalance, raw	1.198
covariate imbalance, preliminary a	0.086
covariate imbalance, augmented ω	0.000

$\omega'\Delta Y$ reproduces the `DRDID` estimate exactly — the difference-in-differences is a signed weighting of observed outcome *changes*, and the diagnostics of this chapter apply to it. Balance has two stages. The preliminary weights a already shrink the largest standardised imbalance from 1.20 to 0.09. The augmented weights ω balance both covariates exactly. The control outcome model is linear in $(1, x_1, x_2)$, so S reproduces those columns, $SX = X$, and $\omega'X = a'X - a'SX = 0$. Exact balance here is a property of the outcome model, not evidence about overlap. And 151 units carry a wrong-sign weight, all of them controls. A treated unit cannot: the outcome model is fit on controls only, so S has no treated columns and every treated weight stays at $a_i > 0$. The flagged controls are ones with small reweighting weight a_i that the adjustment $-(S'a)_i$ pushes above zero. As in the binary-regression case above, that is evidence the fitted comparison extrapolates in this sample, not proof that population support fails.

14.5.1 Staggered designs: TWFE at the observation level

Negative weights in staggered designs are usually described at the level of two-by-two comparisons, and the description needs care, because two different decompositions get run together. Goodman-Bacon (2021) writes TWFE as a weighted average of all two-by-two DiD comparisons, and those comparison weights are *non-negative*: they are built from group shares and treatment-variance terms and they sum to one. What goes wrong there is the estimand, not the weight. A comparison that uses an already-treated cohort as its control subtracts that cohort's own treatment effect whenever effects move over time, so what it contributes is not an ATT. That is the forbidden comparison. De Chaisemartin and D'Haultfœuille (2020) decompose TWFE differently, into a weighted sum of *treatment-effect* parameters, and there some weights genuinely are negative. Both are effect-weight readings. The outcome-weight reading goes one level further down, to the individual y_{it}.

TWFE is still FWL: residualise D_{it} on unit and time fixed effects to get $\tilde{D}_{it}$, and $\hat{\beta} = \sum_{it} \omega_{it} y_{it}$ with $\omega_{it} = \tilde{D}_{it} / \sum_{jt} \tilde{D}^2_{jt}$. A treated observation with $\omega_{it} < 0$ is a cell whose outcome is subtracted from the estimated effect — used as a control even though it is treated. It is the same sign flip seen earlier under thin overlap, caused here by the unit-and-time residualisation pushing $\tilde{D}_{it}$ past zero in late periods of early cohorts, where the fixed-effect prediction of D_{it} exceeds 1. Untreated cells need more care, because a difference-in-differences gives them both signs by design. It adds the control group's pre-period baseline and subtracts its post-period outcomes, the $(+, -, -, +)$ pattern of the 2×2 case. So an untreated cell with $\omega_{it} > 0$ is usually a control baseline, not a wrong sign. In a panel the cross-sectional rule, untreated cells should be negative, does not apply.

Simulation — staggered panel. $N = 300$ units over $T = 8$ periods, in three cohorts: never-treated (20%), early-treated at period 3 with effect $\tau = 3$ (40%), and late-treated at period 6 with effect $\tau = 1$ (40%). Outcomes are $y_{it} = \alpha_i + \gamma_t + \tau_g D_{it} + \varepsilon_{it}$, with unit effects $\alpha_i \sim N(0, 1)$, a

time effect γ_t rising linearly from 0 to 1.4, and $\varepsilon_{it} \sim N(0, 0.5^2)$. The high treated share makes the forbidden comparison visible.

```
library(fixest)
set.seed(5)
N <- 300; TT <- 8
g <- sample(c(0, 3, 6), N, TRUE, prob = c(.2, .4, .4))
panel <- data.frame(id = rep(1:N, each = TT), t = rep(1:TT, N))
panel$g <- g[panel$id]
panel$D <- as.integer(panel$g > 0 & panel$t >= panel$g)
tau <- ifelse(panel$g == 3, 3, ifelse(panel$g == 6, 1, 0))
alpha <- rnorm(N); gamma <- seq(0, 1.4, length.out = TT)
panel$y <- alpha[panel$id] + gamma[panel$t] + tau * panel$D + rnorm(nrow(panel),
 ↪  0, 0.5)

Dtil <- resid(feols(D ~ 1 | id + t, data = panel))
omega_tw <- Dtil / sum(Dtil^2)
n1 <- sum(panel$D)
vtab(c(`true ATT`              = mean(tau[panel$D == 1]),
       `TWFE`                  = coef(feols(y ~ D | id + t, panel))[["D"]],
       `$\\boldsymbol\\omega'\\boldsymbol y$`              = sum(omega_tw *
        ↪  panel$y),
       `treated, $\\omega_{it} < 0$`    = sum(panel$D == 1 & omega_tw < 0),
       `untreated, $\\omega_{it} > 0$`  = sum(panel$D == 0 & omega_tw > 0),
       `  never-treated, periods 1-5` = sum(panel$g == 0 & panel$t <= 5 &
        ↪  omega_tw > 0),
       `  late cohort, periods 1-2`   = sum(panel$g == 6 & panel$t <= 2 &
        ↪  omega_tw > 0)), 3)
```

	value
true ATT	2.426
TWFE	1.915
$\omega' y$	1.915
treated, $\omega_{it} < 0$	384.000
untreated, $\omega_{it} > 0$	551.000
never-treated, periods 1-5	345.000
late cohort, periods 1-2	206.000

The TWFE coefficient sits well below the true ATT, and $\sum \omega_{it} y_{it}$ reproduces it exactly. Hundreds of treated cells carry a negative weight: these are the early cohort's cells in periods 6–8, the ones where the unit-and-time prediction of D exceeds 1 and the residual $\tilde{D}_{it}$ flips sign. Their outcomes are subtracted from $\hat{\beta}$ — the observation-level signature of the Bacon forbidden comparison. The untreated cells with positive weight are all control baselines: never-treated units in periods 1–5 and the late cohort in periods 1–2, before either serves as a post-period control. None of them is a wrong sign.

This simulation shows the negative weights but not their cost. Each cohort's effect is constant over time, so in the comparison of the late cohort against the early one, the early cohort's effect enters both periods and cancels. The gap between TWFE and the true ATT comes from variance weighting instead: the late cohort is treated in 3 of 8 periods, the early one in 6, so the late cohort's treatment varies more within units and its smaller effect gets more weight. With constant effects, TWFE is a non-negatively weighted average of the cohort effects (Goodman-Bacon 2021). If the early cohort's effect grew with time since adoption, the forbidden comparison would subtract that growth and add bias on top.

The wrong-sign diagnostic in a panel is by observation (i, t), not by unit. In the cross-sectional case earlier, the fix was to trim wrong-sign observations and refit. In a panel, trimming specific (i, t) cells removes observations that anchor that unit's fixed effect and distorts the estimates for its remaining cells. The response in the staggered setting is to switch estimators — imputation (Borusyak, Jaravel, and Spiess 2024), Callaway and Sant'Anna (2021), or Sun and Abraham (2021) — that avoid the forbidden comparison by construction. The imputation estimator, for example, fits unit and time fixed effects on untreated observations alone and imputes each treated cell's counterfactual; because its smoother draws only on untreated outcomes, the weight on every treated cell reduces to $1/n_1$ and no treated observation can count as a control.

14.5.2 Synthetic difference-in-differences: a rank-one weighting

Synthetic difference-in-differences is a weighted two-way fixed effects regression, from Arkhangelsky et al. (2021). The estimators so far imply their weights. This one chooses them:

$$(\hat{\tau}, \hat{\mu}, \hat{\alpha}, \hat{\beta}) = \arg\min \sum_i \sum_t \left(Y_{it} - \mu - \alpha_i - \beta_t - W_{it}\tau\right)^2 \omega_i \lambda_t. \tag{14.15}$$

The recipe: fit ω_i on the control units so the reweighted control path tracks the treated path over the pre-period, and fit λ_t on the pre-periods so the reweighted pre-period resembles the post-period. Treated units enter at $1/N_1$, post-periods at $1/T_1$. Both fitted sets lie on the simplex, non-negative and summing to one.

Equation 14.15 is weighted least squares, so the FWL reading of Equation 14.2 goes through with the weights carried along. Then $\hat{\tau}$ is again a weighting of outcomes. Let u_i be $1/N_1$ for a treated unit and $-\omega_i$ for a control. Let v_t be $1/T_1$ for a post-period and $-\lambda_t$ for a pre-period. Then

$$\hat{\tau} = \sum_i \sum_t \omega_{it}\, Y_{it}, \qquad \omega_{it} = u_i\, v_t. \tag{14.16}$$

In words, the weight on a cell is a unit part times a time part. Collect the weights into an $N \times T$ matrix Ω, one entry per cell. Then Equation 14.16 reads $\Omega = uv'$, an outer product, and a matrix of that form has **rank one**. Every row is a multiple of every other row. So each unit carries the same weight profile over time, scaled by its own u_i, and each period weights the units in the same pattern, scaled by its own v_t. There is no unit-by-period interaction in the weights.

That product structure is what makes this a difference-in-differences. A double difference contrasts units and contrasts periods, and multiplying the two contrasts gives a rank-one weighting. We come back to the rank below, where it separates the block designs from the staggered ones.

Multiply Equation 14.16 out and we get the 2×2 of Equation 14.14 with two of its four averages reweighted, the control average by ω and the pre-period average by λ:

$$\hat{\tau} = \Big(\bar{Y}_{T,\text{post}} - \sum_{t \in \text{pre}} \lambda_t \bar{Y}_{T,t}\Big) - \Big(\sum_{i \in C} \omega_i \bar{Y}_{i,\text{post}} - \sum_{i \in C} \sum_{t \in \text{pre}} \omega_i \lambda_t Y_{it}\Big). \tag{14.17}$$

The four blocks keep the signs $(+, -, -, +)$ of Equation 14.14. Only the averages inside two of them change, and they are no longer uniform.

Simulation — the outcome weights of synthetic DiD. A panel of 40 control and 10 treated units over 12 pre- and 6 post-periods. Untreated outcomes are

$$Y_{it}(0) = \alpha_i + \beta_t + f_i' g_t + \varepsilon_{it}, \tag{14.18}$$

with a unit effect $\alpha_i \sim N(0, 0.5^2)$, a time effect β_t rising linearly from 0 to 1, two factors in g_t (one linear in t, one a sine), loadings $f_i \sim N(0, \mathbf{I})$ shifted by 0.8 for the treated units, and $\varepsilon_{it} \sim N(0, 0.3^2)$. The spread in f_i makes the control trends differ from each other, and the shift makes the treated trend differ from all of them. The true ATT is 2. We fit with `synthdid`, then rebuild Equation 14.16 by hand from the fitted ω and λ.

```
library(synthdid)
set.seed(11)
N0 <- 40; N1 <- 10; N <- N0 + N1; T0 <- 12; T1 <- 6; Tt <- T0 + T1
f <- cbind(rnorm(N), rnorm(N)); f[(N0 + 1):N, ] <- f[(N0 + 1):N, ] + 0.8
g <- cbind(seq(0, 2, length = Tt), sin(seq(0, 3, length = Tt)))
Y <- outer(rnorm(N, 0, .5), rep(1, Tt)) + outer(rep(1, N), seq(0, 1, length =
 ↪  Tt)) +
     f %*% t(g) + matrix(rnorm(N * Tt, 0, .3), N, Tt)
W <- matrix(0, N, Tt); W[(N0 + 1):N, (T0 + 1):Tt] <- 1
Y <- Y + 2 * W                                          # true ATT = 2

est <- synthdid_estimate(Y, N0, T0)
wt  <- attr(est, "weights")
u <- c(-wt$omega,  rep(1 / N1, N1))                     # unit part
v <- c(-wt$lambda, rep(1 / T1, T1))                     # time part
Omega <- outer(u, v)                                    # outcome weight on cell
 ↪  (i, t)
```

Three routes should give the same number: what `synthdid` reports, the weighting Equation 14.16 applies to the observed outcomes, and the weighted regression of Equation 14.15.

```
cw <- outer(c(wt$omega, rep(1 / N1, N1)), c(wt$lambda, rep(1 / T1, T1)))
dd <- data.frame(y = as.vector(Y), Wt = as.vector(W), i = factor(rep(1:N, Tt)),
                 t = factor(rep(1:Tt, each = N)), cw = as.vector(cw))
tau3 <- c(`synthdid_estimate()`              = as.numeric(est),
          `$\\sum_{it} \\omega_{it} Y_{it}$, by hand` = sum(Omega * Y),
```

```
          `weighted TWFE regression`        = coef(lm(y ~ Wt + i + t, dd,
                                                      weights = dd$cw))[["Wt"]])
vtab(c(tau3, `largest gap between them` = max(abs(tau3 - tau3[1]))), 6)
```

	value
synthdid_estimate()	2.205444
$\sum_{it} \omega_{it} Y_{it}$, by hand	2.205444
weighted TWFE regression	2.205444
largest gap between them	0.000000

The gap between them is zero to machine precision. The outer-product form is what `synthdid` computes.

Writing the weights this way puts three constraints on them. First, the rank is one, since a cell's weight is a unit part times a time part. Second, each margin sums to zero on its own: ω and λ each sum to one, and the treated and post-period blocks each contribute an offsetting one. Third, no cell carries a wrong sign: no treated post-period cell is subtracted, and no control pre-period cell is subtracted either. We check all three.

```
chk <- rbind(`rank of $\\boldsymbol\\Omega$`     = c(sum(svd(Omega)$d > 1e-12),
 ↪  1),
             `$\\sum_i u_i$`      = c(sum(u), 0),
             `$\\sum_t v_t$`    = c(sum(v), 0),
             `treated post-cells, $\\omega_{it} < 0$` = c(sum(Omega[(N0+1):N,
              ↪  (T0+1):Tt] < 0), 0),
             `control pre-cells, $\\omega_{it} < 0$`  = c(sum(Omega[1:N0, 1:T0] <
              ↪  0), 0))
colnames(chk) <- c("found", "should be")
mtab(chk, 4)
```

	found	should be
rank of Ω	1	1
$\sum_i u_i$	0	0
$\sum_t v_t$	0	0
treated post-cells, $\omega_{it} < 0$	0	0
control pre-cells, $\omega_{it} < 0$	0	0

All three hold. Note that the double zero-sum is stronger than the single sum-to-zero of Equation 14.2, which constrains the weights only in total. Here they vanish along each margin. The signs are fixed by construction too: a treated post-period cell gets $1/(N_1 T_1) > 0$, and a control pre-period cell gets $\omega_i \lambda_t > 0$, because the simplex rules out a negative ω_i. The doubly-robust fit above produced 151 wrong-sign units. Synthetic DiD cannot produce any.

The remaining question is how much of the panel the comparison uses. Counting the non-zero weights is one answer, but it treats a weight of 0.4 and a weight of 0.001 alike. The usual summary that does not is the **effective sample size**, and it is worth deriving rather than asserting. Take m observations with equal variance σ^2. An unweighted mean of them has variance σ^2/m. A weighted mean with weights w summing to one has variance $\sigma^2 \sum w^2$. Ask which m makes the two equal:

$$\frac{\sigma^2}{m} = \sigma^2 \sum_j w_j^2 \qquad \Longrightarrow \qquad m = \frac{1}{\sum_j w_j^2}. \tag{14.19}$$

So Equation 14.19 is the number of equally weighted observations that would deliver the same precision as the weighting actually used. It equals the full count when every weight is equal, and 1 when one observation carries everything. We apply it to each margin separately: to the control weights as $1/\sum_i \omega_i^2$, and to the pre-period weights as $1/\sum_t \lambda_t^2$. Both are sparse here, so both fall below the nominal 40 controls and 12 pre-periods.

```
dsg <- c(sum(wt$omega > 1e-8), 1 / sum(wt$omega^2),
         sum(wt$lambda > 1e-8), 1 / sum(wt$lambda^2), wt$lambda[T0])
names(dsg) <- c(sprintf("controls with non-zero weight (of %d)", N0),
                sprintf("effective controls, $1/\\sum_i \\omega_i^2$ (of %d)",
                 ↪  N0),
                sprintf("pre-periods with non-zero weight (of %d)", T0),
                sprintf("effective pre-periods, $1/\\sum_t \\lambda_t^2$ (of
                 ↪  %d)", T0),
                "weight on the last pre-period")
vtab(dsg)
```

	value
controls with non-zero weight (of 40)	23.000
effective controls, $1/\sum_i \omega_i^2$ (of 40)	14.557
pre-periods with non-zero weight (of 12)	2.000
effective pre-periods, $1/\sum_t \lambda_t^2$ (of 12)	1.091
weight on the last pre-period	0.957

Only 23 of the 40 controls get any weight, and the effective count is 14.6. Three reference points make that number readable. Weighting all 40 equally would give 40, and that is plain difference-in-differences. Weighting the 23 active controls equally, at 1/23 each, would give 23. The comparison actually delivers 14.6, so the drop from 23 to 14.6 measures nothing but the inequality among those 23 weights: a few carry most of the mass and the rest contribute little.

A larger effective size is not automatically better, and it is worth seeing why. By Cauchy–Schwarz, $\sum_i \omega_i^2 \geq 1/N_0$ with equality only when every weight equals $1/N_0$, so the effective size reaches N_0 exactly when the weighting is uniform — that is, when no reweighting has happened. Matching the treated path requires tilting toward the controls that resemble it, and any tilt lowers the effective size. The two things we want pull against each other, and the effective count is the price paid for the match rather than a score to be maximised. The one case where both arrive together is when

the treated path already equals the unweighted control average, and then there was nothing to reweight for.

Synthetic difference-in-differences prices that trade-off explicitly, which is why the effective count is the number to read. Its weights minimise not the pre-period fit alone but

$$\text{(pre-period fit error)} \; + \; \zeta^2 \sum_i \omega_i^2, \tag{14.20}$$

and by Equation 14.19 the penalty $\zeta^2 \sum_i \omega_i^2$ is exactly ζ^2 divided by the effective sample size. So the estimator charges itself for concentration at a rate ζ, which by default rises with the estimated noise in the panel: noisier data is shrunk harder toward uniform, because in noisy data an apparent gain from concentrating is more likely to be fitted noise. The 14.6 above is where that trade-off settled on this panel, and it is computable whether or not we know what generated the data. The time margin is sparser: an effective 1.09 pre-periods, with 0.957 of the mass on the last one. The counterfactual level is anchored on the single period before treatment, so the estimate inherits that period's noise. We can read all of this off the weights. None of it is visible in $\hat{\tau}$.

We plot the two margins to see where the mass sits. The left panel sorts the fitted control weights ω_i from largest to smallest, which shows how many controls the comparison uses. The right panel keeps the pre-period weights λ_t in time order, which shows which periods the counterfactual is built from.

```
par(mfrow = c(1, 2), mar = c(4.2, 4.2, 2.6, 1))
os <- sort(wt$omega, decreasing = TRUE)
barplot(os, col = ifelse(os > 1e-8, teal, "grey85"), border = NA,
        xlab = "control units, sorted", ylab = expression(omega[i]),
        main = "Unit weights")
barplot(wt$lambda, names.arg = 1:T0, border = NA,
        col = ifelse(wt$lambda > 1e-8, rust, "grey85"),
        xlab = "pre-period", ylab = expression(lambda[t]), main = "Time weights")
```




Figure 14.5: The two margins of the synthetic DiD weight matrix. Left: fitted control-unit weights ω_i, sorted. 23 of 40 are non-zero, and the 17 the simplex sets to zero lie flat against the axis. Right: fitted pre-period weights λ_t. Almost all the mass is on the last pre-period, so the counterfactual level is anchored just before treatment.

14.5.3 The rank of the weight matrix

Synthetic difference-in-differences has a rank-one Ω by construction. So does the plain 2×2: set u_i and v_t to the uniform $\pm 1/N_g$ and $\pm 1/T_g$ and Equation 14.14 reappears. Staggered adoption gives a higher rank, bounded by the number of cohorts.

Write the staggered treatment matrix as a sum over adoption cohorts, $W = \sum_{k=1}^{K} c_k s_k'$. Here c_k indicates the units of cohort k, and s_k the periods from its adoption date on. That is a sum of K rank-one terms. Residualising on unit and time dummies is a double centring, $(\mathbf{I} - P)\, W\, (\mathbf{I} - Q)$, and double centring cannot raise a rank. So $\operatorname{rank}(\Omega) \leq K$, with equality generically.

We check the bound directly. Each panel has 60 units over 20 periods, with one to five adoption dates plus a never-treated group. For each we build the staggered TWFE weights, then count the non-zero singular values of Ω.

```
set.seed(4)
Ns <- 60; Ts <- 20
rk <- sapply(list(10, c(8, 14), c(6, 10, 14), c(6, 9, 12, 15), c(5, 8, 11, 14,
 ↪  17)),
  function(dates) {
    coh <- sample(c(dates, Inf), Ns, replace = TRUE)     # Inf = never treated
    Ws  <- matrix(0, Ns, Ts)
    for (i in 1:Ns) if (is.finite(coh[i])) Ws[i, coh[i]:Ts] <- 1
    ds  <- data.frame(Wt = as.vector(Ws), i = factor(rep(1:Ns, Ts)),
                      t = factor(rep(1:Ts, each = Ns)))
    Wtil <- residuals(lm(Wt ~ i + t, ds))
```

```
    Om   <- matrix(Wtil / sum(Wtil^2), Ns, Ts)
    c(`treated cohorts` = length(dates),
      `rank of $\\boldsymbol\\Omega$`   = sum(svd(Om)$d > 1e-9 * max(svd(Om)$d)))
  })
colnames(rk) <- c("1 date", "2 dates", "3 dates", "4 dates", "5 dates")
mtab(rk)
```

	1 date	2 dates	3 dates	4 dates	5 dates
treated cohorts	1	2	3	4	5
rank of Ω	1	2	3	4	5

The rank is at most the number of treated cohorts, and equals it generically. This ties the chapter's difference-in-differences material together. A block design has one adoption date, so its weight matrix is rank one. The plain 2×2 and synthetic difference-in-differences are the same object at two settings of the margins, uniform in the first and fitted in the second. Staggered adoption raises the rank. Once a cell's weight no longer factors into a unit part times a time part, a treated cell can enter negatively. In the block case it cannot, since every treated post-period cell gets $u_i v_t = 1/(N_1 T_1) > 0$. Rank one is not what does that work, though. An outer product uv' has negative entries as soon as either margin has mixed signs, so rank one constrains the *structure* of Ω and not the signs of its entries. What rules out a negative treated cell here is the block design's particular sign pattern, both margins uniform and positive.

14.5.4 Which margin does the work

The weight matrix has two margins, u over units and v over periods, and Equation 14.16 says the whole weighting is their product. So a natural question about the structure is which margin carries the correction when the control trends differ from the treated group's.

It is worth saying first what such a correction can and cannot be. Parallel trends is a statement about two group means, $E[\Delta Y(0) \mid D = 1] = E[\Delta Y(0) \mid D = 0]$, and no single observation either has it or lacks it, so in the model behind Equation 14.14 there is nothing at the unit level for a weight to act on. Weighting bites only under a model where trends vary across units: unit-specific slopes, or a factor structure. Then each unit has its own trend, and reweighting controls toward those whose loadings resemble the treated group's is well defined. That is what synthetic difference-in-differences does, and what the wider family of trajectory-balancing and history-matching estimators does too (Hazlett and Xu 2018; Imai, Kim, and Wang 2023; Doudchenko and Imbens 2016).

We can then hold one margin uniform and fit the other, and read off how much of the correction each supplies.

Simulation — which margin does the work. Two panels of 40 controls and 10 treated over 12 pre- and 6 post-periods, with a true ATT of 2. Both carry a unit effect $\alpha_i \sim N(0, 1)$, shifted up by 1 for the treated units, a time effect β_t rising linearly from 0 to 1.5, and a serially correlated shock. The shock is a stationary AR(1),

$$\varepsilon_{it} = \rho\, \varepsilon_{i,t-1} + \nu_{it}, \qquad \nu_{it} \sim N(0, \sigma^2), \qquad \rho = 0.5, \tag{14.21}$$

started from its stationary distribution, so ε_{it} has standard deviation $\sigma/\sqrt{1-\rho^2}$. We set $\sigma = 0.4$, which makes the shock's own standard deviation 0.46. Note the distinction: σ is the standard deviation of the innovation ν_{it} that arrives each period, not of the shock ε_{it} that accumulates.

The shock is serially correlated rather than independent for a reason, and it is not decoration. The weights are fitted on the pre-period path. Under independent noise, averaging across twelve pre-periods removes most of it, so the weights would recover the systematic path almost exactly and there would be little left to study. Serial correlation means twelve periods carry well under twelve periods' worth of independent information, and that is what leaves the fitted weights short. It is also the realistic case: panel outcomes persist. AR(1) is the smallest model with a persistence parameter, and it nests independent noise at $\rho = 0$. It does not contain a random walk. The process here starts from its stationary distribution, whose variance $\sigma^2/(1-\rho^2)$ diverges as $\rho \to 1$, so the stationary family reaches ever greater persistence without ever arriving at the unit root.

These three terms are common to almost any panel, and none of them breaks parallel trends: α_i differences away and β_t is shared by both groups, so a difference-in-differences removes them whatever the weights are.

Parallel trends fails only when the treated and control groups have *different* untreated trends, so a term that varies by unit and moves with t is what we have to put in. Without one there is nothing for the weights to fix. The two panels are two forms of it. The `factor` panel uses the $f_i' g_t$ of Equation 14.18. The `slopes` panel replaces it with a unit-specific linear trend,

$$Y_{it}(0) = \alpha_i + \beta_t + s_i\, t + \varepsilon_{it}, \qquad s_i \sim N(0,\, 0.08^2), \tag{14.22}$$

with s_i shifted up by 0.12 for the treated units. So each unit has its own growth rate, and the treated units grow faster on average, by about one and a half standard deviations of the control slopes. Note that $s_i t$ is itself a one-factor model, with loading s_i and factor $g_t = t$. It is the simplest member of the class Equation 14.18 describes, which is the class synthetic difference-in-differences is built for, so this is a leading case and not a special one. We run synthetic DiD four ways: both margins uniform, which is plain difference-in-differences; unit weights fitted with time weights uniform; time weights fitted with unit weights uniform; and both fitted. 400 replications.

```
library(parallel)
panel <- function(kind, N0 = 40, N1 = 10, T0 = 12, T1 = 6, sig = .4, rho = .5) {
  N <- N0 + N1; Tt <- T0 + T1
  a <- rnorm(N, 0, 1); a[(N0 + 1):N] <- a[(N0 + 1):N] + 1
  e <- matrix(0, N, Tt); e[, 1] <- rnorm(N, 0, sig / sqrt(1 - rho^2))
  for (t in 2:Tt) e[, t] <- rho * e[, t - 1] + rnorm(N, 0, sig)
  Y0 <- outer(a, rep(1, Tt)) + outer(rep(1, N), seq(0, 1.5, length = Tt)) + e
  if (kind == "factor") {
    f <- cbind(rnorm(N), rnorm(N)); f[(N0 + 1):N, ] <- f[(N0 + 1):N, ] + 0.8
    g <- cbind(seq(0, 2, length = Tt), sin(seq(0, 3, length = Tt)))
    Y0 <- Y0 + f %*% t(g)
  } else {
    s <- rnorm(N, 0, .08); s[(N0 + 1):N] <- s[(N0 + 1):N] + .12   # unit slopes
    Y0 <- Y0 + outer(s, 1:Tt)
  }
```

```
  W <- matrix(0, N, Tt); W[(N0 + 1):N, (T0 + 1):Tt] <- 1
  list(Y = Y0 + 2 * W, N0 = N0, N1 = N1, T0 = T0, T1 = T1)
}
margins <- function(r, kind) {
  set.seed(5000 + r); p <- panel(kind); N0 <- p$N0; T0 <- p$T0
  fx <- function(om, lam) as.numeric(synthdid_estimate(p$Y, N0, T0,
          weights = list(omega = om, lambda = lam)))     # NULL = fit this margin
  unif.o <- rep(1 / N0, N0); unif.l <- rep(1 / T0, T0)
  c(`both uniform (DiD)` = fx(unif.o, unif.l), `unit weights only` = fx(NULL,
   ↪  unif.l),
    `time weights only`  = fx(unif.o, NULL),   `both fitted (SDID)` = fx(NULL,
     ↪  NULL))
}
res <- lapply(c("factor", "slopes"), function(kind) {
  # portable core count: mclapply falls back to serial on Windows
  n_cores <- if (.Platform$OS.type == "windows") 1L else
               max(1L, min(6L, parallel::detectCores() - 1L))
  M <- simplify2array(mclapply(1:400, margins, kind = kind, mc.cores = n_cores))
  cbind(bias = rowMeans(M) - 2, sd = apply(M, 1, sd), rmse = sqrt(rowMeans((M -
   ↪  2)^2)))
})
out <- cbind(res[[1]], res[[2]])
colnames(out) <- paste(rep(c("factor", "slopes"), each = 3), colnames(res[[1]]))
mtab(out)
```

	factor bias	factor sd	factor rmse	slopes bias	slopes sd	slopes rmse
both uniform (DiD)	0.729	0.403	0.833	1.083	0.284	1.120
unit weights only	0.173	0.337	0.379	0.603	0.231	0.645
time weights only	0.082	0.285	0.296	0.536	0.203	0.573
both fitted (SDID)	0.044	0.299	0.302	0.403	0.208	0.453

Reweighting helps, and by a lot. Under the factor structure the bias falls from 0.73 to 0.04 with both margins fitted. Each margin helps alone: 0.17 from the unit weights, 0.08 from the time weights. Which margin matters is not fixed in advance. Here the time weights do slightly more, because the factor paths are smooth, so anchoring on the periods just before treatment removes most of the divergence. The fitted λ above showed the same thing when it put 0.957 on the last pre-period.

The `slopes` panel gains less: 1.08 down to 0.40. The reason is that the weights are fitted on the observed pre-period path, which carries the shock of Equation 14.21 as well as the trend, so they match a noisy version of what they are aiming at and fall short. The size of that shortfall depends on the pre-period length and the noise level rather than on anything intrinsic to reweighting — with a much cleaner pre-period the same estimator gets the bias under 0.05 — so the direction is the finding here and the magnitude is not.

Two limits are worth stating even though they do not bind in this panel. Reweighting controls can only produce a weighted average of the control trends, so a treated trend steeper than every control's is out of reach; that is the same overlap problem seen earlier, now in the design, and the simplex is what makes it a hard boundary rather than a silent extrapolation. And if we are confident the trends really are unit-specific and linear, the direct response is to put unit-specific linear trends in the regression, not to reweight at all. Reweighting is the tool for when we will not commit to a functional form, which is the case Equation 14.18 stands in for.

14.5.5 Implied versus chosen

One distinction cuts across everything above. OLS, AIPW, DML, TWFE, and the imputation estimator all *imply* weights — they fall out of the estimation and are diagnosed afterward. Synthetic controls and synthetic difference-in-differences *choose* weights as part of the design, selecting ω to balance pre-treatment outcomes or covariates. For those estimators $\hat{\tau} = \omega' Y$ is the definition, not a decomposition.

That difference changes how the weights should be read, in three ways. First, chosen weights are fitted on Y itself, so $\hat{\tau}$ is not linear in Y and Equation 14.16 describes the estimate at the fitted weights rather than decomposing it the way Equation 14.2 decomposes OLS. Inference has to account for the weights having been estimated, which is why `synthdid` reports standard errors by jackknife or bootstrap rather than treating ω as fixed.

Second, a clean-looking balance diagnostic is weaker evidence here than it looks. When the weights were chosen to make the pre-period match, the match is not a test the design passed — it is what the design optimised. Fitting weights on a noisy pre-period statistic is a close cousin of a pretest. The analogy is with Roth (2022), who shows that conditioning an analysis on passing a parallel-trends test distorts the estimate and undercovers. Weight-fitting is not literally that procedure, since nothing is accepted or rejected, but it shares the mechanism: noise in the pre-period is allowed to select the analysis. Matching on pre-period *levels* is the sharp case: it induces regression to the mean, with a bias that grows as serial correlation weakens (Daw and Hatfield 2018; Chabé-Ferret 2015, 2017). Synthetic difference-in-differences avoids that particular trap by fitting ω with an intercept, so it never needs to match levels, only trajectories up to a constant.

Third, the effective count of Equation 14.19 is the number that reports the price. A design that matches well by concentrating weight on a few controls has achieved that match by lowering the effective sample size, and Equation 14.20 shows the estimator charging itself for exactly that. Read the two together rather than either alone.

A closing caution. These weights diagnose estimation. They show which comparisons a difference-in-differences rests on, and whether it uses non-overlapping or forbidden ones. They do not test identification. Parallel trends is an assumption about the treated group's unobserved untreated path, and a negative weight and a failure of parallel trends stay independent: either can occur without the other. Chosen weights can give robustness to one class of violations, those the design's trend model can express, and the weights let us see what that cost. No weighting of observed outcomes tests the assumption. To do that we bound the violation instead of reweighting, using the observed pre-period differences to limit how large the post-period departure can be (Rambachan and Roth 2023).

14.6 Summary

1. **A signed weighting of the outcomes.** FWL writes $\hat{\beta} = \sum_i \omega_i y_i$ with $\omega_i = \tilde{V}_i / \sum_j \tilde{V}_j^2$ (Equation 14.2): a signed, computable weighting of the observed outcomes, different from the effect weights $w_i = \tilde{V}_i V_i$.
2. **Two constraints.** The weights sum to zero and the weighted regressor sums to one: $\sum_i \omega_i = 0$ and $\sum_i \omega_i V_i = 1$.
3. **The sign shows direction.** Because ω_i is proportional to $\tilde{V}_i$, not to $\tilde{V}_i^2$, it carries a sign: it shows which rows push $\hat{\beta}$ up and which pull it down, and, under controls with thin overlap, which enter with the sign their raw position would not predict.
4. **The same weights beyond OLS.** Any estimator that can be written $\hat{\tau} = \omega' Y$ has outcome weights, so the same object and its checks reach AIPW, double machine learning, and causal forests (Knaus 2024). PLR's denominator $\tilde{D}'\tilde{D}$ reveals its estimand as the VWATE (Equation 14.11), not the ATE; and the sum-to-zero of OLS need not become a sum to ± 1 within the treated and control groups.
5. **From difference in means to difference-in-differences.** Regressed on the treatment alone, $\hat{\beta}$ is the difference in group means; a 2×2 difference-in-differences is a signed contrast of four cell means. With covariates, a doubly-robust DiD weights the outcome changes, $\widehat{\text{ATT}} = \omega' \Delta Y$ (Sant'Anna and Zhao 2020), open to the same checks.
6. **Staggered designs.** In staggered TWFE the outcome weights flag treated cells used as controls — the observation-level version of the Goodman-Bacon (2021) forbidden comparisons. Untreated cells with positive weight are mostly control baselines, which every DiD adds. The negative weights bias the estimate only when effects change over time; with constant effects the gap from the ATT is variance weighting. Estimators that fit counterfactuals on untreated observations alone (Borusyak, Jaravel, and Spiess 2024; Callaway and Sant'Anna 2021) cannot assign negative weight to any treated cell, so the forbidden-comparison diagnostic is clean by construction.
7. **Rank counts the cohorts.** As an $N \times T$ matrix, the outcome weights of a block design have rank one and factor into a unit part times a time part (Equation 14.16). The plain 2×2 and synthetic difference-in-differences differ only in whether those margins are uniform or fitted. Staggered adoption raises the rank, to at most the number of treated cohorts and generically to exactly that. Rank one is not itself what keeps treated cells positive: in the block design that follows from both margins being uniform and positive, and a rank-one matrix with mixed-sign margins has negative entries.
8. **Both margins carry correction, and the effective count reports its price.** Weighting can offset differing trends only under a model in which trends vary by unit; then either margin helps alone and both together help most. What it costs is effective sample size: by Equation 14.19 the count reaches N_0 only under uniform weights, so concentrating weight lowers the effective count. The optimisation is trading fit against that concentration, and Equation 14.20 shows synthetic difference-in-differences charging itself for exactly that.
9. **Chosen weights are read differently from implied ones.** Weights fitted on Y make $\hat{\tau}$ non-linear in Y, so inference must account for their estimation, and a good pre-period match is what the design optimised rather than a test it passed. Fitting on a noisy pre-period statistic is a close cousin of a pretest rather than the thing itself: Roth (2022) studies conditioning on a pass/fail test, and what the two share is that pre-period noise gets to select the analysis. Matching on pre-period *levels* invites regression to the mean (Daw and Hatfield 2018); fitting with an intercept, so only trajectories need to match, avoids that case.

References

This guide draws on a handful of textbooks, the original papers behind the estimators discussed, and package documentation. The list below collects every work the guide rests on. The bibliography source is `references.bib` in the book repository. Keys shared with the two causal econometrics guides and the blog book are kept unchanged, so a passage can move between the books without a key being renamed.

Besides the works cited in the chapters, the guide draws throughout on the textbooks of Baltagi (2013), Baum (2006) and Greene (1993). Greene is cited from the 1993 edition; chapter, example and table numbers differ across editions.

The R packages loaded in the code examples are `cobalt`, `DRDID`, `fixest`, `ggfortify`, `gmm`, `ivreg`, `knitr`, `MASS`, `OutcomeWeights`, `plm`, `sandwich`, `strucchange`, `survival`, `synthdid`, and `WeightIt`, run under R (R Core Team). Stata examples use Stata (StataCorp).

Allison, Paul D., and Richard P. Waterman. 2002. "Fixed-Effects Negative Binomial Regression Models." *Sociological Methodology* 32 (1): 247–65. https://doi.org/10.1111/1467-9531.00117.

Anderson, T. W., and Cheng Hsiao. 1981. "Estimation of Dynamic Models with Error Components." *Journal of the American Statistical Association* 76 (375): 598–606. https://doi.org/10.1080/01621459.1981.10477691.

Andrews, Donald W. K., Marcelo J. Moreira, and James H. Stock. 2006. "Optimal Two-Sided Invariant Tests for Instrumental Variables Regression." *Econometrica* 74 (3): 715–52. https://doi.org/10.1111/j.1468-0262.2006.00680.x.

Angrist, Joshua D. 1998. "Estimating the Labor Market Impact of Voluntary Military Service Using Social Security Data on Military Applicants." *Econometrica* 66 (2): 249–88. https://doi.org/10.2307/2998558.

Arellano, Manuel, and Stephen Bond. 1991. "Some Tests of Specification for Panel Data." *Review of Economic Studies* 58 (2): 277–97. https://doi.org/10.2307/2297968.

Arkhangelsky, Dmitry, Susan Athey, David A. Hirshberg, Guido W. Imbens, and Stefan Wager. 2021. "Synthetic Difference-in-Differences." *American Economic Review* 111 (12): 4088–118.

Baltagi, Badi H. 2013. *Econometric Analysis of Panel Data*. 5th ed. Wiley.

Baltagi, Badi H., and Dan Levin. 1992. "Cigarette Taxation: Raising Revenues and Reducing Consumption." *Structural Change and Economic Dynamics* 3 (2): 321–35. https://doi.org/10.1016/0954-349x(92)90010-4.

Baum, Christopher F. 2006. *An Introduction to Modern Econometrics Using Stata*. Stata Press.

Blundell, Richard, and Stephen Bond. 1998. "Initial Conditions and Moment Restrictions in Dynamic Panel Data Models." *Journal of Econometrics* 87 (1): 115–43. https://doi.org/10.1016/s0304-4076(98)00009-8.

Borusyak, Kirill, Xavier Jaravel, and Jann Spiess. 2024. "Revisiting Event Study Designs: Robust and Efficient Estimation." *Review of Economic Studies* 91 (6): 3253–85.

Breslow, N. 1974. "Covariance Analysis of Censored Survival Data." *Biometrics* 30 (1): 89–99. https://doi.org/10.2307/2529620.

Buuren, Stef van. 2018. *Flexible Imputation of Missing Data*. 2nd ed. CRC Press. https://doi.org/10.1201/9780429492259.

Callaway, Brantly, and Pedro H. C. Sant'Anna. 2021. "Difference-in-Differences with Multiple Time Periods." *Journal of Econometrics* 225 (2): 200–230.

Cameron, A. Colin, and Pravin K. Trivedi. 2005. *Microeconometrics: Methods and Applications*. Cambridge University Press. https://doi.org/10.1017/CBO9780511811241.

Chabé-Ferret, Sylvain. 2015. "Analysis of the Bias of Matching and Difference-in-Difference Under Alternative Earnings and Selection Processes." *Journal of Econometrics* 185 (1): 110–23. https://doi.org/10.1016/j.jeconom.2014.09.013.

———. 2017. "Should We Combine Difference in Differences with Conditioning on Pre-Treatment Outcomes?" TSE Working Paper 17-824. Toulouse School of Economics.

Chaisemartin, Clément de, and Xavier D'Haultfœuille. 2020. "Two-Way Fixed Effects Estimators with Heterogeneous Treatment Effects." *American Economic Review* 110 (9): 2964–96.

Chattopadhyay, Ambarish, and José R Zubizarreta. 2023. "On the Implied Weights of Linear Regression for Causal Inference." *Biometrika* 110 (3): 615–29. https://doi.org/10.1093/biomet/asac058.

Cochrane, D., and G. H. Orcutt. 1949. "Application of Least Squares Regression to Relationships Containing Auto-Correlated Error Terms." *Journal of the American Statistical Association* 44 (245): 32–61. https://doi.org/10.1080/01621459.1949.10483290.

Cox, David R. 1972. "Regression Models and Life-Tables." *Journal of the Royal Statistical Society: Series B (Methodological)* 34 (2): 187–202.

Crump, R. K., V. J. Hotz, G. W. Imbens, and O. A. Mitnik. 2009. "Dealing with Limited Overlap in Estimation of Average Treatment Effects." *Biometrika* 96 (1): 187–99. https://doi.org/10.1093/biomet/asn055.

Davidson, Russell, and James G. MacKinnon. 2004. *Econometric Theory and Methods*. Oxford University Press.

Daw, Jamie R., and Laura A. Hatfield. 2018. "Matching and Regression to the Mean in Difference-in-Differences Analysis." *Health Services Research* 53 (6): 4138–56. https://doi.org/10.1111/1475-6773.12993.

Donald, Stephen G, and Kevin Lang. 2007. "Inference with Difference-in-Differences and Other Panel Data." *Review of Economics and Statistics* 89 (2): 221–33. https://doi.org/10.1162/rest.89.2.221.

Doudchenko, Nikolay, and Guido Imbens. 2016. "Balancing, Regression, Difference-in-Differences and Synthetic Control Methods: A Synthesis." NBER Working Paper 22791. National Bureau of Economic Research. https://doi.org/10.3386/w22791.

Efron, Bradley. 1977. "The Efficiency of Cox's Likelihood Function for Censored Data." *Journal of the American Statistical Association* 72 (359): 557–65. https://doi.org/10.1080/01621459.1977.10480613.

Frisch, Ragnar, and Frederick V. Waugh. 1933. "Partial Time Regressions as Compared with Individual Trends." *Econometrica* 1 (4): 387–401. https://doi.org/10.2307/1907330.

Gibbons, Charles E., Juan Carlos Suárez Serrato, and Michael B. Urbancic. 2018. "Broken or Fixed Effects?" *Journal of Econometric Methods* 8 (1): 1–12. https://doi.org/10.1515/jem-2017-0002.

Goodman-Bacon, Andrew. 2021. "Difference-in-Differences with Variation in Treatment Timing." *Journal of Econometrics* 225 (2): 254–77.

Greene, William H. 1993. *Econometric Analysis*. 2nd ed. Macmillan.

Grogger, J. T., and R. T. Carson. 1991. "Models for Truncated Counts." *Journal of Applied Econometrics* 6 (3): 225–38. https://doi.org/10.1002/jae.3950060302.

Guimarães, Paulo. 2008. "The Fixed Effects Negative Binomial Model Revisited." *Economics Letters* 99 (1): 63–66. https://doi.org/10.1016/j.econlet.2007.05.030.

Hamilton, James D. 1994. *Time Series Analysis*. Princeton University Press.

Hausman, J. A. 1978. "Specification Tests in Econometrics." *Econometrica* 46 (6): 1251–71. https://doi.org/10.2307/1913827.

Hausman, Jerry, Bronwyn H. Hall, and Zvi Griliches. 1984. "Econometric Models for Count Data with an Application to the Patents-R&D Relationship." *Econometrica* 52 (4): 909–38. https://doi.org/10.2307/1911191.

Hazlett, Chad, and Yiqing Xu. 2018. "Trajectory Balancing: A General Reweighting Approach to Causal Inference with Time-Series Cross-Sectional Data." Working paper 3214231. SSRN. https://doi.org/10.2139/ssrn.3214231.

Heckman, James J. 1979. "Sample Selection Bias as a Specification Error." *Econometrica* 47 (1): 153–61. https://doi.org/10.2307/1912352.

Imai, Kosuke, In Song Kim, and Erik H. Wang. 2023. "Matching Methods for Causal Inference with Time-Series Cross-Sectional Data." *American Journal of Political Science* 67 (3): 587–605. https://doi.org/10.1111/ajps.12685.

Imbens, Guido W., and Joshua D. Angrist. 1994. "Identification and Estimation of Local Average Treatment Effects." *Econometrica* 62 (2): 467–75.

Judson, Ruth A, and Ann L Owen. 1999. "Estimating Dynamic Panel Data Models: A Guide for Macroeconomists." *Economics Letters* 65 (1): 9–15. https://doi.org/10.1016/s0165-1765(99)00130-5.

Knaus, Michael C. 2024. "Treatment Effect Estimators as Weighted Outcomes." https://arxiv.org/abs/2411.11559.

Li, Fan, Kari Lock Morgan, and Alan M. Zaslavsky. 2018. "Balancing Covariates via Propensity Score Weighting." *Journal of the American Statistical Association* 113 (521): 390–400. https://doi.org/10.1080/01621459.2016.1260466.

Lovell, Michael C. 1963. "Seasonal Adjustment of Economic Time Series and Multiple Regression Analysis." *Journal of the American Statistical Association* 58 (304): 993–1010. https://doi.org/10.1080/01621459.1963.10480682.

Montiel Olea, José Luis, and Carolin Pflueger. 2013. "A Robust Test for Weak Instruments." *Journal of Business & Economic Statistics* 31 (3): 358–69. https://doi.org/10.1080/00401706.2013.806694.

Moreira, Marcelo J. 2003. "A Conditional Likelihood Ratio Test for Structural Models." *Econometrica* 71 (4): 1027–48. https://doi.org/10.1111/1468-0262.00438.

Moulton, Brent R. 1990. "An Illustration of a Pitfall in Estimating the Effects of Aggregate Variables on Micro Units." *Review of Economics and Statistics* 72 (2): 334–38. https://doi.org/10.2307/2109724.

Mundlak, Yair. 1978. "On the Pooling of Time Series and Cross Section Data." *Econometrica* 46 (1): 69–85. https://doi.org/10.2307/1913646.

Newey, W. K., and K. D. West. 1994. "Automatic Lag Selection in Covariance Matrix Estimation." *Review of Economic Studies* 61 (4): 631–53. https://doi.org/10.2307/2297912.

Newey, Whitney K., and Kenneth D. West. 1987. "A Simple, Positive Semi-Definite, Heteroskedasticity and Autocorrelation Consistent Covariance Matrix." *Econometrica* 55 (3): 703–8. https://doi.org/10.2307/1913610.

Nickell, Stephen. 1981. "Biases in Dynamic Models with Fixed Effects." *Econometrica* 49 (6): 1417–26. https://doi.org/10.2307/1911408.

Peto, Richard. 1972. "Contribution to the Discussion of Paper by D. R. Cox." *Journal of the Royal Statistical Society, Series B* 34 (2): 205–7.

Prais, S. J., and C. B. Winsten. 1954. "Trend Estimators and Serial Correlation." Cowles Commission Discussion Paper 383. Cowles Commission.

Rambachan, Ashesh, and Jonathan Roth. 2023. "A More Credible Approach to Parallel Trends." *Review of Economic Studies* 90 (5): 2555–91. https://doi.org/10.1093/restud/rdad018.

Roth, Jonathan. 2022. "Pretest with Caution: Event-Study Estimates After Testing for Parallel Trends." *American Economic Review: Insights* 4 (3): 305–22. https://doi.org/10.1257/aeri.20210236.

Rubin, Donald B. 1987. *Multiple Imputation for Nonresponse in Surveys*. Wiley. https://doi.org/10.1002/9780470316696.

Sant'Anna, Pedro H. C., and Jun Zhao. 2020. "Doubly Robust Difference-in-Differences Estimators." *Journal of Econometrics* 219 (1): 101–22. https://doi.org/10.1016/j.jeconom.2020.06.003.

Sargan, J. D. 1958. "The Estimation of Economic Relationships Using Instrumental Variables." *Econometrica* 26 (3): 393–415. https://doi.org/10.2307/1907619.

Shinkre, Tanvi, and Chad Hazlett. 2024. "Demystifying and Avoiding the OLS "Weighting Problem": Unmodeled Heterogeneity and Straightforward Solutions." https://doi.org/10.48550/arXiv.2403.03299.

Singer, Judith D., and John B. Willett. 1993. "It's about Time: Using Discrete-Time Survival Analysis to Study Duration and the Timing of Events." *Journal of Educational Statistics* 18 (2): 155–95. https://doi.org/10.3102/10769986018002155.

Słoczyński, Tymon. 2022. "Interpreting OLS Estimands When Treatment Effects Are Heterogeneous: Smaller Groups Get Larger Weights." *The Review of Economics and Statistics* 104 (3): 501–9.

Stock, James H., and Motohiro Yogo. 2005. "Testing for Weak Instruments in Linear IV Regression." In *Identification and Inference for Econometric Models*, 80–108. Cambridge University Press. https://doi.org/10.1017/cbo9780511614491.006.

Sun, Liyang, and Sarah Abraham. 2021. "Estimating Dynamic Treatment Effects in Event Studies with Heterogeneous Treatment Effects." *Journal of Econometrics* 225 (2): 175–99.

Swamy, P. A. V. B., and S. S. Arora. 1972. "The Exact Finite Sample Properties of the Estimators of Coefficients in the Error Components Regression Models." *Econometrica* 40 (2): 261–75. https://doi.org/10.2307/1909405.

White, Halbert. 1980. "A Heteroskedasticity-Consistent Covariance Matrix Estimator and a Direct Test for Heteroskedasticity." *Econometrica* 48 (4): 817–38. https://doi.org/10.2307/1912934.

Wooldridge, Jeffrey M. 1999. "Distribution-Free Estimation of Some Nonlinear Panel Data Models." *Journal of Econometrics* 90 (1): 77–97. https://doi.org/10.1016/S0304-4076(98)00033-5.

———. 2010. *Econometric Analysis of Cross Section and Panel Data*. 2nd ed. MIT Press.