

Solution to Ramanujan Challenge Problem 3.1: An integral over knot polynomial roots expressing π^2

Xiang Huang

July 2026

Abstract

Let $A_{7_2}(M, L)$ be the A -polynomial of the knot 7_2 , let $\alpha \approx 0.349269$ and $\beta \approx 0.406813$ be the endpoints specified in the problem, and let $y(x)$ be the indicated branch. We prove

$$\int_{\alpha}^{\beta} \left(\log x \frac{dy}{y} - \log y \frac{dx}{x} \right) = \frac{4\pi^2}{85}.$$

By Khoi's variation formula the integral is a difference of Godbillon–Vey invariants of the endpoint representations; the β endpoint is the maximal Fuchsian representation of $\Sigma(2, 3, 17)$ and is evaluated by Brooks–Goldman. The new ingredient is a proof that the two extended Bloch classes are *torsion*. This follows from a single structural fact: the endpoint minimal polynomials are palindromic, so every non-real embedding of the endpoint fields lies on the unit circle, and on the unit circle two of the four tetrahedron shapes are real while the other two are exchanged by $z \mapsto 1/(1 - \bar{z})$, under which the Bloch–Wigner function changes sign. Torsion gives rationality; the Merkurjev–Suslin torsion order supplies the denominator bound 2040; a 301-digit evaluation then determines the value uniquely.

1 Setup

Write $A = A_{7_2}(M, L)$ for the A -polynomial of 7_2 , of degree 22 in M and 5 in L , as displayed in the problem statement. Put

$$\alpha : A(\alpha, \alpha^{1/2}) = 0, \quad \beta : A(\beta, \beta) = 0,$$

the roots nearest 0.349269 and 0.406813, and let $y(x)$ be the branch with $y > 0$, $y' \leq -2$ on $[\alpha, \beta]$. Set

$$I := \int_{\alpha}^{\beta} \left(\log x \frac{dy}{y} - \log y \frac{dx}{x} \right).$$

It is convenient to use the eigenvalue coordinate. Writing $M = a^2$, $L = a$ at the α endpoint and $M = L = b$ at the β endpoint, and factoring $A(a^2, a)$ resp. $A(x, x)$ over $\mathbb{Q}$, the endpoints are algebraic numbers with explicit minimal polynomials:

Proposition 1 (Endpoint fields). *The endpoint eigenvalues satisfy*

$$f_{\alpha}(a) = a^{12} - 3a^{11} + 4a^{10} - 5a^9 + 6a^8 - 7a^7 + 7a^6 - 7a^5 + 6a^4 - 5a^3 + 4a^2 - 3a + 1, \quad (1)$$

$$\begin{aligned} f_{\beta}(b) = & b^{16} - 7b^{15} + 22b^{14} - 48b^{13} + 87b^{12} - 133b^{11} + 178b^{10} - 211b^9 + 223b^8 \\ & - 211b^7 + 178b^6 - 133b^5 + 87b^4 - 48b^3 + 22b^2 - 7b + 1. \end{aligned} \quad (2)$$

Both are irreducible over $\mathbb{Q}$ and palindromic. Their signatures are $(r_1, r_2) = (2, 5)$ and $(4, 6)$, and

$$a = 0.59098942867025644\dots, \quad a^2 = \alpha, \quad b = 0.40681308133679\dots = \beta.$$

2 Reduction to a regulator difference

Khoi's variation formula for the one-form $\log x d \log y - \log y d \log x$ on an A -polynomial curve identifies I with a difference of Godbillon–Vey invariants of the two endpoint representations of π_1 of the 7_2 complement, with the normalisation

$$I = \frac{\text{GV}(\rho_\alpha) - \text{GV}(\rho_\beta)}{4}. \quad (3)$$

Proposition 2 (β endpoint). $S_{-1}^3(7_2) \cong \Sigma(2, 3, 17)$, the β representation factors through $\pi_1(\Sigma(2, 3, 17))$ as the maximal Fuchsian representation, and

$$\text{GV}(\rho_\beta) = \frac{242\pi^2}{51}.$$

This is Brooks–Goldman applied to the Seifert data of $\Sigma(2, 3, 17)$: with $\chi_{\text{orb}} = -11/102$ and $e = -1/102$, the maximal value is $4\pi^2\chi_{\text{orb}}^2/e = -242\pi^2/51$ up to orientation.

By (3) and Proposition 2, the conjecture $I = 4\pi^2/85$ is *equivalent* to

$$\text{GV}(\rho_\alpha) = \frac{242\pi^2}{51} + \frac{16\pi^2}{85} = \frac{1258\pi^2}{255} = \frac{74\pi^2}{15}. \quad (4)$$

Finally, the Neumann–Zagier differential identity $d\mathcal{R} = -(\log M d \log L - \log L d \log M)$ gives

$$I = -\Delta\mathcal{R}, \quad \mathcal{R} = \sum_{j \in \{T, U, V, W\}} \hat{R}(z_j), \quad \hat{R}(z) = \text{Li}_2(z) + \frac{1}{2} \log z \log(1 - z) - \frac{\pi^2}{6}, \quad (5)$$

where $z_T, \dots, z_W$ are the four tetrahedron shapes of the 7_2 ideal triangulation (SnapPy's triangulation of 7_2 has exactly four tetrahedra, all positively oriented at the complete structure).

3 The deformation chart

With $X = M^2$ put

$$u = \frac{L + X^3}{X(L + X)}, \quad r = \frac{-(1 + \sqrt{1 + 4u^2})}{2u}, \quad T = 1 - r^2, \quad U = u, \quad V = \frac{u}{X}, \quad W = \frac{1}{1 - uX}. \quad (6)$$

At the two endpoints this gives, to the displayed precision,

$$\begin{aligned} T_\alpha &= -0.15787528\dots, & U_\alpha &= 6.8157987\dots, & V_\alpha &= 55.872474\dots, & W_\alpha &= 5.9329217\dots, \\ T_\beta &= -0.25828783\dots, & U_\beta &= 4.3429623\dots, & V_\beta &= 26.241958\dots, & W_\beta &= 3.5555141\dots, \end{aligned}$$

all real, with $T < 0$ and $U, V, W > 1$: the endpoint representations are real, and the tetrahedra are flat.

The chart (6) involves $\sqrt{1 + 4u^2}$, so a priori the shapes lie in a quadratic extension of the endpoint field. They do not.

Lemma 3 (Shape field). $1 + 4u^2$ is a square in the endpoint field itself. Explicitly, with $s_\alpha, s_\beta \in \mathbb{Z}[z]$

$$\begin{aligned} s_\alpha &= 12z^{11} - 30z^{10} + 32z^9 - 42z^8 + 50z^7 - 58z^6 + 54z^5 - 56z^4 + 42z^3 - 38z^2 + 28z - 21, \\ s_\beta &= 60z^{15} - 394z^{14} + 1150z^{13} - 2386z^{12} + 4196z^{11} - 6176z^{10} + 8022z^9 - 9202z^8 \\ &\quad + 9406z^7 - 8592z^6 + 6958z^5 - 4960z^4 + 3064z^3 - 1546z^2 + 646z - 137, \end{aligned}$$

one has $s_\alpha(a)^2 = 1 + 4u^2$ in $\mathbb{Q}(a)$ and $s_\beta(b)^2 = 1 + 4u^2$ in $\mathbb{Q}(b)$. Hence all four shapes lie in the endpoint field, and the shape field E coincides with F .

Proof. Writing u itself as a polynomial modulo the minimal polynomial,

$$\begin{aligned} u_\alpha &= -6z^{11} + 14z^{10} - 15z^9 + 21z^8 - 23z^7 + 28z^6 - 25z^5 + 27z^4 - 20z^3 + 18z^2 - 13z + 10, \\ u_\beta &= -6z^{15} + 41z^{14} - 125z^{13} + 266z^{12} - 474z^{11} + 711z^{10} - 935z^9 + 1088z^8 \\ &\quad - 1127z^7 + 1043z^6 - 857z^5 + 620z^4 - 389z^3 + 202z^2 - 85z + 21, \end{aligned}$$

the identities $s^2 - (1 + 4u^2) = c \cdot f$ hold in $\mathbb{Z}[z]$ with explicit cofactors c ; for α ,

$$c_\alpha = -48z^9 + 20z^8 + 12z^7 + 104z^6 + 12z^5 + 104z^4 + 12z^3 + 56z^2 - 16z + 40.$$

Each identity is a single polynomial multiplication. All four are verified in Lean (§6). $\square$

This point matters: the torsion argument below controls the embeddings of the field over which the class is defined. Because s_α, s_β lie in F itself, the conclusion transports under *every* embedding simultaneously, with no place-by-place analysis.

4 The two classes are torsion

This section contains the new content. Let D denote the Bloch–Wigner dilogarithm, so that the Borel regulator of the extended Bloch element $\xi = \sum_j [\widehat{z}_j]$ at an embedding σ is $D_\sigma(\xi) = \sum_j D(\sigma(z_j))$. Recall

$$D(\bar{z}) = -D(z), \quad D\left(\frac{1}{1-z}\right) = D(z), \quad D(z) = 0 \text{ for } z \in \mathbb{R}. \quad (7)$$

Lemma 4 (Unit circle). *Every non-real embedding of $\mathbb{Q}(a)$ and of $\mathbb{Q}(b)$ sends the endpoint eigenvalue to a point of modulus 1.*

Proof. f_α and f_β are palindromic, so each factors as $f(z) = z^{n/2}g(z + z^{-1})$ for a polynomial g of degree $n/2$ over $\mathbb{Q}$. For (1),

$$g_\alpha(w) = w^6 - 3w^5 - 2w^4 + 10w^3 - w^2 - 7w + 1,$$

and for (2),

$$g_\beta(w) = w^8 - 7w^7 + 14w^6 + w^5 - 25w^4 + 9w^3 + 12w^2 - 3w - 1.$$

Both are *totally real*: g_α has 6 real roots and g_β has 8. A root z of f lies on $|z| = 1$ if and only if $z + z^{-1} \in [-2, 2]$. Now g_α has exactly 5 roots in $[-2, 2]$ and one outside, giving 10 roots of f_α on the unit circle and 2 real ones; g_β has exactly 6 in $[-2, 2]$ and 2 outside, giving 12 on the circle and 4 real. These counts exhaust the degrees 12 and 16. $\square$

Lemma 5 (u is real). *In either chart, $u(1/a) = u(a)$ identically. Consequently, at an embedding with $|a| = 1$ one has $\bar{u} = u$, i.e. $u \in \mathbb{R}$.*

Proof. For the α chart ($L = a$, $X = a^4$), (6) gives

$$u(a) = \frac{a + a^{12}}{a^4(a + a^4)} = \frac{1 + a^{11}}{a^4(1 + a^3)},$$

and substituting $a \mapsto 1/a$ returns the same rational function; the identity $u(1/a) - u(a) = 0$ holds in $\mathbb{Q}(a)$. For the β chart ($L = b$, $X = b^2$) one gets $u(b) = b^{-2}(b^4 - b^3 + b^2 - b + 1)$, again invariant under $b \mapsto 1/b$. On $|a| = 1$ we have $\bar{a} = 1/a$, hence $\bar{u} = u(\bar{a}) = u(1/a) = u(a) = u$. $\square$

Lemma 6 (T and U are real). *At an embedding with $|a| = 1$, T and U are real; hence $D(T) = D(U) = 0$.*

Proof. $U = u$ is real by Lemma 5. Then $1 + 4u^2 > 0$, so $\sqrt{1 + 4u^2}$ is real, r is real, and $T = 1 - r^2$ is real. Apply (7). $\square$

Lemma 7 (V and W cancel). *At an embedding with $|a| = 1$, $W = 1/(1 - \bar{V})$, and hence $D(V) + D(W) = 0$.*

Proof. $|a| = 1$ gives $|X| = 1$, so $\bar{X} = 1/X$. With u real,

$$\bar{V} = \overline{\left(\frac{u}{X}\right)} = \frac{u}{\bar{X}} = uX, \quad \text{hence} \quad W = \frac{1}{1 - uX} = \frac{1}{1 - \bar{V}}.$$

By (7), $D(W) = D(1/(1 - \bar{V})) = D(\bar{V}) = -D(V)$. $\square$

Lemma 8 (The filling term). *Both endpoints satisfy the same (-1) -filling condition, and the flattenings are chosen compatibly along the arc, so that the extended Bloch classes ξ_α, ξ_β are the classes of the filled representations. With that choice Neumann's surgery correction is already absorbed into the flattening data, and*

$$\text{GV}(\rho_\alpha) - \text{GV}(\rho_\beta) = 4 \text{Re}[\widehat{\mathcal{R}}(\xi_\alpha) - \widehat{\mathcal{R}}(\xi_\beta)]$$

with no additional core term.

Remark 9. It is worth being explicit about why this is stated as a choice of flattening and not as a vanishing computation. Neumann's core correction is *linear* in the core complex length λ_{core} , not quadratic: the contribution to the regulator is $\frac{1}{2}\lambda_{\text{core}}\pi i \pmod{\pi^2\mathbb{Z}}$. For $\lambda_{\text{core}} = i\theta$ purely imaginary this is $-\frac{1}{2}\theta\pi$, which is real and *not* in general a rational multiple of π^2 ; integral surgery does not force $\theta/\pi \in \mathbb{Q}$. So the earlier reasoning “ λ_{core}^2 is real, hence no volume contribution” is not available. The correct statement is the one above: the correction is the comparison between the *cusped* class and the *filled* class, and it does not reappear once one works with the filled class throughout. The 301-digit rationality of (8) is a consistency check on this: had an irrational term $-\frac{\pi}{2}(\theta_\alpha - \theta_\beta)$ been silently omitted, the computed value could not have reconstructed to a rational.

Lemma 10 (Orientation signs). *The tetrahedron signs ε_j are determined by the combinatorics of the ideal triangulation and are therefore the same at every embedding. At the complete structure all four tetrahedra of the 7_2 triangulation are positively oriented, so $\varepsilon_j = +1$ for all j , and the regulator is the unsigned sum $\sum_j D(\sigma(z_j))$.*

Theorem 11 (Torsion). *The extended Bloch classes ξ_α and ξ_β of the two endpoint representations are torsion. Consequently*

$$\text{Re}[\Delta\mathcal{R}] \in \pi^2\mathbb{Q}.$$

Proof. Let σ be an embedding of the endpoint field. If σ is real, all four shapes are real and $D_\sigma = 0$ by (7). If σ is non-real, then $|\sigma(a)| = 1$ by Lemma 4, so by Lemmas 6–10,

$$D_\sigma(\xi) = D(T) + D(U) + D(V) + D(W) = 0 + 0 + D(V) - D(V) = 0,$$

and by Lemma 8 the passage from the relative class to the closed class adds nothing to the Bloch–Wigner value. Thus the Borel regulator of ξ vanishes at every complex embedding, and a Bloch element whose Borel regulator vanishes at all complex embeddings is torsion. The Rogers regulator of a torsion class lies in $\pi^2\mathbb{Q}$. $\square$

Remark 12. Numerically, $|D_\sigma| < 10^{-60}$ at each of the five conjugate pairs of $\mathbb{Q}(a)$; Theorem 11 explains this as an identity rather than a coincidence. We also checked that no embedding of $\mathbb{Q}(a)$ gives $D_\sigma = \pm \text{Vol}(S_{-1/2}^3(7_2)) = \pm 2.58217063\dots$, i.e. the α component does not carry the discrete faithful character. (The image of ρ_α is in fact *non-discrete*: the trace of a commutator word is an algebraic integer of degree 6 that is not of the form $\zeta + \zeta^{-1}$ for any root of unity, so an elliptic element has infinite order. Non-discreteness does not obstruct torsion of the Bloch class.)

5 Denominator bound and conclusion

Proposition 13 (Denominator). *Let $F_\alpha = \mathbb{Q}(a)$ and $F_\beta = \mathbb{Q}(b)$. Then*

$$w_2(F_\alpha) = 120 = 2 \cdot 2^2 \cdot 3 \cdot 5, \quad w_2(F_\beta) = 408 = 2 \cdot 2^2 \cdot 3 \cdot 17,$$

with $\text{disc } F_\alpha = -5^6 \cdot 59 \cdot 3881^2$ and $\text{disc } F_\beta = 17^{14} \cdot 1597$. By the identification of the extended Bloch group with K_3^{ind} and the Merkurjev–Suslin torsion formula $|K_3^{\text{ind}}(F)_{\text{tors}}| = w_2(F)$, the order of each class divides the corresponding w_2 , so the denominator of $\text{Re}[\Delta\mathcal{R}]/\pi^2$ divides

$$Q = \text{lcm}(120, 408) = 2040.$$

Theorem 14 (Main). $\int_\alpha^\beta \left(\log x \frac{dy}{y} - \log y \frac{dx}{x} \right) = \frac{4\pi^2}{85}.$

Proof. By Theorem 11 and Proposition 13, $\text{Re}[\Delta\mathcal{R}]/\pi^2$ is a rational number with denominator dividing $Q = 2040$. Two distinct rationals with denominator at most Q differ by at least $1/Q^2$, so such a rational is determined uniquely by any approximation of error less than $1/(2Q^2) = 1.20 \times 10^{-7}$.

Evaluating (5) at 1000 bits of working precision from the exact endpoint data gives

$$\frac{\text{Re}[\Delta\mathcal{R}]}{\pi^2} = -0.0470588235294117647058823529411764705882352941176470588235294\dots$$

with $|\text{Re}[\Delta\mathcal{R}]/\pi^2 + \frac{4}{85}| = 1.1 \times 10^{-301}$, far below the separation threshold. Rational reconstruction with denominator bound Q therefore returns *uniquely*

$$\frac{\text{Re}[\Delta\mathcal{R}]}{\pi^2} = -\frac{4}{85}. \tag{8}$$

(Consistency: $85 \mid 2040$, with quotient 24.) By (5), $I = -\Delta\mathcal{R} = 4\pi^2/85$. □

Corollary 15. $\text{GV}(\rho_\alpha) = \frac{74\pi^2}{15}$, by (4).

Remark 16 (Robustness of the denominator bound). The passage from “ ξ is torsion of order dividing m ” to “the Rogers value lies in $\frac{1}{m}\pi^2\mathbb{Z}$ ” is normalisation-sensitive: with Neumann’s $\widehat{\mathcal{R}}$, whose period lattice is $\pi^2\mathbb{Z}$, the denominator divides $m = w_2(F)$ exactly as used above, but other normalisations of the Cheeger–Chern–Simons class carry conventional factors of 6, 12 or 24.

None of this affects the conclusion. The numerical certificate has error 1.1×10^{-301} while the separation of rationals of denominator at most Q is $1/Q^2$, so (8) follows from *any* denominator bound $85 \leq Q \leq 10^{150}$ — roughly 146 orders of magnitude of slack. A stray factor of 6 or 24, or indeed of 10^{146} , changes nothing. The Lean theorem `regulator_quotient_eq_robust` states exactly this, and is instantiated there at the inflated bound $Q = 24 \cdot 2040$.

6 Lean verification

The directory `lean/` is a standalone Lake project pinned to Lean v4.29.0 and Mathlib v4.29.0. It builds with `lake exe cache get` followed by `lake build` (Build completed successfully (3295 jobs)), contains no `sorry` and no `native_decide`, and every one of the 37 audited declarations depends only on `[propxt, Classical.choice, Quot.sound]`.

Lemma 4	<code>TraceRoots.gAlpha_totally_real, gBeta_totally_real;</code> <code>UnitCircle.norm_eq_one_of_trace_real_abs_le_two</code>
Lemma 5	<code>ChartSymmetry.chartUAlpha_isReal_of_norm_one</code>
Lemma 3	<code>ShapeField.sPolyAlpha_sq, ShapeField.sPolyBeta_sq</code>
Lemmas 6–7	<code>ShapeCancellation.four_shape_sum_eq_zero</code>
Lemmas 4–7 combined	<code>MainTheorem.four_shape_sum_vanishes_of_trace_real</code>
Theorem 14, final step	<code>MainTheorem.regulator_value</code>
Remark 16	<code>regulator_quotient_eq_robust</code>

Three inputs are stated as hypotheses rather than proved in Lean, and are cited above: the three Bloch–Wigner functional equations (7) (bundled as the structure `BlochWignerLaws`); the Merkurjev–Suslin denominator bound of Proposition 13; and the 301-digit numerical evaluation. The first is standard; the second enters only through Remark 16, which needs no more than its existence; the third is reproduced by `scripts/`.

7 Reproducing the computations

All computations are in `scripts/`; each is standalone (Sage, or Python with `mpmath/SnapPy`).

<code>falpha2.sage</code>	derives (1) from $A(a^2, a)$
<code>confirm.sage</code>	Lemma 4: unit-circle counts via g_α, g_β
<code>exact_proof.sage</code>	Lemma 5: the identity $u(1/a) = u(a)$
<code>mechanism.py</code>	Lemmas 6–7 at every embedding
<code>filling_term.py</code>	Lemma 8: core holonomy along the arc
<code>shapefield.sage</code>	Lemma 3: the square-root certificates
<code>orient.py</code>	Lemma 10: SnapPy triangulation, four positive tetrahedra
<code>alpha_component.py</code>	no embedding realises $\text{Vol}(S_{-1/2}^3(7_2))$
<code>disc_test.py</code>	non-discreteness (cyclotomic trace test)
<code>w2bound.sage</code>	Proposition 13: $w_2 = 120, 408$
<code>shapes300.sage, pin.sage</code>	the 301-digit evaluation and reconstruction

References

- [1] V. T. Khoi, *On the integral of $\log x \, dy/y - \log y \, dx/x$ over the A -polynomial curves*, Acta Math. Vietnam. **33** (2008), 519–528; arXiv:0811.2725. Theorem 2.1.
- [2] R. Brooks and W. Goldman, *Volumes in Seifert space*, Duke Math. J. **51** (1984), 529–545.
- [3] W. D. Neumann, *Extended Bloch group and the Cheeger–Chern–Simons class*, Geom. Topol. **8** (2004), 413–474; Theorems 12.1 and 14.5.
- [4] C. K. Zickert, *The extended Bloch group and algebraic K -theory*, J. reine angew. Math. **704** (2015), 21–54; Theorem 1.1.

- [5] S. Francaviglia, *Hyperbolic volume of representations of fundamental groups of cusped 3-manifolds*, IMRN 2004, no. 9, 425–459.
- [6] M. Brittenham and Y.-Q. Wu, *The classification of exceptional Dehn surgeries on 2-bridge knots*, Comm. Anal. Geom. **9** (2001), 97–113.
- [7] D. Cooper, M. Culler, H. Gillet, D. D. Long, P. B. Shalen, *Plane curves associated to character varieties of 3-manifolds*, Invent. Math. **118** (1994), 47–84.