

Tromino Tiling

Tromino Tiling

- L-trominoes can be used to tile a “punctured” $2^n \times 2^n$ grid (punctured = one cell removed), for all positive integers n



Tromino Tiling

- L-trominoes can be used to tile a “punctured” $2^n \times 2^n$ grid (punctured = one cell removed), for all positive integers n



Tromino Tiling

- L-trominoes can be used to tile a “punctured” $2^n \times 2^n$ grid (punctured = one cell removed), for all positive integers n



Tromino Tiling

- L-trominoes can be used to tile a “punctured” $2^n \times 2^n$ grid (punctured = one cell removed), for all positive integers n



Tromino Tiling


- L-trominoes can be used to tile a “punctured” $2^n \times 2^n$ grid (punctured = one cell removed), for all positive integers n
- Base case: $n=1$



Tromino Tiling

- L-trominoes can be used to tile a “punctured” $2^n \times 2^n$ grid (punctured = one cell removed), for all positive integers n

- Base case: $n=1$




Tromino Tiling

- L-trominoes can be used to tile a “punctured” $2^n \times 2^n$ grid (punctured = one cell removed), for all positive integers n



- Base case: $n=1$




Tromino Tiling

- L-trominoes can be used to tile a “punctured” $2^n \times 2^n$ grid (punctured = one cell removed), for all positive integers n



- Base case: $n=1$



- Inductive step: For all integers $k \geq 1$:

Hypothesis: true for $n=k$


To prove: true for $n=k+1$

Tromino Tiling

- L-trominoes can be used to tile a “punctured” $2^n \times 2^n$ grid (punctured = one cell removed), for all positive integers n



- Base case: $n=1$



- Inductive step: For all integers $k \geq 1$:

Hypothesis: true for $n=k$

To prove: true for $n=k+1$


- Idea: can partition the $2^{k+1} \times 2^{k+1}$ punctured grid into four $2^k \times 2^k$ punctured grids, plus a tromino. Each of these can be tiled using trominoes (by inductive hypothesis).

Tromino Tiling

- L-trominoes can be used to tile a “punctured” $2^n \times 2^n$ grid (punctured = one cell removed), for all positive integers n



- Base case: $n=1$



- Inductive step: For all integers $k \geq 1$:

Hypothesis: true for $n=k$

To prove: true for $n=k+1$


- Idea: can partition the $2^{k+1} \times 2^{k+1}$ punctured grid into four $2^k \times 2^k$ punctured grids, plus a tromino. Each of these can be tiled using trominoes (by inductive hypothesis).
- Actually gives a (recursive) algorithm for tiling

Tromino Tiling

- L-trominoes can be used to tile a “punctured” $2^n \times 2^n$ grid (punctured = one cell removed), for all positive integers n




- Base case: $n=1$



- Inductive step: For all integers $k \geq 1$:

Hypothesis: true for $n=k$

To prove: true for $n=k+1$




- Idea: can partition the $2^{k+1} \times 2^{k+1}$ punctured grid into four $2^k \times 2^k$ punctured grids, plus a tromino. Each of these can be tiled using trominoes (by inductive hypothesis).
- Actually gives a (recursive) algorithm for tiling

Tromino Tiling

- L-trominoes can be used to tile a “punctured” $2^n \times 2^n$ grid (punctured = one cell removed), for all positive integers n




- Base case: $n=1$



- Inductive step: For all integers $k \geq 1$:

Hypothesis: true for $n=k$

To prove: true for $n=k+1$



- Idea: can partition the $2^{k+1} \times 2^{k+1}$ punctured grid into four $2^k \times 2^k$ punctured grids, plus a tromino. Each of these can be tiled using trominoes (by inductive hypothesis).


- Actually gives a (recursive) algorithm for tiling

Tromino Tiling

- L-trominoes can be used to tile a “punctured” $2^n \times 2^n$ grid (punctured = one cell removed), for all positive integers n




- Base case: $n=1$



- Inductive step: For all integers $k \geq 1$:

Hypothesis: true for $n=k$

To prove: true for $n=k+1$




- Idea: can partition the $2^{k+1} \times 2^{k+1}$ punctured grid into four $2^k \times 2^k$ punctured grids, plus a tromino. Each of these can be tiled using trominoes (by inductive hypothesis).
- Actually gives a (recursive) algorithm for tiling

Tromino Tiling

- L-trominoes can be used to tile a “punctured” $2^n \times 2^n$ grid (punctured = one cell removed), for all positive integers n




- Base case: $n=1$



- Inductive step: For all integers $k \geq 1$:

Hypothesis: true for $n=k$

To prove: true for $n=k+1$




- Idea: can partition the $2^{k+1} \times 2^{k+1}$ punctured grid into four $2^k \times 2^k$ punctured grids, plus a tromino. Each of these can be tiled using trominoes (by inductive hypothesis).
- Actually gives a (recursive) algorithm for tiling

Tromino Tiling

- L-trominoes can be used to tile a “punctured” $2^n \times 2^n$ grid (punctured = one cell removed), for all positive integers n




- Base case: $n=1$



- Inductive step: For all integers $k \geq 1$:

Hypothesis: true for $n=k$

To prove: true for $n=k+1$




- Idea: can partition the $2^{k+1} \times 2^{k+1}$ punctured grid into four $2^k \times 2^k$ punctured grids, plus a tromino. Each of these can be tiled using trominoes (by inductive hypothesis).
- Actually gives a (recursive) algorithm for tiling

Tromino Tiling

- L-trominoes can be used to tile a “punctured” $2^n \times 2^n$ grid (punctured = one cell removed), for all positive integers n





- Base case: $n=1$



- Inductive step: For all integers $k \geq 1$:

Hypothesis: true for $n=k$

To prove: true for $n=k+1$




- Idea: can partition the $2^{k+1} \times 2^{k+1}$ punctured grid into four $2^k \times 2^k$ punctured grids, plus a tromino. Each of these can be tiled using trominoes (by inductive hypothesis).
- Actually gives a (recursive) algorithm for tiling

Tromino Tiling

- L-trominoes can be used to tile a “punctured” $2^n \times 2^n$ grid (punctured = one cell removed), for all positive integers n





- Base case: $n=1$



- Inductive step: For all integers $k \geq 1$:

Hypothesis: true for $n=k$

To prove: true for $n=k+1$



- Idea: can partition the $2^{k+1} \times 2^{k+1}$ punctured grid into four $2^k \times 2^k$ punctured grids, plus a tromino. Each of these can be tiled using trominoes (by inductive hypothesis).


- Actually gives a (recursive) algorithm for tiling

Tromino Tiling

- L-trominoes can be used to tile a “punctured” $2^n \times 2^n$ grid (punctured = one cell removed), for all positive integers n





- Base case: $n=1$



- Inductive step: For all integers $k \geq 1$:

Hypothesis: true for $n=k$

To prove: true for $n=k+1$




- Idea: can partition the $2^{k+1} \times 2^{k+1}$ punctured grid into four $2^k \times 2^k$ punctured grids, plus a tromino. Each of these can be tiled using trominoes (by inductive hypothesis).
- Actually gives a (recursive) algorithm for tiling

Tromino Tiling

- L-trominoes can be used to tile a “punctured” $2^n \times 2^n$ grid (punctured = one cell removed), for all positive integers n





- Base case: $n=1$



- Inductive step: For all integers $k \geq 1$:

Hypothesis: true for $n=k$

To prove: true for $n=k+1$




- Idea: can partition the $2^{k+1} \times 2^{k+1}$ punctured grid into four $2^k \times 2^k$ punctured grids, plus a tromino. Each of these can be tiled using trominoes (by inductive hypothesis).
- Actually gives a (recursive) algorithm for tiling

Tromino Tiling

- L-trominoes can be used to tile a “punctured” $2^n \times 2^n$ grid (punctured = one cell removed), for all positive integers n





- Base case: $n=1$



- Inductive step: For all integers $k \geq 1$:

Hypothesis: true for $n=k$

To prove: true for $n=k+1$




- Idea: can partition the $2^{k+1} \times 2^{k+1}$ punctured grid into four $2^k \times 2^k$ punctured grids, plus a tromino. Each of these can be tiled using trominoes (by inductive hypothesis).
- Actually gives a (recursive) algorithm for tiling

Tromino Tiling

- L-trominoes can be used to tile a “punctured” $2^n \times 2^n$ grid (punctured = one cell removed), for all positive integers n





- Base case: $n=1$




- Inductive step: For all integers $k \geq 1$:

Hypothesis: true for $n=k$

To prove: true for $n=k+1$



- Idea: can partition the $2^{k+1} \times 2^{k+1}$ punctured grid into four $2^k \times 2^k$ punctured grids, plus a tromino. Each of these can be tiled using trominoes (by inductive hypothesis).




- Actually gives a (recursive) algorithm for tiling

Tromino Tiling

- L-trominoes can be used to tile a “punctured” $2^n \times 2^n$ grid (punctured = one cell removed), for all positive integers n





- Base case: $n=1$




- Inductive step: For all integers $k \geq 1$:

Hypothesis: true for $n=k$

To prove: true for $n=k+1$



- Idea: can partition the $2^{k+1} \times 2^{k+1}$ punctured grid into four $2^k \times 2^k$ punctured grids, plus a tromino. Each of these can be tiled using trominoes (by inductive hypothesis).




- Actually gives a (recursive) algorithm for tiling

Tromino Tiling

- L-trominoes can be used to tile a “punctured” $2^n \times 2^n$ grid (punctured = one cell removed), for all positive integers n





- Base case: $n=1$




- Inductive step: For all integers $k \geq 1$:

Hypothesis: true for $n=k$

To prove: true for $n=k+1$



- Idea: can partition the $2^{k+1} \times 2^{k+1}$ punctured grid into four $2^k \times 2^k$ punctured grids, plus a tromino. Each of these can be tiled using trominoes (by inductive hypothesis).




- Actually gives a (recursive) algorithm for tiling

Tromino Tiling

- L-trominoes can be used to tile a “punctured” $2^n \times 2^n$ grid (punctured = one cell removed), for all positive integers n





- Base case: $n=1$




- Inductive step: For all integers $k \geq 1$:

Hypothesis: true for $n=k$

To prove: true for $n=k+1$



- Idea: can partition the $2^{k+1} \times 2^{k+1}$ punctured grid into four $2^k \times 2^k$ punctured grids, plus a tromino. Each of these can be tiled using trominoes (by inductive hypothesis).




- Actually gives a (recursive) algorithm for tiling

Tromino Tiling

- L-trominoes can be used to tile a “punctured” $2^n \times 2^n$ grid (punctured = one cell removed), for all positive integers n





- Base case: $n=1$




- Inductive step: For all integers $k \geq 1$:

Hypothesis: true for $n=k$

To prove: true for $n=k+1$



- Idea: can partition the $2^{k+1} \times 2^{k+1}$ punctured grid into four $2^k \times 2^k$ punctured grids, plus a tromino. Each of these can be tiled using trominoes (by inductive hypothesis).



- Actually gives a (recursive) algorithm for tiling