

# Real-Time Computability of Real Numbers by Chemical Reaction Networks

---

Xiang Huang<sup>1</sup>, Titus H. Klinge<sup>2</sup>, James I. Lathrop<sup>1</sup>,  
Xiaoyuan Li<sup>1</sup>, Jack H. Lutz<sup>1</sup>

April 21, 2018

<sup>1</sup>Computer Science Department, Iowa State University

<sup>2</sup>Computer Science Department, Grinnell College

# Introduction and Motivation

---

- Abbreviated “CRNs”

# Chemical Reaction Networks

- Abbreviated “CRNs”
- An abstract **mathematical model** of how chemicals interact in a well-mixed solution.

# Chemical Reaction Networks

- Abbreviated “CRNs”
- An abstract **mathematical model** of how chemicals interact in a well-mixed solution.
- Used for over a half century

# Chemical Reaction Networks

- Abbreviated “CRNs”
- An abstract **mathematical model** of how chemicals interact in a well-mixed solution.
- Used for over a half century


## Hydrazine Combustion




# Chemical Reaction Networks

- Abbreviated “CRNs”
- An abstract **mathematical model** of how chemicals interact in a well-mixed solution.
- Used for over a half century

## Hydrazine Combustion



## Abstract CRN



- CRNs have become the programming language of choice for many molecular programming applications.



# Computing Real Numbers

- CRNs have become the programming language of choice for many molecular programming applications.
- How powerful are they?

- CRNs have become the programming language of choice for many molecular programming applications.
- How powerful are they?
- Benchmark: we are asking which real numbers can be computed by CRNs.

# Computing Real Numbers

- CRNs have become the programming language of choice for many molecular programming applications.
- How powerful are they?
- Benchmark: we are asking which real numbers can be computed by CRNs.
- In fact, which real numbers can be computed **quickly by CRNs**?

## Today's Theorem

*Every algebraic number can be computed by a CRN in real time.*

## Today's Theorem

*Every algebraic number can be computed by a CRN in real time.*

**algebraic**

$$\sqrt{2}, \frac{1+\sqrt{5}}{2}, \frac{\sqrt{\sqrt{3}+\sqrt{5}}}{2}$$

**not algebraic (transcendental)**

$$\pi, e$$

- Real-time Computability

- Real-time Computability
- Lyapunov Computability

- Real-time Computability
- Lyapunov Computability
- Algebraic Numbers




- Real-time Computability
- Lyapunov Computability
- Algebraic Numbers
- Large Population Protocol
- Transcendental Number

# Real-time Computability

---

## More about CRNs



$A, B, C, X$  are **species** (abstract molecule types).  $r, s$  are **rate constants**. The **rates** of these two reactions at time  $t$  are

$$r \cdot a(t) \cdot b(t)^2$$

and


$$s.$$

where  $a(t)$  = the **concentration** of  $A$  at time  $t$ , etc.

# CRN and its ordinary differential equations


Once we have the definition for rates of reactions, we can write a CRN into a system of ordinary differential equations (ODEs).

**Example:**




with initial value  $x(0) = 0, y(0) = 0$ .

# Plot




# Streamplot



**Figure 1:** Streamplot

# Streamplot




**Figure 1:** Streamplot



**Figure 2:** duck

# Streamplot



**Figure 1:** Streamplot



**Figure 2:** duck

Put a rubber duck at point  $(0,0)$  at time 0. It will eventually reach the red spot, i.e.,  $(2, \frac{1}{2})$ .



## Real-time computability: the idea

We view the long-term behavior (limit) of a CRN as a process of **computation**.

## Real-time computability: the idea

We view the long-term behavior (limit) of a CRN as a process of **computation**. In the previous example, we say

- $X$  computes 2

## Real-time computability: the idea


We view the long-term behavior (limit) of a CRN as a process of **computation**. In the previous example, we say

- $X$  computes 2
- $Y$  computes  $\frac{1}{2}$

# Real-time computability: the idea

We view the long-term behavior (limit) of a CRN as a process of **computation**. In the previous example, we say

- $X$  computes 2
- $Y$  computes  $\frac{1}{2}$



## Real-time computability: the definition

A number  $\alpha$  is real-time computable, if it can be computed by a CRN  $N$  with the following properties:

- (*zero initialized*). All species should have initial value 0.

# Real-time computability: the definition

A number  $\alpha$  is real-time computable, if it can be computed by a CRN  $N$  with the following properties:

- (*zero initialized*). All species should have initial value 0.
- (*integrality*). All the rate constants of any reaction should be integers.

## Real-time computability: the definition

A number  $\alpha$  is real-time computable, if it can be computed by a CRN  $N$  with the following properties:

- (*zero initialized*). All species should have initial value 0.
- (*integrality*). All the rate constants of any reaction should be integers.
- (*boundedness*). All the concentrations of any species are bounded.

# Real-time computability: the definition

A number  $\alpha$  is real-time computable, if it can be computed by a CRN  $N$  with the following properties:


- (*zero initialized*). All species should have initial value 0.
- (*integrality*). All the rate constants of any reaction should be integers.
- (*boundedness*). All the concentrations of any species are bounded.
- (*real-time convergence*). There is a species  $X$ , such that for all  $t \in [\tau, \infty)$  for some  $\tau$ ,

$$|x(t) - |\alpha|| \leq 2^{-t} \tag{1}$$



- This prevents the CRN from coding too much (infinite) information in its initial values.  
For example, encode an uncomputable number in the initial value to “compute” it.

Suppose we want to compute a real number  $\alpha$ , without the constraint, one can use the following simple CRN



to compute  $\alpha$ , which is just like “cheating”.

We denote the set of real-time computable reals as  $\mathbb{R}_{RTCRN}$ .

The following are true for this class:

We denote the set of real-time computable reals as  $\mathbb{R}_{RTCRN}$ .

The following are true for this class:

- $\mathbb{R}_{RTCRN}$  is a countable set.

We denote the set of real-time computable reals as  $\mathbb{R}_{RTCRN}$ .

The following are true for this class:

- $\mathbb{R}_{RTCRN}$  is a countable set.
- $\mathbb{Q} \subseteq \mathbb{R}_{RTCRN}$ , where  $\mathbb{Q}$  is the set of rational numbers.

We denote the set of real-time computable reals as  $\mathbb{R}_{RTCRN}$ .

The following are true for this class:

- $\mathbb{R}_{RTCRN}$  is a countable set.
- $\mathbb{Q} \subseteq \mathbb{R}_{RTCRN}$ , where  $\mathbb{Q}$  is the set of rational numbers.

# Lyapunov CRN-Computable Reals: A Bridge

---

# A Bridge Crosses Nonlinearity

- It is not easy to show real-time computability directly, due to the high nonlinearity of CRNs.

---

<sup>1</sup>Teschl, Gerald. *Ordinary Differential Equations and Dynamical Systems*.



# A Bridge Crosses Nonlinearity

- It is not easy to show real-time computability directly, due to the high nonlinearity of CRNs.
- However, we can take advantage of Lyapunov's **exponential stability theorem**<sup>1</sup>, which requires only derivative and eigenvalue analysis.

---

<sup>1</sup>Teschl, Gerald. *Ordinary Differential Equations and Dynamical Systems*.

# A Bridge Crosses Nonlinearity

- It is not easy to show real-time computability directly, due to the high nonlinearity of CRNs.
- However, we can take advantage of Lyapunov's **exponential stability theorem**<sup>1</sup>, which requires only derivative and eigenvalue analysis.
- The path: instead of showing

$$\mathbf{Algebraic} \subseteq \mathbb{R}_{RTCRN},$$

we define a new set  $\mathbb{R}_{LCRN}$  in between and show

$$\mathbf{Algebraic} \subseteq \mathbb{R}_{LCRN} \subseteq \mathbb{R}_{RTCRN},$$

---

<sup>1</sup>Teschl, Gerald. *Ordinary Differential Equations and Dynamical Systems*.

If  $N = (S, R)$  is a CRN with  $S = \{Y_1, \dots, Y_n\}$ , then the ODEs of each species can be written as

$$\begin{aligned}y_1' &= f_1(y_1, \dots, y_n), \\ &\vdots \\ y_n' &= f_n(y_1, \dots, y_n),\end{aligned}$$

where  $f_1, \dots, f_n : \mathbb{R}^n \rightarrow \mathbb{R}$  are polynomials.

# Jacobian Matrix

If  $N = (S, R)$  is a CRN with  $S = \{Y_1, \dots, Y_n\}$ , then the ODEs of each species can be written as

$$\begin{aligned}y_1' &= f_1(y_1, \dots, y_n), \\&\vdots \\y_n' &= f_n(y_1, \dots, y_n),\end{aligned}$$

where  $f_1, \dots, f_n : \mathbb{R}^n \rightarrow \mathbb{R}$  are polynomials.

The **Jacobian matrix** of the CRN  $N$  is the  $n \times n$  matrix

$$J_N = \begin{pmatrix} \frac{\partial f_1}{\partial y_1} & \cdots & \frac{\partial f_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial f_n}{\partial y_1} & \cdots & \frac{\partial f_n}{\partial y_n} \end{pmatrix}.$$

## The Bridge: $\mathbb{R}_{LCRN}$

We define the set of real  $\mathbb{R}_{LCRN}$  basically like  $\mathbb{R}_{RTCNR}$ ., but instead of requiring exponential convergence, we require:

- $x \in \mathbb{R}_{LCRN}$  then all eigenvalues of the Jacobian at  $x$  must have **negative real part**.
- $x$  is a limit point of the system (in our rubber duck sense).

Lyapunov's exponential stability theorem then guarantees exponential convergence **near  $x$** .

## The Bridge: $\mathbb{R}_{LCRN}$

We define the set of real  $\mathbb{R}_{LCRN}$  basically like  $\mathbb{R}_{RTCRN}$ :, but instead of requiring exponential convergence, we require:

- $x \in \mathbb{R}_{LCRN}$  then all eigenvalues of the Jacobian at  $x$  must have **negative real part**.
- $x$  is a limit point of the system (in our rubber duck sense).

Lyapunov's exponential stability theorem then guarantees exponential convergence **near  $x$** .



**Figure 3:** whirlpool

# Algebraic Numbers

---

A real number  $\alpha \in \mathbb{R}$  is **algebraic** if there is a polynomial  $P(x)$  with integer coefficients such that  $\alpha$  is a root of  $P$ , i.e.,  $P(\alpha) = 0$ . Here we demonstrate by an example of how to compute algebraic numbers. Actually, along this path, one can show that algebraic numbers are in  $\mathbb{R}_{LCRN}$ .

**Example:**  $\alpha = 2 - \sqrt{3}$  is a root of

$$x^2 - 4x + 1 = 0$$



A real number  $\alpha \in \mathbb{R}$  is **algebraic** if there is a polynomial  $P(x)$  with integer coefficients such that  $\alpha$  is a root of  $P$ , i.e.,  $P(\alpha) = 0$ . Here we demonstrate by an example of how to compute algebraic numbers. Actually, along this path, one can show that algebraic numbers are in  $\mathbb{R}_{LCRN}$ .

**Example:**  $\alpha = 2 - \sqrt{3}$  is a root of

$$x^2 - 4x + 1 = 0$$

We can implement a CRN such that:

$$x' = x^2 - 4x + 1 \quad \Longleftrightarrow$$

## Algebraic are in $\mathbb{R}_{NECRN}$

A real number  $\alpha \in \mathbb{R}$  is **algebraic** if there is a polynomial  $P(x)$  with integer coefficients such that  $\alpha$  is a root of  $P$ , i.e.,  $P(\alpha) = 0$ . Here we demonstrate by an example of how to compute algebraic numbers. Actually, along this path, one can show that algebraic numbers are in  $\mathbb{R}_{LCRN}$ .

**Example:**  $\alpha = 2 - \sqrt{3}$  is a root of

$$x^2 - 4x + 1 = 0$$

We can implement a CRN such that:



## Algebraic are in $\mathbb{R}_{NECRN}$

A real number  $\alpha \in \mathbb{R}$  is **algebraic** if there is a polynomial  $P(x)$  with integer coefficients such that  $\alpha$  is a root of  $P$ , i.e.,  $P(\alpha) = 0$ . Here we demonstrate by an example of how to compute algebraic numbers. Actually, along this path, one can show that algebraic numbers are in  $\mathbb{R}_{LCRN}$ .

**Example:**  $\alpha = 2 - \sqrt{3}$  is a root of

$$x^2 - 4x + 1 = 0$$

We can implement a CRN such that:



Note that in this process:

- $\alpha$  is the **first positive root** of the polynomial.

## Algebraic are in $\mathbb{R}_{NECRN}$

A real number  $\alpha \in \mathbb{R}$  is **algebraic** if there is a polynomial  $P(x)$  with integer coefficients such that  $\alpha$  is a root of  $P$ , i.e.,  $P(\alpha) = 0$ . Here we demonstrate by an example of how to compute algebraic numbers. Actually, along this path, one can show that algebraic numbers are in  $\mathbb{R}_{LCRN}$ .

**Example:**  $\alpha = 2 - \sqrt{3}$  is a root of

$$x^2 - 4x + 1 = 0$$


We can implement a CRN such that:



Note that in this process:

- $\alpha$  is the **first positive root** of the polynomial.
- constant term i.e.,  $P(0)$ , of the polynomial should be positive.

# First positive root




**Figure 4:** First positive root

## Other roots: shift

What if we want to compute the second root,  $\beta = 2 + \sqrt{3}$ ?


## Other roots: shift

What if we want to compute the second root,  $\beta = 2 + \sqrt{3}$ ?



**Figure 5:** Other roots

## Other roots: shift and flip



**Figure 6:** Other roots



Let species  $X$  computes  $\alpha$ . How to compute  $\frac{1}{\alpha}$ ?

Let species  $X$  computes  $\alpha$ . How to compute  $\frac{1}{\alpha}$ ?

Ans:  $Y' = 1 - XY$

Let species  $X$  computes  $\alpha$ . How to compute  $\frac{1}{\alpha}$ ?

Ans:  $Y' = 1 - XY$

That is, we add the follows reaction to the original network.



# Subtraction

Let  $X$  computes  $\alpha$ ,  $Y$  computes  $\beta$  in a CRN. How to compute  $\alpha - \beta$ ?

# Subtraction

Let  $X$  computes  $\alpha$ ,  $Y$  computes  $\beta$  in a CRN. How to compute  $\alpha - \beta$ ?

Ans:

$$Z' = 1 - (X - Y)Z,$$

# Subtraction

Let  $X$  computes  $\alpha$ ,  $Y$  computes  $\beta$  in a CRN. How to compute  $\alpha - \beta$ ?

Ans:

$$Z' = 1 - (X - Y)Z,$$

$$Z'_r = 1 - ZZ_r$$

Addition and multiplication can also be implemented easily.

# Compare to Other Similar Models

Large Population Protocols (LPP).

- It is a simplify CRN model.

---

<sup>2</sup>*Computing with Large Populations Using Interactions*

# Compare to Other Similar Models

Large Population Protocols (LPP).

- It is a simplify CRN model.
- Its reactions look like  $X + Y \rightarrow Z + W$

---

<sup>2</sup>*Computing with Large Populations Using Interactions*



# Compare to Other Similar Models

Large Population Protocols (LPP).

- It is a simplify CRN model.
- Its reactions look like  $X + Y \rightarrow Z + W$
- Bournez et. al. <sup>2</sup> prove that in this model all you can compute are exactly the algebraic numbers.

---

<sup>2</sup>*Computing with Large Populations Using Interactions*

# Compare to Other Similar Models

Large Population Protocols (LPP).

- It is a simplify CRN model.
- Its reactions look like  $X + Y \rightarrow Z + W$
- Bournez et. al. <sup>2</sup> prove that in this model all you can compute are exactly the algebraic numbers.
- However, they assume their protocol must have finitely many equilibrium points. Then one can do a standard *quantifier elimination*.

---

<sup>2</sup>*Computing with Large Populations Using Interactions*

# Compare to Other Similar Models

Large Population Protocols (LPP).

- It is a simplify CRN model.
- Its reactions look like  $X + Y \rightarrow Z + W$
- Bournez et. al. <sup>2</sup> prove that in this model all you can compute are exactly the algebraic numbers.
- However, they assume their protocol must have finitely many equilibrium points. Then one can do a standard *quantifier elimination*.

Generalized Ehrenfest urns:

- A restricted model of LPP

---

<sup>2</sup>Computing with Large Populations Using Interactions

# Compare to Other Similar Models

Large Population Protocols (LPP).

- It is a simplify CRN model.
- Its reactions look like  $X + Y \rightarrow Z + W$
- Bournez et. al. <sup>2</sup> prove that in this model all you can compute are exactly the algebraic numbers.
- However, they assume their protocol must have finitely many equilibrium points. Then one can do a standard *quantifier elimination*.

Generalized Ehrenfest urns:

- A restricted model of LPP
- Reactions look like  $X + Y \rightarrow Z + Z$
- It can not compute all the algebraic numbers, specifically, it can not compute some *rationals*.


---

<sup>2</sup>Computing with Large Populations Using Interactions

But what if we have infinitely many equilibrium points? A continuum of them?

But what if we have infinitely many equilibrium points? A continuum of them?

Well, then we can compute transcendental numbers! The following CRN uses species  $U$ ,  $V$ , such that  $\lim_{t \rightarrow \infty} U(t) - V(t) = e - 1$ . Actually, it is just a fancy way to encode  $y = e^{1-e^{-t}} - 1$  in CRN.



The ODE for the previous CRN is:


$$x' = 1 - x$$

$$u' = 1 + u - xu - uv$$

$$v' = x + v - xv - uv$$

Its equilibrium points are on the curve  $\{x = 1, uv = 1\}$ .

# Simulation by Simbiology





**Thank you**

---