

Xiangying Chen

# The geometry of conditional independence structures and their Coxeter friends

Leipzig, 09.08.2021

**Institut für Algebra und Geometrie**  
Otto-von-Guericke-Universität Magdeburg



DFG-Graduiertenkolleg  
**MATHEMATISCHE  
KOMPLEXITÄTSREDUKTION**

# Overview



# Overview

linear independence

algebraic independence, subforests,

matchings in bipartite graphs, ...

matroids



# Overview

linear independence

algebraic independence, subforests,

matchings in bipartite graphs, ...

matroids

Coxeter matroids



# Overview

linear independence

algebraic independence, subforests,

matchings in bipartite graphs, ...

conditional independence

separation in graphs and topological spaces,

not compared pairs in rank tests, ...

matroids

CI-structures

Coxeter matroids



# Overview

linear independence

algebraic independence, subforests,

matchings in bipartite graphs, ...

conditional independence

separation in graphs and topological spaces,

not compared pairs in rank tests, ...

matroids

CI-structures

Coxeter matroids

Coxeter CI-structures



# Overview

linear independence

algebraic independence, subforests,

matchings in bipartite graphs, ...

matroids

Coxeter matroids

“Combinatorial Erlangen Program”

conditional independence

separation in graphs and topological spaces,

not compared pairs in rank tests, ...

CI-structures

Coxeter CI-structures

“Conditional Erlangen Program”



# Separation and connection

in topological spaces






# Separation and connection

## in topological spaces

Let  $X$  be a topological space.

$\mathcal{G} := \{(ij|K) : K \subseteq X, i \neq j \in X \setminus K \text{ such that } i \text{ and } j \text{ are separated by } K\}.$




# Separation and connection

## in topological spaces

Let  $X$  be a topological space.

$\mathcal{G} := \{(ij|K) : K \subseteq X, i \neq j \in X \setminus K \text{ such that } i \text{ and } j \text{ are separated by } K\}.$



Then  $\mathcal{G}$  satisfies




# Separation and connection

## in topological spaces

Let  $X$  be a topological space.

$\mathcal{G} := \{(ij|K) : K \subseteq X, i \neq j \in X \setminus K \text{ such that } i \text{ and } j \text{ are separated by } K\}.$



Then  $\mathcal{G}$  satisfies

(Ascension)  $(ij|L) \in \mathcal{G} \Rightarrow (ij|kL) \in \mathcal{G},$

(Intersection)  $(ij|kL), (ik|jL) \in \mathcal{G} \Rightarrow (ij|L) \in \mathcal{G},$

(Transtivity)  $(ij|L) \in \mathcal{G} \Rightarrow (ik|L) \in \mathcal{G} \text{ or } (jk|L) \in \mathcal{G}.$



# Separation and connection

in graphs



# Separation and connection

## in graphs

Let  $G$  be a graph with vertex set  $V$  and  $K \subseteq V$ ,  $i \neq j \in V \setminus K$ .

$\mathcal{G} := \{(ij|K) : i, j \text{ are in different conn. comp. in the induced subgraph } G[V \setminus K]\}.$



# Separation and connection

## in graphs

Let  $G$  be a graph with vertex set  $V$  and  $K \subseteq V$ ,  $i \neq j \in V \setminus K$ .

$\mathcal{G} := \{(ij|K) : i, j \text{ are in different conn. comp. in the induced subgraph } G[V \setminus K]\}$ .

Then  $\mathcal{G}$  also satisfies

(Ascension)  $(ij|L) \in \mathcal{G} \Rightarrow (ij|kL) \in \mathcal{G}$ ,

(Intersection)  $(ij|kL), (ik|jL) \in \mathcal{G} \Rightarrow (ij|L) \in \mathcal{G}$ ,

(Transtivity)  $(ij|L) \in \mathcal{G} \Rightarrow (ik|L) \in \mathcal{G} \text{ or } (jk|L) \in \mathcal{G}$ .



# Separation and connection

## in graphs

Let  $G$  be a graph with vertex set  $V$  and  $K \subseteq V$ ,  $i \neq j \in V \setminus K$ .


$\mathcal{G} := \{(ij|K) : i, j \text{ are in different conn. comp. in the induced subgraph } G[V \setminus K]\}$ .

Then  $\mathcal{G}$  also satisfies

(Ascension)  $(ij|L) \in \mathcal{G} \Rightarrow (ij|kL) \in \mathcal{G}$ ,

(Intersection)  $(ij|kL), (ik|jL) \in \mathcal{G} \Rightarrow (ij|L) \in \mathcal{G}$ ,

(Transitivity)  $(ij|L) \in \mathcal{G} \Rightarrow (ik|L) \in \mathcal{G} \text{ or } (jk|L) \in \mathcal{G}$ .



The cycle graph  $C_5$ , which is not topologizable.



# Graphical models

Let  $X_{[n]} = (X_1, \dots, X_n)$  be an  $n$ -dimensional random vector.





# Graphical models

Let  $X_{[n]} = (X_1, \dots, X_n)$  be an  $n$ -dimensional random vector.

For  $K \subseteq [n]$  and  $i \neq j \in [n] \setminus K$  we write

$$X_i \perp\!\!\!\perp X_j | X_K$$

iff the random variables  $X_i$  and  $X_j$  are independent under the condition  $\{X_k : k \in K\}$ .



# Graphical models

Let  $X_{[n]} = (X_1, \dots, X_n)$  be an  $n$ -dimensional random vector.

For  $K \subseteq [n]$  and  $i \neq j \in [n] \setminus K$  we write

$$X_i \perp\!\!\!\perp X_j | X_K$$

iff the random variables  $X_i$  and  $X_j$  are independent under the condition  $\{X_k : k \in K\}$ .

$X_{[n]}$  satisfies the **Markov property** associated to the graph  $G = ([n], E)$  if

$$X_i \perp\!\!\!\perp X_j | X_K \Leftrightarrow \text{vertices } i \text{ and } j \text{ are separated by } K \text{ in } G.$$



# Graphical models

Let  $X_{[n]} = (X_1, \dots, X_n)$  be an  $n$ -dimensional random vector.

For  $K \subseteq [n]$  and  $i \neq j \in [n] \setminus K$  we write

$$X_i \perp\!\!\!\perp X_j | X_K$$

iff the random variables  $X_i$  and  $X_j$  are independent under the condition  $\{X_k : k \in K\}$ .

$X_{[n]}$  satisfies the **Markov property** associated to the graph  $G = ([n], E)$  if

$$X_i \perp\!\!\!\perp X_j | X_K \Leftrightarrow \text{vertices } i \text{ and } j \text{ are separated by } K \text{ in } G.$$

**Remark:** A regular Gaussian random vector  $X_{[n]}$  satisfies the Markov property iff its covariance matrix  $\Sigma$  satisfies

$$(\Sigma^{-1})_{ij} = 0 \quad \text{for all } i \neq j \text{ and } ij \notin E.$$



# Gaussians

Let  $X_{[n]} = (X_1, \dots, X_n)$  be an  $n$ -dimensional Gaussian distributed random vector with positive definite covariance matrix  $\Sigma$ .



# Gaussians

Let  $X_{[n]} = (X_1, \dots, X_n)$  be an  $n$ -dimensional Gaussian distributed random vector with positive definite covariance matrix  $\Sigma$ .

$$\mathcal{G} = [[\Sigma]] := \{(ij|K) : X_i \perp\!\!\!\perp X_j | X_K\} = \{(ij|K) : \det(\Sigma_{ij|K}) = 0\}.$$



# Gaussians

Let  $X_{[n]} = (X_1, \dots, X_n)$  be an  $n$ -dimensional Gaussian distributed random vector with positive definite covariance matrix  $\Sigma$ .

$$\mathcal{G} = [[\Sigma]] := \{(ij|K) : X_i \perp\!\!\!\perp X_j | X_K\} = \{(ij|K) : \det(\Sigma_{ij|K}) = 0\}.$$

Lemma ([Mat02])

$$\det(\Sigma_{kL}) \det(\Sigma_{ij|L}) = \det(\Sigma_L) \det(\Sigma_{ij|kL}) + \det(\Sigma_{ik|L}) \det(\Sigma_{jk|L}).$$



# Gaussians

Let  $X_{[n]} = (X_1, \dots, X_n)$  be an  $n$ -dimensional Gaussian distributed random vector with positive definite covariance matrix  $\Sigma$ .

$$\mathcal{G} = [[\Sigma]] := \{(ij|K) : X_i \perp\!\!\!\perp X_j | X_K\} = \{(ij|K) : \det(\Sigma_{ij|K}) = 0\}.$$

## Lemma ([Mat02])

$$\det(\Sigma_{kL}) \det(\Sigma_{ij|L}) = \det(\Sigma_L) \det(\Sigma_{ij|kL}) + \det(\Sigma_{ik|L}) \det(\Sigma_{jk|L}).$$

Therefore  $\mathcal{G}$  satisfies

(Semigraphoid)  $\{(ij|L), (ik|jL)\} \subseteq \mathcal{G} \Rightarrow \{(ik|L), (ij|kL)\} \subseteq \mathcal{G}$ ,

(Intersection)  $\{(ij|kL), (ik|jL)\} \subseteq \mathcal{G} \Rightarrow \{(ij|L), (ik|L)\} \subseteq \mathcal{G}$ ,

(Composition)  $\{(ij|L), (ik|L)\} \subseteq \mathcal{G} \Rightarrow \{(ij|kL), (ik|jL)\} \subseteq \mathcal{G}$ ,

(Weak transition)  $\{(ij|L), (ij|kL)\} \subseteq \mathcal{G} \Rightarrow (ik|L) \text{ or } (jk|L) \in \mathcal{G}$ .



# Semigraphoids, graphoids, gaussoids

Let  $\mathcal{A}_n$  be the set  $\{(ij|K) : K \subseteq [n], i \neq j \in [n] \setminus K\}$  of conditional independence statements. A conditional independence structure is a subset of  $\mathcal{A}_n$ .





# Semigraphoids, graphoids, gaussoids

Let  $\mathcal{A}_n$  be the set  $\{(ij|K) : K \subseteq [n], i \neq j \in [n] \setminus K\}$  of **conditional independence statements**.

A **conditional independence structure** is a subset of  $\mathcal{A}_n$ .

A CI-structure  $\mathcal{G} \subseteq \mathcal{A}_n$  is

- a **semigraphoid** if it satisfies (Semigraphoid),
- and is a **graphoid** if it satisfies additionally (Intersection),
- and is a **gaussoid** if it satisfies additionally (Composition) and (Weak transition).



# Semigraphoids, graphoids, gaussoids

Let  $\mathcal{A}_n$  be the set  $\{(ij|K) : K \subseteq [n], i \neq j \in [n] \setminus K\}$  of **conditional independence statements**.

A **conditional independence structure** is a subset of  $\mathcal{A}_n$ .

A CI-structure  $\mathcal{G} \subseteq \mathcal{A}_n$  is

- a **semigraphoid** if it satisfies (Semigraphoid),
- and is a **graphoid** if it satisfies additionally (Intersection),
- and is a **gaussoid** if it satisfies additionally (Composition) and (Weak transition).

$[[X]] := \{(ij|K) : i \perp\!\!\!\perp j|K\}$  is a semigraphoid for any random vector  $X$ .

$[[X]]$  is a graphoid for any random vector  $X$  with positive density function.

$[[X]]$  is a gaussoid for any regular Gaussian random vector  $X$ .



# Gaussians, revisited



## Gaussians, revisited

A set function  $\omega: 2^{[n]} \rightarrow \mathbb{R}$  is called **submodular** if for all  $A, B \subseteq [n]$ ,

$$\omega(A) + \omega(B) \geq \omega(A \cap B) + \omega(A \cup B).$$



## Gaussians, revisited

A set function  $\omega: 2^{[n]} \rightarrow \mathbb{R}$  is called **submodular** if for all  $A, B \subseteq [n]$ ,

$$\omega(A) + \omega(B) \geq \omega(A \cap B) + \omega(A \cup B).$$



Figure: Nonsubmodular price



# Gaussians, revisited



## Gaussians, revisited

The principal minors of a positive definite matrix  $\Sigma$  satisfy the Hadamard-Fischer inequalities

$$\det(\Sigma_{I \cap J}) \cdot \det(\Sigma_{I \cup J}) \leq \det(A_I) \cdot \det(A_J) \quad \text{for all } I, J \subseteq [n].$$

That is, the map  $2^{[n]} \rightarrow \mathbb{R}, I \mapsto \log \det(\Sigma_I)$  is submodular.



# Gaussians, revisited

The principal minors of a positive definite matrix  $\Sigma$  satisfy the Hadamard-Fischer inequalities

$$\det(\Sigma_{I \cap J}) \cdot \det(\Sigma_{I \cup J}) \leq \det(\Sigma_I) \cdot \det(\Sigma_J) \quad \text{for all } I, J \subseteq [n].$$

That is, the map  $2^{[n]} \rightarrow \mathbb{R}, I \mapsto \log \det(\Sigma_I)$  is submodular.

## Lemma (Dodgson Condensation)

$$\det(\Sigma_{ij|K})^2 = \det(\Sigma_{iK}) \det(\Sigma_{jK}) - \det(\Sigma_{iK}) \det(\Sigma_{jK}).$$





## Gaussians, revisited

The principal minors of a positive definite matrix  $\Sigma$  satisfy the Hadamard-Fischer inequalities

$$\det(\Sigma_{I \cap J}) \cdot \det(\Sigma_{I \cup J}) \leq \det(\Sigma_I) \cdot \det(\Sigma_J) \quad \text{for all } I, J \subseteq [n].$$

That is, the map  $2^{[n]} \rightarrow \mathbb{R}, I \mapsto \log \det(\Sigma_I)$  is submodular.

### Lemma (Dodgson Condensation)

$$\det(\Sigma_{ij|K})^2 = \det(\Sigma_{iK}) \det(\Sigma_{jK}) - \det(\Sigma_{ijK}) \det(\Sigma_K).$$

Therefore,

$$\begin{aligned} [[\Sigma]] &= \{(ij|K) \in \mathcal{A}_n : X_i \perp\!\!\!\perp X_j | X_K\} = \{(ij|K) \in \mathcal{A}_n : \det(\Sigma_{ij|K}) = 0\} \\ &= \{(ij|K) \in \mathcal{A}_n : \log \det(\Sigma_{iK}) + \log \det(\Sigma_{jK}) = \log \det(\Sigma_{ijK}) + \log \det(\Sigma_K)\}. \end{aligned}$$



# Semimatroid

## Definition

A semigraphoid  $\mathcal{G} \subseteq \mathcal{A}_n$  is a **semimatroid** if there is a submodular function  $\omega : 2^{[n]} \rightarrow \mathbb{R}$  with  $\omega(\emptyset) = 0$  such that

$$\mathcal{G} = [[\omega]] := \{(ij|K) \in \mathcal{A}_n : \omega(iK) + \omega(jK) = \omega(ijK) + \omega(K)\}.$$



# The permutohedral fan

The **permutohedral fan**  $\Sigma_{A_{n-1}}$  is the normal fan of the **permutohedron**

$$\Pi_{n-1} = \text{conv} \left\{ (\delta^{-1}(n), \dots, \delta^{-1}(1))^{\top} : \delta \in \mathfrak{S}_n \right\} = \frac{n}{2} \mathbf{1} + \sum_{1 \leq i < j \leq n} \left[ -\frac{\mathbf{e}_j - \mathbf{e}_i}{2}, \frac{\mathbf{e}_j - \mathbf{e}_i}{2} \right]$$

It can also be defined by the hyperplane arrangement  $\{\{x_i = x_j\} : 1 \leq i < j \leq n\}$  in  $\mathbb{R}^n$ .



# The permutohedral fan

The **permutohedral fan**  $\Sigma_{A_{n-1}}$  is the normal fan of the **permutohedron**

$$\Pi_{n-1} = \text{conv} \left\{ (\delta^{-1}(n), \dots, \delta^{-1}(1))^{\top} : \delta \in \mathfrak{S}_n \right\} = \frac{n}{2} \mathbf{1} + \sum_{1 \leq i < j \leq n} \left[ -\frac{\mathbf{e}_j - \mathbf{e}_i}{2}, \frac{\mathbf{e}_j - \mathbf{e}_i}{2} \right]$$

It can also be defined by the hyperplane arrangement  $\{\{x_i = x_j\} : 1 \leq i < j \leq n\}$  in  $\mathbb{R}^n$ .

- The **chambers** of  $\Sigma_{A_{n-1}}$  are  $\{\mathbf{x} : x_{\delta(1)} \geq \dots \geq x_{\delta(n)}\} =: (\delta(1) | \dots | \delta(n)), \delta \in \mathfrak{S}_n$ .



# The permutohedral fan

The **permutohedral fan**  $\Sigma_{A_{n-1}}$  is the normal fan of the **permutohedron**

$$\Pi_{n-1} = \text{conv} \left\{ (\delta^{-1}(n), \dots, \delta^{-1}(1))^{\top} : \delta \in \mathfrak{S}_n \right\} = \frac{n}{2} \mathbf{1} + \sum_{1 \leq i < j \leq n} \left[ -\frac{\mathbf{e}_j - \mathbf{e}_i}{2}, \frac{\mathbf{e}_j - \mathbf{e}_i}{2} \right]$$

It can also be defined by the hyperplane arrangement  $\{\{x_i = x_j\} : 1 \leq i < j \leq n\}$  in  $\mathbb{R}^n$ .

- The **chambers** of  $\Sigma_{A_{n-1}}$  are  $\{\mathbf{x} : x_{\delta(1)} \geq \dots \geq x_{\delta(n)}\} =: (\delta(1) | \dots | \delta(n))$ ,  $\delta \in \mathfrak{S}_n$ .
- A **wall** is of the form  $\{\mathbf{x} : x_{\delta(1)} \geq \dots \geq x_{\delta(i)} = x_{\delta(i+1)} \geq \dots \geq x_{\delta(n)}\}$ , which is the intersection of two chambers

$$(\delta(1) | \dots | \delta(i) | \delta(i+1) | \dots | \delta(n)) \quad \text{and} \quad (\delta(1) | \dots | \delta(i+1) | \delta(i) | \dots | \delta(n)).$$



# The permutohedral fan

The **permutohedral fan**  $\Sigma_{A_{n-1}}$  is the normal fan of the **permutohedron**

$$\Pi_{n-1} = \text{conv} \left\{ (\delta^{-1}(n), \dots, \delta^{-1}(1))^{\top} : \delta \in \mathfrak{S}_n \right\} = \frac{n}{2} \mathbf{1} + \sum_{1 \leq i < j \leq n} \left[ -\frac{\mathbf{e}_j - \mathbf{e}_i}{2}, \frac{\mathbf{e}_j - \mathbf{e}_i}{2} \right]$$

It can also be defined by the hyperplane arrangement  $\{\{x_i = x_j\} : 1 \leq i < j \leq n\}$  in  $\mathbb{R}^n$ .

- The **chambers** of  $\Sigma_{A_{n-1}}$  are  $\{\mathbf{x} : x_{\delta(1)} \geq \dots \geq x_{\delta(n)}\} =: (\delta(1) | \dots | \delta(n))$ ,  $\delta \in \mathfrak{S}_n$ .
- A **wall** is of the form  $\{\mathbf{x} : x_{\delta(1)} \geq \dots \geq x_{\delta(i)} = x_{\delta(i+1)} \geq \dots \geq x_{\delta(n)}\}$ , which is the intersection of two chambers

$$(\delta(1) | \dots | \delta(i) | \delta(i+1) | \dots | \delta(n)) \quad \text{and} \quad (\delta(1) | \dots | \delta(i+1) | \delta(i) | \dots | \delta(n)).$$

We associate this wall to a CI statement  $\delta(i) \perp\!\!\!\perp \delta(i+1) | \delta(1) \dots \delta(i-1)$  in  $\mathcal{A}_n$ .



# The permutohedral fan

The **permutohedral fan**  $\Sigma_{A_{n-1}}$  is the normal fan of the **permutohedron**

$$\Pi_{n-1} = \text{conv} \left\{ (\delta^{-1}(n), \dots, \delta^{-1}(1))^{\top} : \delta \in \mathfrak{S}_n \right\} = \frac{n}{2} \mathbf{1} + \sum_{1 \leq i < j \leq n} \left[ -\frac{\mathbf{e}_j - \mathbf{e}_i}{2}, \frac{\mathbf{e}_j - \mathbf{e}_i}{2} \right]$$

It can also be defined by the hyperplane arrangement  $\{\{x_i = x_j\} : 1 \leq i < j \leq n\}$  in  $\mathbb{R}^n$ .

- The **chambers** of  $\Sigma_{A_{n-1}}$  are  $\{\mathbf{x} : x_{\delta(1)} \geq \dots \geq x_{\delta(n)}\} =: (\delta(1) | \dots | \delta(n))$ ,  $\delta \in \mathfrak{S}_n$ .
- A **wall** is of the form  $\{\mathbf{x} : x_{\delta(1)} \geq \dots \geq x_{\delta(i)} = x_{\delta(i+1)} \geq \dots \geq x_{\delta(n)}\}$ , which is the intersection of two chambers

$$(\delta(1) | \dots | \delta(i) | \delta(i+1) | \dots | \delta(n)) \quad \text{and} \quad (\delta(1) | \dots | \delta(i+1) | \delta(i) | \dots | \delta(n)).$$

We associate this wall to a CI statement  $\delta(i) \perp\!\!\!\perp \delta(i+1) | \delta(1) \dots \delta(i-1)$  in  $\mathcal{A}_n$ .  
Every CI statement  $(ij | K) \in \mathcal{A}_n$  corresponds to  $|K|!(n - |K| - 2)!$  walls of  $\Sigma_{A_{n-1}}$ .



# The permutohedral fan

- **Ridges:** Let  $s_\ell$  be the reflection across the hyperplane  $\{x_\ell = x_{\ell+1}\}$ ,  $\ell = 1, \dots, n-1$ . Then  $\langle s_1, \dots, s_{n-1} \rangle = \mathfrak{S}_n$ , and a 2-face of  $\Pi_{n-1}$  is a coset  $\delta \cdot \langle s_\ell, s_{\ell'} \rangle$ .

**Case 1:**  $\ell' > \ell + 1$

$$\begin{array}{ccccc}
 (L_1|j|i|L_2|i'|j'|L_3) & i \perp\!\!\!\perp j|L_1 & (L_1|i|j|L_2|i'|j'|L_3) & & \\
 & & & \square & \\
 i' \perp\!\!\!\perp j'|L_1ijL_2 & & i' \perp\!\!\!\perp j'|L_1ijL_2 & & \\
 & & & & \\
 (L_1|j|i|L_2|j'|i'|L_3) & i \perp\!\!\!\perp j|L_1 & (L_1|i|j|L_2|j'|i'|L_3) & & 
 \end{array}$$







# The permutohedral fan

- Ridges:** Let  $s_\ell$  be the reflection across the hyperplane  $\{x_\ell = x_{\ell+1}\}$ ,  $\ell = 1, \dots, n-1$ . Then  $\langle s_1, \dots, s_{n-1} \rangle = \mathfrak{S}_n$ , and a 2-face of  $\Pi_{n-1}$  is a coset  $\delta \cdot \langle s_\ell, s_{\ell'} \rangle$ .

**Case 2:**  $\ell' = \ell + 1$




# Semigraphoids



(Semigraphoid)  $\{(ij|L), (ik|jL)\} \subseteq \mathcal{G} \Rightarrow \{(ik|L), (ij|kL)\} \subseteq \mathcal{G}$



# Semigraphoids




(Semigraphoid)  $\{(ij|L), (ik|jL)\} \subseteq \mathcal{G} \Rightarrow \{(ik|L), (ij|kL)\} \subseteq \mathcal{G}$

A set  $M$  of edges of the permutohedron  $\Pi_{n-1}$  is a semigraphoid iff it satisfies



# Semigraphoids



(Semigraphoid)  $\{(ij|L), (ik|jL)\} \subseteq \mathcal{G} \Rightarrow \{(ik|L), (ij|kL)\} \subseteq \mathcal{G}$


A set  $M$  of edges of the permutohedron  $\Pi_{n-1}$  is a semigraphoid iff it satisfies

(Square) if an edge of a square is in  $M$ , then the opposite edge is also in  $M$ .

(Hexagon) if two adjacent edges of a hexagon are in  $M$ , then the two opposite edges are also in  $M$ .



# Semigraphoids



(Semigraphoid)  $\{(ij|L), (ik|jL)\} \subseteq \mathcal{G} \Rightarrow \{(ik|L), (ij|kL)\} \subseteq \mathcal{G}$

A set  $M$  of edges of the permutohedron  $\Pi_{n-1}$  is a semigraphoid iff it satisfies

(Square) if an edge of a square is in  $M$ , then the opposite edge is also in  $M$ .

(Hexagon) if two adjacent edges of a hexagon are in  $M$ , then the two opposite edges are also in  $M$ .

## Theorem ([Mor+09])

A set of walls of the permutohedral fan  $\Sigma_{A_{n-1}}$  is a semigraphoid iff removing them from  $\Sigma_{A_{n-1}}$  results in a fan.

# Semimatroids



# Semimatroids

A polytope  $P \subseteq \mathbb{R}^n$  is a **generalized permutohedron** if its normal fan coarsens  $\Sigma_{A_{n-1}}$ . Equivalently, there exists a submodular function  $\omega : 2^{[n]} \rightarrow \mathbb{R}$  with  $\omega(\emptyset) = 0$  such that

$$P = \left\{ \mathbf{x} \in \mathbb{R}^n : \sum_{i \in I} x_i \leq \omega(I) \ \forall \emptyset \neq I \subseteq [n], \sum_{i \in [n]} x_i = \omega([n]) \right\}. \quad (1)$$

## Theorem ([Mor+09])

*A semigraphoid is a semimatroid iff the corresponding coarsening of the permutohedral fan is polytopal. In particular, it is the normal fan of the generalized permutohedron (1) defined by the submodular function.*



# Semimatroid

$$\begin{aligned} \{\text{generalized permutohedra}\} &\overset{1:1}{\longleftrightarrow} \{\text{submodular functions}\} \\ \left\{ \begin{array}{l} \text{combinatorial types of} \\ \text{generalized permutohedra} \end{array} \right\} &\overset{1:1}{\longleftrightarrow} \{\text{faces of submodularity cone}\} \overset{1:1}{\longleftrightarrow} \{\text{semimatroids}\} \\ &\hspace{15em} \{\text{facets of submodularity cone}\} \overset{1:1}{\longleftrightarrow} \{\text{CI-statements}\} \end{aligned}$$





# Root systems



# Root systems

A **root system**  $\Phi \subset V$  is a finite set of vectors, called **roots**, which satisfies

(R0)  $\text{span}(\Phi) = V$ ,

(R1)  $\mathbb{R}\alpha \cap \Phi = \{\alpha, -\alpha\}$  for any  $\alpha \in \Phi$ ,

(R2)  $s_\alpha(\Phi) = \Phi$  for any  $\alpha \in \Phi$ .



# Root systems

A **root system**  $\Phi \subset V$  is a finite set of vectors, called **roots**, which satisfies

(R0)  $\text{span}(\Phi) = V$ ,

(R1)  $\mathbb{R}\alpha \cap \Phi = \{\alpha, -\alpha\}$  for any  $\alpha \in \Phi$ ,

(R2)  $s_\alpha(\Phi) = \Phi$  for any  $\alpha \in \Phi$ .

## Theorem


*The irreducible root systems can be completely classified into four infinite families  $A_d, B_d, C_d, D_d$ , the exceptional types  $E_6, E_7, E_8, F_4, G_2, H_3, H_4$  in the dimensions indicated by their subscripts, and  $I_2(m)$  for  $m \geq 3$ .*



# Root systems

## Example

$$\Phi = C_3 = \{\pm \mathbf{e}_1 \pm \mathbf{e}_2, \pm \mathbf{e}_1 \pm \mathbf{e}_3, \pm \mathbf{e}_2 \pm \mathbf{e}_3, \pm 2\mathbf{e}_1, \pm 2\mathbf{e}_2, \pm 2\mathbf{e}_3\}$$



Let  $\Phi$  be a root system.

The **Coxeter complex**  $\Sigma_\Phi$  is the simplicial fan defined by the hyperplane arrangements with the roots as normal vectors.

Fix a chamber  $D$  of  $\Sigma_\Phi$  called the **fundamental domain**.



# Root systems


## Example

$$\Phi = C_3 = \{\pm \mathbf{e}_1 \pm \mathbf{e}_2, \pm \mathbf{e}_1 \pm \mathbf{e}_3, \pm \mathbf{e}_2 \pm \mathbf{e}_3, \pm 2\mathbf{e}_1, \pm 2\mathbf{e}_2, \pm 2\mathbf{e}_3\}$$

$$\Delta_{C_3} = \{\mathbf{e}_1 - \mathbf{e}_2, \mathbf{e}_2 - \mathbf{e}_3, 2\mathbf{e}_3\}$$

$$\lambda_1 = \mathbf{e}_1, \lambda_2 = \mathbf{e}_1 + \mathbf{e}_2,$$

$$\lambda_3 = \mathbf{e}_1 + \mathbf{e}_2 + \mathbf{e}_3$$



The **simple roots**  $\Delta = \{\alpha_1, \dots, \alpha_d\} \subseteq \Phi$  are the roots in  $\Phi$  which are the inner normals of the walls in  $D$ .

The **fundamental weights**  $(\lambda_1, \dots, \lambda_d)$  is the basis of  $V$  dual to the simple **coroots**  $(\alpha_1^\vee, \dots, \alpha_d^\vee)$ , that is,  $\langle \lambda_i, \alpha_j^\vee \rangle = \delta_{ij}$ , where  $\alpha^\vee = \frac{2}{\langle \alpha, \alpha \rangle} \alpha$ .



# Root systems


## Example

$$\Phi = C_3 = \{\pm \mathbf{e}_1 \pm \mathbf{e}_2, \pm \mathbf{e}_1 \pm \mathbf{e}_3, \pm \mathbf{e}_2 \pm \mathbf{e}_3, \pm 2\mathbf{e}_1, \pm 2\mathbf{e}_2, \pm 2\mathbf{e}_3\}$$

$$\Delta_{C_3} = \{\mathbf{e}_1 - \mathbf{e}_2, \mathbf{e}_2 - \mathbf{e}_3, 2\mathbf{e}_3\}$$

$$\lambda_1 = \mathbf{e}_1, \lambda_2 = \mathbf{e}_1 + \mathbf{e}_2,$$

$$\lambda_3 = \mathbf{e}_1 + \mathbf{e}_2 + \mathbf{e}_3$$



$$W_{C_3} \cong \mathbb{Z}_2 \rtimes \mathfrak{S}_3$$

$s_1$  = exchange the 1st and 2nd coord.

$s_2$  = exchange the 2nd and 3rd coord.

$s_3$  = change the sign of the 3rd coord.

$$\langle s_1, s_2 \rangle \cong \mathfrak{S}_3 \cong D_6$$

$$\langle s_1, s_3 \rangle \cong \mathbb{Z}_2 \times \mathbb{Z}_2 \cong D_4$$

$$\langle s_2, s_3 \rangle \cong \mathbb{Z}_2 \rtimes \mathfrak{S}_2 \cong D_8$$

The **Weyl group** of  $\Phi$  is  $W_\Phi := \langle s_\alpha : \alpha \in \Phi \rangle = \langle s_\alpha : \alpha \in \Delta \rangle \subseteq \text{GL}(V)$ .

The **parabolic subgroups** of  $W_\Phi$  are the subgroups

$$(W_\Phi)_I := \langle s_\alpha : \alpha \in I \rangle \subseteq W_\Phi \text{ for } I \subseteq \Delta.$$



## $\Phi$ -permutohedra

The Coxeter complex  $\Sigma_\Phi$  is the normal fan of the  $\Phi$ -permutohedron

$$\Pi_\Phi := \sum_{\alpha \in \Phi_+} [-\alpha/2, \alpha/2] = \text{conv}\{w \cdot \rho : w \in W\},$$

where  $\rho := \frac{1}{2}(\sum_{\alpha \in \Phi_+} \alpha) = \lambda_1 + \dots + \lambda_d$  is the sum of fundamental weights.




Figure: The  $A_3$ ,  $B_3$  ( $C_3$ ) and  $H_3$  permutohedra



# $\Phi$ -permutohedra




Figure: The  $A_3$ ,  $B_3$  ( $C_3$ ) and  $H_3$  permutohedra

polytope	fan	toric variety
truncation	stellar subdivision	blow-up
omnitruncation	Coxeter complex	wonderful compactification





# $\Phi$ -permutohedra




Figure: The  $A_3$ ,  $B_3$  ( $C_3$ ) and  $H_3$  permutohedra

$$\left\{ \begin{array}{c} k\text{-codimensional} \\ \text{cones in the} \\ \text{Coxeter complex } \Sigma_\Phi \end{array} \right\} \xleftrightarrow{1:1} \left\{ \begin{array}{c} k\text{-dimensional} \\ \text{faces in the} \\ \Phi\text{-permutohedron } \Pi_\Phi \end{array} \right\} \xleftrightarrow{1:1} \left\{ \begin{array}{c} \text{cosets of parabolic} \\ \text{subgroups generated} \\ \text{by } k \text{ simple reflections} \end{array} \right\}$$



# $\Phi$ -semigraphoid



# $\Phi$ -semigraphoid

## Definition

Let  $\Phi$  be a root system. A  $\Phi$ -semigraphoid is a fan which is a coarsening of the Coxeter complex  $\Sigma_\Phi$ .



# $\Phi$ -semigraphoid

## Definition

Let  $\Phi$  be a root system. A  $\Phi$ -semigraphoid is a fan which is a coarsening of the Coxeter complex  $\Sigma_\Phi$ .

By the main theorem from [Rea12], we have

## Corollary

*A set of edges  $\mathcal{G}$  of  $\Pi_\Phi$  is a  $\Phi$ -semigraphoid iff it satisfies*

*For every  $2k$ -gonal 2-face  $F$  of  $\Pi_\Phi$ , whenever  $\mathcal{G}$  contains any  $k - 1$  consecutive edges of  $F$ , then  $\mathcal{G}$  also contains the opposite  $k - 1$  consecutive edges of  $F$ .*



# $\Phi$ -semimatroid



# $\Phi$ -semimatroid

## Definition

A  $\Phi$ -semigraphoid, regarded as a fan, is a  $\Phi$ -semimatroid if it is a polytopal fan. That is, a fan coarsening  $\Sigma_\Phi$  which is the normal fan of a polytope  $Q$ . Such a polytope  $Q$  is called a generalized  $\Phi$ -permutohedron.



# $\Phi$ -semimatroid

## Definition

A  $\Phi$ -semigraphoid, regarded as a fan, is a  $\Phi$ -semimatroid if it is a polytopal fan. That is, a fan coarsening  $\Sigma_\Phi$  which is the normal fan of a polytope  $Q$ . Such a polytope  $Q$  is called a **generalized  $\Phi$ -permutohedron**.

A function  $h: \mathcal{R}_\Phi \rightarrow \mathbb{R}$  is  $\Phi$ -submodular if it is convex when regarded as a piecewise linear function  $|\Sigma_\Phi| \rightarrow \mathbb{R}$ . Equivalently, it is the support function of a generalized  $\Phi$ -permutohedron.



## $\Phi$ -semimatroid

A  $\Phi$ -semigraphoid is a  $\Phi$ -semimatroid iff there is a  $\Phi$ -submodular function  $h : \mathcal{R}_\Phi \rightarrow \mathbb{R}$  such that the equality is attained in the **local  $\Phi$ -submodularity inequalities** exactly at its elements.

### Theorem ([Ard+20])

*A function  $h : \mathcal{R}_\Phi \rightarrow \mathbb{R}$  is  $\Phi$ -submodular iff for every  $w \in W_\Phi$  and every simple reflection  $s_i$ , the following local  $\Phi$ -submodularity inequalities hold:*

$$h(w\lambda_i) + h(ws_i\lambda_i) \geq \sum_{j \in N(i)} -2 \frac{\langle \alpha_j, \alpha_i \rangle}{\langle \alpha_j, \alpha_j \rangle} h(w\lambda_j).$$





## $\Phi$ -semimatroid

A  $\Phi$ -semigraphoid is a  $\Phi$ -semimatroid iff there is a  $\Phi$ -submodular function  $h : \mathcal{R}_\Phi \rightarrow \mathbb{R}$  such that the equality is attained in the **local  $\Phi$ -submodularity inequalities** exactly at its elements.

### Theorem ([Ard+20])

*A function  $h : \mathcal{R}_\Phi \rightarrow \mathbb{R}$  is  $\Phi$ -submodular iff for every  $w \in W_\Phi$  and every simple reflection  $s_i$ , the following local  $\Phi$ -submodularity inequalities hold:*

$$h(w\lambda_i) + h(ws_i\lambda_i) \geq \sum_{j \in N(i)} -2 \frac{\langle \alpha_j, \alpha_i \rangle}{\langle \alpha_j, \alpha_j \rangle} h(w\lambda_j).$$



# $\Phi$ -CI-statements



## $\Phi$ -CI-statements

The cone  $SF_\Phi$  of  $\Phi$ -submodular functions: parameter space of generalized  $\Phi$ -permutohedra



## $\Phi$ -CI-statements

The cone  $SF_\Phi$  of  $\Phi$ -submodular functions: parameter space of generalized  $\Phi$ -permutohedra  
Faces of  $SF_\Phi$ :  $\Phi$ -semimatroids



## $\Phi$ -CI-statements

The cone  $SF_\Phi$  of  $\Phi$ -submodular functions: parameter space of generalized  $\Phi$ -permutohedra

Faces of  $SF_\Phi$ :  $\Phi$ -semimatroids

Facets of  $SF_\Phi$ :  $\Phi$ -CI-statements



# $\Phi$ -CI-statements

The cone  $SF_\Phi$  of  $\Phi$ -submodular functions: parameter space of generalized  $\Phi$ -permutohedra

Faces of  $SF_\Phi$ :  $\Phi$ -semimatroids

Facets of  $SF_\Phi$ :  $\Phi$ -CI-statements

## Theorem ([Ard+20])

*Each local  $\Phi$ -submodularity inequality associated with a pair  $(w, s_i)$ , for  $w \in W$  and  $i \in [d]$ , gives a facet of the  $\Phi$ -submodular cone  $SF_\Phi$ . Two pairs  $(w, s_i)$  and  $(w', s_{i'})$  define the same facet iff  $i = i'$  and  $w^{-1}w' \in W_{[d] \setminus N(i)}$ .*



## $\Phi$ -CI-statements

The cone  $SF_\Phi$  of  $\Phi$ -submodular functions: parameter space of generalized  $\Phi$ -permutohedra

Faces of  $SF_\Phi$ :  $\Phi$ -semimatroids

Facets of  $SF_\Phi$ :  $\Phi$ -CI-statements

### Theorem ([Ard+20])

*Each local  $\Phi$ -submodularity inequality associated with a pair  $(w, s_i)$ , for  $w \in W$  and  $i \in [d]$ , gives a facet of the  $\Phi$ -submodular cone  $SF_\Phi$ . Two pairs  $(w, s_i)$  and  $(w', s_{i'})$  define the same facet iff  $i = i'$  and  $w^{-1}w' \in W_{[d] \setminus N(i)}$ .*

The  $\Phi$ -CI statements are exactly the orbits

$$\mathcal{A}_\Phi = \{W_{[d] \setminus N(i)} \cdot wW_{\{i\}} : w \in W, i \in [d]\}.$$



## $\Phi$ -CI-statements

The cone  $SF_\Phi$  of  $\Phi$ -submodular functions: parameter space of generalized  $\Phi$ -permutohedra

Faces of  $SF_\Phi$ :  $\Phi$ -semimatroids

Facets of  $SF_\Phi$ :  $\Phi$ -CI-statements

### Theorem ([Ard+20])

*Each local  $\Phi$ -submodularity inequality associated with a pair  $(w, s_i)$ , for  $w \in W$  and  $i \in [d]$ , gives a facet of the  $\Phi$ -submodular cone  $SF_\Phi$ . Two pairs  $(w, s_i)$  and  $(w', s_{i'})$  define the same facet iff  $i = i'$  and  $w^{-1}w' \in W_{[d] \setminus N(i)}$ .*

The  $\Phi$ -CI statements are exactly the orbits

$$\mathcal{A}_\Phi = \{W_{[d] \setminus N(i)} \cdot wW_{\{i\}} : w \in W, i \in [d]\}.$$

A subset  $\mathcal{G}$  of  $\mathcal{A}_\Phi$  is called a **semigraphoid** resp. **semimatroid** if  $\bigcup \mathcal{G}$  is a semigraphoid resp. semimatroid as a set of edges of  $\Pi_\Phi$ .





## Type $B$ and $C$

Write  $[\pm n] := \{1, \dots, n, \bar{1}, \dots, \bar{n}\}$ , and  $S \sqsubseteq [\pm n]$  if  $\emptyset \neq S \subseteq [\pm n]$  and  $j \in S \Rightarrow \bar{j} \notin S$ .

We define the set of **C-conditional independence (CI) statements** to be

$$\mathcal{C}_n := \{(ij|K) : K \sqsubseteq [\pm n], \{i, j\} \sqsubseteq [\pm n] \setminus (K \cup \bar{K}), i \neq j\} \cup \\ \cup \{(i\bar{i}|K) : K \sqsubseteq [\pm n], |K| = n - 1, i \in [\pm n] \setminus (K \cup \bar{K})\}.$$

A **C-semigraphoid** on  $[n]$  is a subset  $\mathcal{G} \subseteq \mathcal{C}_n$  which satisfies (**Semigraphoid**), and for every  $L \sqsubseteq [\pm n]$ ,  $|L| = n - 2$ ,  $\{i, j\} \sqsubseteq [\pm n] \setminus (L \cup \bar{L})$ ,  $i \neq j$ :

$$(\text{CSG1}) \quad \{(ij|L), (j\bar{j}|iL), (i\bar{j}|L)\} \subseteq \mathcal{G} \Rightarrow \{(\bar{i}\bar{j}|L), (j\bar{j}|\bar{i}L), (\bar{i}\bar{j}|L)\} \subseteq \mathcal{G},$$

$$(\text{CSG2}) \quad \{(i\bar{i}|jL), (\bar{i}\bar{j}|L), (j\bar{j}|\bar{i}L)\} \subseteq \mathcal{G} \Rightarrow \{(i\bar{i}|\bar{j}L), (\bar{i}\bar{j}|L), (j\bar{j}|iL)\} \subseteq \mathcal{G}.$$



## Type $B$ and $C$

For  $f : \mathcal{R}_{C_n} = \{\mathbf{e}_S : S \sqsubseteq [\pm n]\} \rightarrow \mathbb{R}$ , write  $f(S) = f(\mathbf{e}_S)$  for any  $S \sqsubseteq [\pm n]$ .

The function  $f$  is **bisubmodular** if it satisfies the local  $C_n$ -submodularity inequalities

$$\begin{cases} f(Sa) + f(Sb) \geq f(S) + f(Sab) & S \sqsubseteq [\pm n], |S| \leq n-2, ab \sqsubseteq [\pm n] \setminus (S\bar{S}), \\ f(Sa) + f(S\bar{a}) \geq 2f(S) & S \sqsubseteq [\pm n], |S| = n-1, a \in [\pm n] \setminus (S\bar{S}). \end{cases}$$

A  $C$ -semigraphoid  $\mathcal{G}$  on  $[n]$  is a  **$C$ -semimatroid** if there is a bisubmodular function  $f : \{S \sqsubseteq [\pm n]\} \rightarrow \mathbb{R}$  such that the equality is attended in the local  $C_n$ -bisubmodularity inequalities exactly at the triples  $(ij|K) \in \mathcal{G}$ .



## Type $D$

Let

$$\tilde{\mathcal{D}}_n := \{(ij|K) : K \sqsubseteq [\pm n], |K| \leq n-2, \{i, j\} \sqsubseteq [\pm n] \setminus K\bar{K}, i \neq j\} \subseteq \mathcal{C}_n.$$

The set of  **$D$ -CI-statements** is

$$\mathcal{D}_n := \tilde{\mathcal{D}}_n / \sim$$

where  $\sim$  is the equivalence relation in  $\mathcal{D}_n$  defined by

$$\sim := \{((ij|K), (\overline{ij}|K)) \in \tilde{\mathcal{D}}_n \times \tilde{\mathcal{D}}_n : |K| = n-2\}.$$

By abusing of notations, we write an element of  $\tilde{\mathcal{D}}_n$  for its class in  $\mathcal{D}_n$ . In other words, we identify  $(ij|K)$  with  $(\overline{ij}|K)$  for  $|K| = n-2$ .



# Type $D$

A  $D$ -semigraphoid on  $[n]$  is a subset  $\mathcal{G} \subseteq \mathcal{D}_n$  satisfying

(DSG1)  $\{(ij|L), (ik|jL)\} \subseteq \mathcal{G} \Rightarrow \{(ik|L), (ij|kL)\} \subseteq \mathcal{G}$ .

A function  $f : \mathcal{R}_{D_n} \rightarrow \mathbb{R}$  is **disubmodular** if

$$\begin{cases} f(Sa) + f(Sb) \geq f(S) + f(Sab) & S \sqsubseteq [\pm n], |S| \leq n-4, ab \sqsubseteq [\pm n] \setminus S\bar{S}, \\ f(Sa) + f(Sb) \geq f(S) + g(Sabc) + g(Sab\bar{c}) & S \sqsubseteq [\pm n], |S| = n-3, abc \sqsubseteq [\pm n] \setminus S\bar{S}, \\ g(Sab) + g(S\bar{a}\bar{b}) \geq f(S) & S \sqsubseteq [\pm n], |S| = n-2, ab \sqsubseteq [\pm n] \setminus S\bar{S}, \end{cases}$$

where  $f(S) := f(\mathbf{e}_S)$  and  $g(S) := f(\frac{1}{2}\mathbf{e}_S)$ .

A  $D$ -semigraphoid  $\mathcal{G}$  is a  **$D$ -semimatroid** if there is a disubmodular function  $f : \mathcal{R}_{D_n} \rightarrow \mathbb{R}$  such that the equality in the local  $D_n$ -submodularity inequalities is attained exactly at the elements of  $\mathcal{G}$ .



# Rank tests

Assume that we go to the Klein Pizzahaus,



# Rank tests



Assume that we go to the Klein Pizzahaus, the boss gives us 5 Pizzas.



# Rank tests



Assume that we go to the Klein Pizzahaus, the boss gives us 5 Pizzas. He wants us to try them and to score them, so that he can conclude the (partial) ranking of the 5 pizzas from the scores.



# Rank tests



Assume that we go to the Klein Pizzahaus, the boss gives us 5 Pizzas. He wants us to try them and to score them, so that he can conclude the (partial) ranking of the 5 pizzas from the scores.

Then he gives us 5 pizzas and a copy of these 5 pizzas which are exactly so bad as how good the original pizzas are...





# Rank tests



Assume that we go to the Klein Pizzahaus, the boss gives us 5 Pizzas. He wants us to try them and to score them, so that he can conclude the (partial) ranking of the 5 pizzas from the scores.

Then he gives us 5 pizzas and a copy of these 5 pizzas which are exactly so bad as how good the original pizzas are...

Then he gives us 5 pizzas and a copy of these 5 pizzas which are exactly so bad as how good the original pizzas are, but nobody want to distinguish the most similar pair of a pizza and its copy...



# Thank you!





Federico Ardila et al. “Coxeter submodular functions and deformations of Coxeter permutahedra”. In: *Advances in Mathematics* 365 (2020), p. 107039 (cit. on pp. 64–72).



František Matúš. “Conditional independences in Gaussian vectors and rings of polynomials”. In: *International Workshop on Conditionals, Information, and Inference*. Springer. 2002, pp. 152–161 (cit. on pp. 20–23).



Jason Morton et al. “Convex rank tests and semigraphoids”. In: *SIAM Journal on Discrete Mathematics* 23.3 (2009), pp. 1117–1134 (cit. on pp. 42–47).



Nathan Reading. “Coarsening polyhedral complexes”. In: *Proceedings of the American Mathematical Society* 140.10 (2012), pp. 3593–3605 (cit. on pp. 58–60).

