

# MATH CAMP ASSIGNMENT

on Difference Equations

Xiankang Wang

August 20, 2025

## 1 Q1

**Problem.** A multiplier–accelerator model is given by

$$S_t = \alpha Y_t, \quad I_{t+1} = \beta (Y_{t+1} - Y_t), \quad S_t = I_t,$$

with constants  $\beta > \alpha > 0$ . Deduce a difference equation for  $\{Y_t\}$ , given  $Y_0$ , and solve it.

**Solution.** Market clearing holds each period, hence also at  $t + 1$ :

$$S_{t+1} = I_{t+1} \Rightarrow \alpha Y_{t+1} = \beta (Y_{t+1} - Y_t).$$

Rearranging,

$$(\beta - \alpha)Y_{t+1} - \beta Y_t = 0 \iff Y_{t+1} = \lambda Y_t, \quad \lambda := \frac{\beta}{\beta - \alpha} > 1.$$

Therefore  $\{Y_t\}$  satisfies a first–order homogeneous linear difference equation with constant coefficient  $\lambda$ .

With  $Y_0$  given,

$$Y_t = Y_0 \lambda^t = Y_0 \left( \frac{\beta}{\beta - \alpha} \right)^t, \quad t = 0, 1, 2, \dots$$

Since  $\beta > \alpha > 0$ , we have  $\lambda > 1$ , so the path is explosive (monotone growth).

## 2 Q2

(a) Solve  $x_{t+1} = 2x_t + 4$  with  $x_0 = 1$ .

*Method.* Write  $x_{t+1} - 2x_t = 4$ . A constant particular solution satisfies  $x = 2x + 4$ , so  $x^* = -4$ . Let  $y_t := x_t - x^* = x_t + 4$ . Then

$$y_{t+1} = x_{t+1} + 4 = 2x_t + 4 + 4 = 2(x_t + 4) = 2y_t,$$

hence  $y_t = y_0 2^t = (x_0 + 4)2^t = 5 \cdot 2^t$ . Therefore

$$x_t = -4 + 5 \cdot 2^t, \quad t \geq 0.$$

(b) Solve  $x_{t+1} - x_t + 3 = 0$  with  $x_0 = 3$ .

*Method.* The recursion is  $x_{t+1} = x_t - 3$ . Iterating,

$$x_t = x_0 - 3t = 3 - 3t, \quad t \geq 0,$$

so

$$x_t = 3 - 3t, \quad t \geq 0.$$

### 3 Q3

**Problem.** Investigate the stability of

$$x_{t+2} - \frac{1}{6}x_{t+1} - \frac{1}{6}x_t = c_t.$$

**Solution.** First, consider the homogeneous equation

$$x_{t+2} - \frac{1}{6}x_{t+1} - \frac{1}{6}x_t = 0.$$

The characteristic equation is

$$r^2 - \frac{1}{6}r - \frac{1}{6} = 0 \quad \Longleftrightarrow \quad 6r^2 - r - 1 = 0,$$

whose roots are

$$r_1 = \frac{1}{2}, \quad r_2 = -\frac{1}{3}.$$

Therefore, the homogeneous solution is

$$x_t^{(h)} = A \left(\frac{1}{2}\right)^t + B \left(-\frac{1}{3}\right)^t \xrightarrow[t \rightarrow \infty]{} 0.$$

Both roots lie inside the unit circle, so the difference operator is *asymptotically stable*. Thus:

- When  $c_t \equiv 0$ , any solution converges to 0.

- When  $\{c_t\}$  is bounded, since the system is BIBO (Bounded-Input, Bounded-Output) stable, there exists a unique bounded solution, and all solutions are given by the sum of a bounded particular solution and the decaying homogeneous solution.
- If  $c_t \equiv c$  is a constant, let the steady state  $x^*$  satisfy  $x^* - \frac{1}{6}x^* - \frac{1}{6}x^* = c$ , then  $x^* = \frac{3}{2}c$ , and

$$x_t = x^* + A \left(\frac{1}{2}\right)^t + B \left(-\frac{1}{3}\right)^t \xrightarrow{t \rightarrow \infty} x^*.$$

In summary, the equation is stable; with a constant exogenous term, it converges exponentially to the steady state  $\frac{3}{2}c$ .

## 4 Q4

**Problem.** Model

$$C_t = cY_{t-1}, \quad K_t = \sigma Y_{t-1}, \quad Y_t = C_t + K_t - K_{t-1},$$

where  $c, \sigma > 0$ . Derive the second-order difference equation for  $Y_t$ , and give the necessary and sufficient condition for the solution to exhibit *explosive oscillations*.

**Solution.** Substituting gives

$$Y_t = (c + \sigma)Y_{t-1} - \sigma Y_{t-2} \iff Y_t - (c + \sigma)Y_{t-1} + \sigma Y_{t-2} = 0.$$

The characteristic equation is

$$r^2 - (c + \sigma)r + \sigma = 0.$$

If the roots are a complex conjugate pair  $r = \rho e^{\pm i\theta}$ , then

$$\rho^2 = r_1 r_2 = \sigma, \quad 2\rho \cos \theta = r_1 + r_2 = c + \sigma.$$

Therefore

$$(\text{complex roots}) \iff (c + \sigma)^2 < 4\sigma, \quad \rho = \sqrt{\sigma}.$$

The general solution can be written as

$$Y_t = \rho^t [A \cos(t\theta) + B \sin(t\theta)], \quad \rho = \sqrt{\sigma}, \quad \cos \theta = \frac{c + \sigma}{2\sqrt{\sigma}}.$$

So-called *explosive oscillations* mean that the solution oscillates and the amplitude diverges, which is equiv-

alent to having complex roots with modulus  $\rho > 1$ . Thus

$$\text{Explosive oscillations} \iff \begin{cases} (c + \sigma)^2 < 4\sigma, \\ \sigma > 1. \end{cases}$$

When  $\sigma < 1$ , the oscillations are damped. When  $(c + \sigma)^2 \geq 4\sigma$ , the roots are real, and the trajectory is non-oscillatory (it may converge/diverge monotonically or alternate in sign, but not in a sinusoidal manner).

## 5 Q5

**Problem.** Given the system

$$y_t = 0.49 y_{t-1} + 0.68 i_{t-1}, \quad i_t = 0.032 y_{t-1} + 0.43 i_{t-1},$$

where  $y_t$  is output and  $i_t$  is investment.

**(a) Derive the second-order difference equation for  $y_t$  and its characteristic equation**

From the first equation, we get  $i_{t-1} = (y_t - 0.49y_{t-1})/0.68$ . Substituting into the second equation gives

$$i_t = 0.032 y_{t-1} + 0.43 \frac{y_t - 0.49y_{t-1}}{0.68}.$$

Substituting this back into  $y_{t+1} = 0.49y_t + 0.68i_t$  and simplifying, we get

$$y_{t+1} = 0.92 y_t - 0.18894 y_{t-1}.$$

Therefore

$$y_{t+1} - 0.92 y_t + 0.18894 y_{t-1} = 0$$

Its characteristic equation is

$$r^2 - 0.92 r + 0.18894 = 0.$$

### (b) Approximate characteristic roots and the general solution

Solving the above quadratic equation yields

$$r_{1,2} = \frac{0.92 \pm \sqrt{0.92^2 - 4 \times 0.18894}}{2} \approx 0.61053, 0.30947.$$

Both roots are inside the unit circle, so the system is stable. The general solution for  $y_t$  is

$$y_t = A r_1^t + B r_2^t, \quad r_1 \approx 0.61053, \quad r_2 \approx 0.30947,$$

where  $A$  and  $B$  are determined by initial values. From  $y_{t+1} = 0.49y_t + 0.68i_t$ , we can express  $i_t$  as

$$i_t = \frac{y_{t+1} - 0.49y_t}{0.68} = A \frac{r_1 - 0.49}{0.68} r_1^t + B \frac{r_2 - 0.49}{0.68} r_2^t \approx A (0.1773) r_1^t + B (-0.2655) r_2^t.$$

This gives the general solution representation for the two-dimensional system.