# Having a Successful Math Conversation Intuition Behind Analysis

Howard Xiao

June 20, 2023

• Why Language?

- Why Language?
- Words and Emotions


- Why Language?
- Words and Emotions
- Conversations and Information

- Why Language?
- Words and Emotions
- Conversations and Information
- Is It Good?

- Why Language?
- Words and Emotions
- Conversations and Information
- Is It Good?
- Future Work

# {Natural, Programming, Math} Language

• Formalism (Syntax)



# {Natural, Programming, Math} Language

- Formalism (Syntax)
- Logic

# {Natural, Programming, Math} Language

- Formalism (Syntax)
- Logic
- Psychologism

ullet R is a field. Alphabet  $\Leftrightarrow$  Words.

- ullet R is a field. Alphabet  $\Leftrightarrow$  Words.
- ullet R is an **ordered** field. Emotions (Norm)

- $\bullet$   $\mathbb{R}$  is a field. Alphabet  $\Leftrightarrow$  Words.
- ullet R is an **ordered** field. Emotions (Norm)
- $\bullet \ \mathbb{R}$  is a complete ordered field. sup, inf

 $\bullet$  Proper nouns  $\mathbb{Z} = \{\ldots, -3, -2, -1, 0, 1, 2, 3, \ldots\}$ 


- Proper nouns  $\mathbb{Z} = \{\ldots, -3, -2, -1, 0, 1, 2, 3, \ldots\}$
- All nouns  $\mathbb{Q} = \{ \frac{p}{q} : p, q \in \mathbb{Z} \}$

- Proper nouns  $\mathbb{Z} = \{\ldots, -3, -2, -1, 0, 1, 2, 3, \ldots\}$
- ullet All nouns  $\mathbb{Q}=\{rac{p}{q}: p,q\in\mathbb{Z}\}$
- "Good" proper nouns  $\mathbb{N} = \{1, 2, 3, \ldots\}$

• 1 input, 1 response conversation. Emotion values recorded.

- 1 input, 1 response conversation. Emotion values recorded.
- 1-1 Conversations  $f: \mathbb{R} \to \mathbb{R}$

- 1 input, 1 response conversation. Emotion values recorded.
- 1-1 Conversations  $f: \mathbb{R} \to \mathbb{R}$
- Group Group conversations  $g: \mathbb{R}^n \to \mathbb{R}^n$



• Types of Conversations: injective, surjective, bijective

- Types of Conversations: injective, surjective, bijective
- Composition


- Types of Conversations: injective, surjective, bijective
- Composition
- Examples of convesations: sequences, norms, etc.

• Emotional Distances (Metric)

- Emotional Distances (Metric)
- $\forall \epsilon > 0.\exists \delta > 0. \forall x \in A \subset \mathbb{R}^n. 0 < |x a| < \delta \Rightarrow |f(x) I| < \epsilon.$

- Emotional Distances (Metric)
- $\forall \epsilon > 0.\exists \delta > 0. \forall x \in A \subset \mathbb{R}^n. 0 < |x a| < \delta \Rightarrow |f(x) I| < \epsilon.$
- I = f(a). Emotions predictable and logged.
- $a_n \to a \Rightarrow f(a_n) \to f(a)$ .

- Emotional Distances (Metric)
- $\forall \epsilon > 0.\exists \delta > 0. \forall x \in A \subset \mathbb{R}^n. 0 < |x a| < \delta \Rightarrow |f(x) I| < \epsilon.$
- I = f(a). Emotions predictable and logged.
- $a_n \to a \Rightarrow f(a_n) \to f(a)$ .
- Unpredictable emotions.



Limit Properties

- Limit Properties
- Three Important Theorems

- Limit Properties
- Three Important Theorems
- Uniform Continuity

$$\bullet \lim_{x\to a} \frac{f(x)-f(a)}{x-a}.$$

$$\bullet \lim_{x \to a} \frac{f(x) - f(a)}{x - a}.$$

• Predictable emotional change

$$\bullet \lim_{x\to a} \frac{f(x)-f(a)}{x-a}.$$


- Predictable emotional change
- Chain rule:  $(f^*g)' = f'^*g + f^*g'$

$$\bullet \lim_{x\to a}\frac{f(x)-f(a)}{x-a}.$$

- Predictable emotional change
- Chain rule:  $(f^*g)' = f'^*g + f^*g'$
- Emotional change as conversation to input


$$\bullet \lim_{x\to a} \frac{f(x)-f(a)}{x-a}.$$

- Predictable emotional change
- Chain rule:  $(f^*g)' = f'^*g + f^*g'$
- Emotional change as conversation to input
- Unpredictable emotional change



- Increasing/Decreasing, Convex/Concave
- Critical Points
- Mean Value Theorem:  $f'(x) = \frac{f(b) f(a)}{b a}$
- Differentiable ⇒ Continuous

• Partitions, Lower Integral, Upper Integral



Partitions, Lower Integral, Upper Integral

• 
$$F(x) = \int_{a}^{x} f(t)dt$$
 "Information"

Partitions, Lower Integral, Upper Integral

• 
$$F(x) = \int_{a}^{x} f(t)dt$$
 "Information"

• 
$$F(b) - F(a) = \int_a^b f(x) dx$$

- Partitions, Lower Integral, Upper Integral
- $F(x) = \int_{a}^{x} f(t)dt$  "Information"
- $F(b) F(a) = \int_a^b f(x) dx$
- Can define log, exp




- Partitions, Lower Integral, Upper Integral
- $F(x) = \int_{a}^{x} f(t)dt$  "Information"
- $F(b) F(a) = \int_a^b f(x) dx$
- Can define log, exp
- Convergence / Improper Integrals

# Analyzing Multiple Conversations

Uniform Convergence

# Analyzing Multiple Conversations

- Uniform Convergence
- Monotone Convergence Theorem



### Is It Good?

Highly Inmature

### Is It Good?

- Highly Inmature
- Infinity is not intuitive enough.

#### Is It Good?

- Highly Inmature
- Infinity is not intuitive enough.
- Difficult to incorporate Topology
- Hard for computational results and theorems.

## Future of This Project

- Supplement to other analysis materials?
- Write something?
- xiaohoward.github.io/web

