

Lecture 6: Linear Regression

Reading: Sections 3.1-3

STATS 202: Data mining and analysis

Jonathan Taylor, 10/5

Slide credits: Sergio Bacallado

Simple linear regression

Model:

$$y_i = \beta_0 + \beta_1 x_i + \varepsilon_i$$

$$\varepsilon_i \sim \mathcal{N}(0, \sigma) \quad \text{i.i.d.}$$




Figure 3.1

Simple linear regression

Model:

$$y_i = \beta_0 + \beta_1 x_i + \varepsilon_i$$

$$\varepsilon_i \sim \mathcal{N}(0, \sigma) \quad \text{i.i.d.}$$




Figure 3.1

The estimates $\hat{\beta}_0$ and $\hat{\beta}_1$ are chosen to minimize the residual sum of squares (RSS):

$$\text{RSS} = \sum_{i=1}^n (y_i - \hat{y}_i)^2$$

Simple linear regression

Model:

$$y_i = \beta_0 + \beta_1 x_i + \varepsilon_i$$

$$\varepsilon_i \sim \mathcal{N}(0, \sigma) \quad \text{i.i.d.}$$




Figure 3.1

The estimates $\hat{\beta}_0$ and $\hat{\beta}_1$ are chosen to minimize the residual sum of squares (RSS):

$$\begin{aligned} \text{RSS} &= \sum_{i=1}^n (y_i - \hat{y}_i)^2 \\ &= \sum_{i=1}^n (y_i - \beta_0 - \beta_1 x_i)^2. \end{aligned}$$

Estimates $\hat{\beta}_0$ and $\hat{\beta}_1$

A little calculus shows that the minimizers of the RSS are:

$$\hat{\beta}_1 = \frac{\sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})}{\sum_{i=1}^n (x_i - \bar{x})^2},$$
$$\hat{\beta}_0 = \bar{y} - \hat{\beta}_1 \bar{x}.$$

Assesing the accuracy of $\hat{\beta}_0$ and $\hat{\beta}_1$

The Standard Errors for the parameters are:

$$SE(\hat{\beta}_0)^2 = \sigma^2 \left[\frac{1}{n} + \frac{\bar{x}^2}{\sum_{i=1}^n (x_i - \bar{x})^2} \right]$$

$$SE(\hat{\beta}_1)^2 = \frac{\sigma^2}{\sum_{i=1}^n (x_i - \bar{x})^2}.$$




Figure 3.3

Assesing the accuracy of $\hat{\beta}_0$ and $\hat{\beta}_1$

The Standard Errors for the parameters are:

$$SE(\hat{\beta}_0)^2 = \sigma^2 \left[\frac{1}{n} + \frac{\bar{x}^2}{\sum_{i=1}^n (x_i - \bar{x})^2} \right]$$

$$SE(\hat{\beta}_1)^2 = \frac{\sigma^2}{\sum_{i=1}^n (x_i - \bar{x})^2}.$$

The 95% confidence intervals:

$$\hat{\beta}_0 \pm 2 \cdot SE(\hat{\beta}_0)$$

$$\hat{\beta}_1 \pm 2 \cdot SE(\hat{\beta}_1)$$

Calculations depend on the model
above




Figure 3.3

Hypothesis test

H_0 : There is no relationship between X and Y .

H_a : There is some relationship between X and Y .

Hypothesis test

$$H_0: \beta_1 = 0.$$

$$H_a: \beta_1 \neq 0.$$

Hypothesis test

$$H_0: \beta_1 = 0.$$

$$H_a: \beta_1 \neq 0.$$

$$\text{Test statistic: } t = \frac{\hat{\beta}_1 - 0}{\text{SE}(\hat{\beta}_1)}.$$

Under the null hypothesis (special case of the model), this has a t -distribution with $n - 2$ degrees of freedom.

Hypothesis test

$$H_0: \beta_1 = 0.$$

$$H_a: \beta_1 \neq 0.$$

$$\text{Test statistic: } t = \frac{\hat{\beta}_1 - 0}{\text{SE}(\hat{\beta}_1)}.$$

Under the null hypothesis (special case of the model), this has a t -distribution with $n - 2$ degrees of freedom.

	Coefficient	Std. error	t-statistic	p-value
Intercept	7.0325	0.4578	15.36	< 0.0001
TV	0.0475	0.0027	17.67	< 0.0001

TABLE 3.1. For the Advertising data, coefficients of the least squares model for the regression of number of units sold on TV advertising budget. An increase of \$1,000 in the TV advertising budget is associated with an increase in sales by around 50 units (Recall that the sales variable is in thousands of units, and the TV variable is in thousands of dollars).

Interpreting the hypothesis test

- ▶ If we reject the null hypothesis, can we assume there is *an exact* linear relationship?

Interpreting the hypothesis test

- ▶ If we reject the null hypothesis, can we assume there is *an exact* linear relationship?
 - ▶ No. A quadratic relationship may be a better fit, for example. There is evidence of linear association.

Interpreting the hypothesis test

- ▶ If we reject the null hypothesis, can we assume there is *an exact* linear relationship?
 - ▶ No. A quadratic relationship may be a better fit, for example. There is evidence of linear association.
- ▶ If we don't reject the null hypothesis, can we assume there is no relationship between X and Y ?

Interpreting the hypothesis test

- ▶ If we reject the null hypothesis, can we assume there is *an exact* linear relationship?
 - ▶ No. A quadratic relationship may be a better fit, for example. There is evidence of linear association.
- ▶ If we don't reject the null hypothesis, can we assume there is no relationship between X and Y ?
 - ▶ No. This test is only powerful against certain monotone alternatives. There could be more complex non-linear relationships.

Multiple linear regression

$$Y = \beta_0 + \beta_1 X_1 + \cdots + \beta_p X_p + \varepsilon$$

$$\varepsilon \sim \mathcal{N}(0, \sigma) \quad \text{i.i.d.}$$




Figure 3.4

Multiple linear regression




Figure 3.4

$$Y = \beta_0 + \beta_1 X_1 + \cdots + \beta_p X_p + \varepsilon$$

$$\varepsilon \sim \mathcal{N}(0, \sigma) \quad \text{i.i.d.}$$

or, in matrix notation:

$$E\mathbf{y} = \mathbf{X}\boldsymbol{\beta},$$

where $\mathbf{y} = (y_1, \dots, y_n)^T$,
 $\boldsymbol{\beta} = (\beta_0, \dots, \beta_p)^T$ and \mathbf{X} is our
usual data matrix with an extra
column of ones on the left to
account for the intercept.

Multiple linear regression can answer several questions

- ▶ Is at least one of the variables X_i useful for predicting the outcome Y ?

Multiple linear regression can answer several questions

- ▶ Is at least one of the variables X_i useful for predicting the outcome Y ?
- ▶ Which subset of the predictors is most important?

Multiple linear regression can answer several questions

- ▶ Is at least one of the variables X_i useful for predicting the outcome Y ?
- ▶ Which subset of the predictors is most important?
- ▶ How good is a linear model for these data?

Multiple linear regression can answer several questions

- ▶ Is at least one of the variables X_i useful for predicting the outcome Y ?
- ▶ Which subset of the predictors is most important?
- ▶ How good is a linear model for these data?
- ▶ Given a set of predictor values, what is a likely value for Y , and how accurate is this prediction?

The estimates $\hat{\beta}$

Our goal again is to minimize the RSS:

$$\text{RSS} = \sum_{i=1}^n (y_i - \hat{y}_i)^2$$

The estimates $\hat{\beta}$

Our goal again is to minimize the RSS:

$$\begin{aligned}\text{RSS} &= \sum_{i=1}^n (y_i - \hat{y}_i)^2 \\ &= \sum_{i=1}^n (y_i - \beta_0 - \beta_1 x_{i,1} - \cdots - \beta_p x_{i,p})^2.\end{aligned}$$

The estimates $\hat{\beta}$

Our goal again is to minimize the RSS:

$$\begin{aligned}\text{RSS} &= \sum_{i=1}^n (y_i - \hat{y}_i)^2 \\ &= \sum_{i=1}^n (y_i - \beta_0 - \beta_1 x_{i,1} - \cdots - \beta_p x_{i,p})^2.\end{aligned}$$

One can show that this is minimized by the vector $\hat{\beta}$:

$$\hat{\beta} = (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{y}.$$

We also write RSS for the *minimized* sum of squares.

Which variables are important?

Consider the hypothesis:

H_0 : The last q predictors have no relation with Y .

Which variables are important?

Consider the hypothesis:

$$H_0 : \beta_{p-q+1} = \beta_{p-q+2} = \cdots = \beta_p = 0.$$

Which variables are important?

Consider the hypothesis:

$$H_0 : \beta_{p-q+1} = \beta_{p-q+2} = \cdots = \beta_p = 0.$$

Let RSS_0 be the residual sum of squares for the model which excludes these variables.

Which variables are important?

Consider the hypothesis:

$$H_0 : \beta_{p-q+1} = \beta_{p-q+2} = \cdots = \beta_p = 0.$$

Let RSS_0 be the residual sum of squares for the model which excludes these variables. The F -statistic is defined by:

$$F = \frac{(RSS_0 - RSS)/q}{RSS/(n - p - 1)}.$$

Which variables are important?

Consider the hypothesis:

$$H_0 : \beta_{p-q+1} = \beta_{p-q+2} = \cdots = \beta_p = 0.$$

Let RSS_0 be the residual sum of squares for the model which excludes these variables. The F -statistic is defined by:

$$F = \frac{(RSS_0 - RSS)/q}{RSS/(n - p - 1)}.$$

Under the null hypothesis, this has an F -distribution.

Which variables are important?

Consider the hypothesis:

$$H_0 : \beta_{p-q+1} = \beta_{p-q+2} = \cdots = \beta_p = 0.$$

Let RSS_0 be the residual sum of squares for the model which excludes these variables. The F -statistic is defined by:

$$F = \frac{(RSS_0 - RSS)/q}{RSS/(n - p - 1)}.$$

Under the null hypothesis, this has an F -distribution.

Example: If $q = p$, we test whether any of the variables is important.

$$RSS_0 = \sum_{i=1}^n (y_i - \bar{y})^2$$

Which variables are important?

A multiple linear regression in R has the following output:

```
Residuals:
    Min       1Q   Median       3Q      Max
-15.594  -2.730  -0.518   1.777   26.199

Coefficients:
            Estimate Std. Error t value Pr(>|t|)
(Intercept)  3.646e+01  5.103e+00   7.144 3.28e-12 ***
crim         -1.080e-01  3.286e-02  -3.287 0.001087 **
zn           4.642e-02  1.373e-02   3.382 0.000778 ***
indus        2.056e-02  6.150e-02   0.334 0.738288
chas         2.687e+00  8.616e-01   3.118 0.001925 **
nox          -1.777e+01  3.820e+00  -4.651 4.25e-06 ***
rm           3.810e+00  4.179e-01   9.116 < 2e-16 ***
age          6.922e-04  1.321e-02   0.052 0.958229
dis          -1.476e+00  1.995e-01  -7.398 6.01e-13 ***
rad           3.060e-01  6.635e-02   4.613 5.07e-06 ***
tax          -1.233e-02  3.761e-03  -3.280 0.001112 **
ptratio      -9.527e-01  1.308e-01  -7.283 1.31e-12 ***
black         9.312e-03  2.686e-03   3.467 0.000573 ***
lstat        -5.248e-01  5.072e-02  -10.347 < 2e-16 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 4.745 on 492 degrees of freedom
Multiple R-Squared:  0.7406,    Adjusted R-squared:  0.7338
F-statistic: 108.1 on 13 and 492 DF,  p-value: < 2.2e-16
```

Which variables are important?

The t -statistic associated to the i th predictor is the square root of the F -statistic for the null hypothesis which sets only $\beta_i = 0$.

Which variables are important?

The t -statistic associated to the i th predictor is the square root of the F -statistic for the null hypothesis which sets only $\beta_i = 0$.

A low p -value indicates that the predictor is important (it adds something to the model not explained by other variables).

Which variables are important?

The t -statistic associated to the i th predictor is the square root of the F -statistic for the null hypothesis which sets only $\beta_i = 0$.

A low p -value indicates that the predictor is important (it adds something to the model not explained by other variables).

P-hacking warning: If there are many predictors, even under the null hypothesis, some of the t -tests will have low p -values.

How many variables are important?

When we select a subset of the predictors, we have 2^p choices.

How many variables are important?

When we select a subset of the predictors, we have 2^p choices.

A way to simplify the choice is to define a range of models with an increasing number of variables, then select the best.

How many variables are important?

When we select a subset of the predictors, we have 2^p choices.

A way to simplify the choice is to define a range of models with an increasing number of variables, then select the best.

- ▶ **Forward selection:** Starting from a *null model*, include variables one at a time, minimizing the RSS at each step.

How many variables are important?

When we select a subset of the predictors, we have 2^p choices.

A way to simplify the choice is to define a range of models with an increasing number of variables, then select the best.

- ▶ **Forward selection:** Starting from a *null model*, include variables one at a time, minimizing the RSS at each step.
- ▶ **Backward selection:** Starting from the *full model*, eliminate variables one at a time, choosing the one with the largest p-value at each step.

How many variables are important?

When we select a subset of the predictors, we have 2^p choices.

A way to simplify the choice is to define a range of models with an increasing number of variables, then select the best.

- ▶ **Forward selection:** Starting from a *null model*, include variables one at a time, minimizing the RSS at each step.
- ▶ **Backward selection:** Starting from the *full model*, eliminate variables one at a time, choosing the one with the largest p-value at each step.
- ▶ **Mixed selection:** Starting from a *null model*, include variables one at a time, minimizing the RSS at each step. If the p-value for some variable goes beyond a threshold, eliminate that variable.

How many variables are important?

When we select a subset of the predictors, we have 2^p choices.

A way to simplify the choice is to define a range of models with an increasing number of variables, then select the best.

- ▶ **Forward selection:** Starting from a *null model*, include variables one at a time, minimizing the RSS at each step.
- ▶ **Backward selection:** Starting from the *full model*, eliminate variables one at a time, choosing the one with the largest p-value at each step.
- ▶ **Mixed selection:** Starting from a *null model*, include variables one at a time, minimizing the RSS at each step. If the p-value for some variable goes beyond a threshold, eliminate that variable.

Choosing model this way is a form of *tuning*. P-hacking: hypothesis tests, confidence intervals not valid after selection!

How good is the fit?

To assess the fit, we focus on the residuals.

How good is the fit?

To assess the fit, we focus on the residuals.

- ▶ The RSS always decreases as we add more variables.

How good is the fit?

To assess the fit, we focus on the residuals.

- ▶ The RSS always decreases as we add more variables.
- ▶ The residual standard error (RSE) corrects this:

$$\text{RSE} = \sqrt{\frac{1}{n - p - 1} \text{RSS}}.$$


How good is the fit?

To assess the fit, we focus on the residuals.

- ▶ The RSS always decreases as we add more variables.
- ▶ The residual standard error (RSE) corrects this:

$$\text{RSE} = \sqrt{\frac{1}{n - p - 1} \text{RSS}}.$$

- ▶ Visualizing the residuals can reveal phenomena that are not accounted for by the model; eg. synergies or interactions:



How good are the predictions?

The function `predict` in R output predictions from a linear model:

```
> predict(lm.fit, data.frame(lstat=c(5,10,15))),  
       interval="confidence")  
      fit    lwr    upr  
1 29.80 29.01 30.60  
2 25.05 24.47 25.63  
3 20.30 19.73 20.87
```

Confidence intervals reflect the uncertainty on $\hat{\beta}$.

How good are the predictions?

The function `predict` in R output predictions from a linear model:

```
> predict(lm.fit, data.frame(lstat=(c(5,10,15))),  
          interval="confidence")  
      fit   lwr   upr  
1 29.80 29.01 30.60  
2 25.05 24.47 25.63  
3 20.30 19.73 20.87
```


Confidence intervals reflect the uncertainty on $\hat{\beta}$.

```
> predict(lm.fit, data.frame(lstat=(c(5,10,15))),  
          interval="prediction")  
      fit   lwr   upr  
1 29.80 17.566 42.04  
2 25.05 12.828 37.28  
3 20.30  8.078 32.53
```

Prediction intervals reflect uncertainty on $\hat{\beta}$ and the irreducible error ε as well.


Dealing with categorical or qualitative predictors

Example: Credit dataset



Dealing with categorical or qualitative predictors

Example: Credit dataset



In addition, there are 4 qualitative variables:

- ▶ gender: male, female.
- ▶ student: student or not.
- ▶ status: married, single, divorced.
- ▶ ethnicity: African American, Asian, Caucasian.

Dealing with categorical or qualitative predictors

For each qualitative predictor, e.g. `ethnicity`:

- ▶ Choose a baseline category, e.g. African American

Dealing with categorical or qualitative predictors

For each qualitative predictor, e.g. ethnicity:

- ▶ Choose a baseline category, e.g. African American
- ▶ For every other category, define a new predictor:
 - ▶ X_{Asian} is 1 if the person is Asian and 0 otherwise.
 - ▶ $X_{\text{Caucasian}}$ is 1 if the person is Caucasian and 0 otherwise.

Dealing with categorical or qualitative predictors

For each qualitative predictor, e.g. ethnicity:

- ▶ Choose a baseline category, e.g. African American
- ▶ For every other category, define a new predictor:
 - ▶ X_{Asian} is 1 if the person is Asian and 0 otherwise.
 - ▶ $X_{\text{Caucasian}}$ is 1 if the person is Caucasian and 0 otherwise.

The model will be:

$$Y = \beta_0 + \beta_1 X_1 + \cdots + \beta_7 X_7 + \beta_{\text{Asian}} X_{\text{Asian}} + \beta_{\text{Caucasian}} X_{\text{Caucasian}} + \varepsilon.$$

Dealing with categorical or qualitative predictors

For each qualitative predictor, e.g. ethnicity:

- ▶ Choose a baseline category, e.g. African American
- ▶ For every other category, define a new predictor:
 - ▶ X_{Asian} is 1 if the person is Asian and 0 otherwise.
 - ▶ $X_{\text{Caucasian}}$ is 1 if the person is Caucasian and 0 otherwise.

The model will be:

$$Y = \beta_0 + \beta_1 X_1 + \cdots + \beta_7 X_7 + \beta_{\text{Asian}} X_{\text{Asian}} + \beta_{\text{Caucasian}} X_{\text{Caucasian}} + \varepsilon.$$

β_{Asian} is the relative effect on balance for being Asian compared to the baseline category.

Dealing with categorical or qualitative predictors

- ▶ The model fit and predictions are independent of the choice of the baseline category.

Dealing with categorical or qualitative predictors

- ▶ The model fit and predictions are independent of the choice of the baseline category.
- ▶ However, hypothesis tests derived from these variables are affected by the choice.

Dealing with categorical or qualitative predictors

- ▶ The model fit and predictions are independent of the choice of the baseline category.
- ▶ However, hypothesis tests derived from these variables are affected by the choice.
 - ▶ **Solution:** To check whether ethnicity is important, use an F -test for the hypothesis $\beta_{\text{Asian}} = \beta_{\text{Caucasian}} = 0$. This does not depend on the coding.

Dealing with categorical or qualitative predictors

- ▶ The model fit and predictions are independent of the choice of the baseline category.
- ▶ However, hypothesis tests derived from these variables are affected by the choice.
 - ▶ **Solution:** To check whether ethnicity is important, use an F -test for the hypothesis $\beta_{\text{Asian}} = \beta_{\text{Caucasian}} = 0$. This does not depend on the coding.
- ▶ Other ways to encode qualitative predictors produce the same fit \hat{f} , but the coefficients have different interpretations.

Recap

So far, we have:

- ▶ Defined Multiple Linear Regression

Recap

So far, we have:

- ▶ Defined Multiple Linear Regression
- ▶ Discussed how to test the importance of variables.

Recap

So far, we have:

- ▶ Defined Multiple Linear Regression
- ▶ Discussed how to test the importance of variables.
- ▶ Described one approach to choose a subset of variables.

Recap

So far, we have:

- ▶ Defined Multiple Linear Regression
- ▶ Discussed how to test the importance of variables.
- ▶ Described one approach to choose a subset of variables.
- ▶ Explained how to code qualitative variables.

Recap

So far, we have:

- ▶ Defined Multiple Linear Regression
- ▶ Discussed how to test the importance of variables.
- ▶ Described one approach to choose a subset of variables.
- ▶ Explained how to code qualitative variables.
- ▶ Now, how do we evaluate model fit? Is the linear model any good? What can go wrong?

How good is the fit?

To assess the fit, we focus on the residuals.

How good is the fit?

To assess the fit, we focus on the residuals.

- ▶ $R^2 = \text{Corr}(Y, \hat{Y})$, always increases as we add more variables.

How good is the fit?

To assess the fit, we focus on the residuals.

- ▶ $R^2 = \text{Corr}(Y, \hat{Y})$, always increases as we add more variables.
- ▶ The residual standard error (RSE) does not always improve with more predictors:

$$\text{RSE} = \sqrt{\frac{1}{n - p - 1} \text{RSS}}.$$

How good is the fit?

To assess the fit, we focus on the residuals.

- ▶ $R^2 = \text{Corr}(Y, \hat{Y})$, always increases as we add more variables.
- ▶ The residual standard error (RSE) does not always improve with more predictors:

$$\text{RSE} = \sqrt{\frac{1}{n - p - 1} \text{RSS}}.$$

- ▶ **Visualizing the residuals** can reveal phenomena that are not accounted for by the model.

Potential issues in linear regression

1. Interactions between predictors
2. Non-linear relationships
3. Correlation of error terms
4. Non-constant variance of error (heteroskedasticity).
5. Outliers
6. High leverage points
7. Colinearity

Interactions between predictors

Linear regression has an *additive* assumption:

$$\text{sales} = \beta_0 + \beta_1 \times \text{tv} + \beta_2 \times \text{radio} + \varepsilon$$

Interactions between predictors

Linear regression has an *additive* assumption:

$$\text{sales} = \beta_0 + \beta_1 \times \text{tv} + \beta_2 \times \text{radio} + \varepsilon$$

i.e. An increase of \$100 dollars in TV ads causes a fixed increase in sales, regardless of how much you spend on radio ads.


Interactions between predictors

Linear regression has an *additive* assumption:

$$\text{sales} = \beta_0 + \beta_1 \times \text{tv} + \beta_2 \times \text{radio} + \varepsilon$$

i.e. An increase of \$100 dollars in TV ads causes a fixed increase in sales, regardless of how much you spend on radio ads.

If we visualize the residuals, it is clear that this is false:



Interactions between predictors

One way to deal with this is to include multiplicative variables in the model:

$$\text{sales} = \beta_0 + \beta_1 \times \text{tv} + \beta_2 \times \text{radio} + \beta_3 \times (\text{tv} \cdot \text{radio}) + \varepsilon$$

The **interaction variable** is high when both tv and radio are high.

Interactions between predictors

R makes it easy to include interaction variables in the model:

```
> lm.fit=lm(Sales~.+Income:Advertising+Price:Age,data=Carseats)
> summary(lm.fit)


Call:
lm(formula = Sales ~ . + Income:Advertising + Price:Age, data =
    Carseats)

Residuals:
    Min       1Q   Median       3Q      Max
-2.921  -0.750   0.018   0.675   3.341

Coefficients:
              Estimate Std. Error t value Pr(>|t|)
(Intercept)    6.575565    1.008747     6.52 2.2e-10 ***
CompPrice      0.092937    0.004118    22.57 < 2e-16 ***
Income         0.010894    0.002604     4.18 3.6e-05 ***
Advertising    0.070246    0.022609     3.11 0.00203 **
Population     0.000159    0.000368     0.43 0.66533
Price         -0.100806    0.007440   -13.55 < 2e-16 ***
ShelveLocGood  4.848676    0.152838    31.72 < 2e-16 ***
ShelveLocMedium 1.953262    0.125768    15.53 < 2e-16 ***
Age           -0.057947    0.015951    -3.63 0.00032 ***
Education     -0.020852    0.019613    -1.06 0.28836
UrbanYes      0.140160    0.112402     1.25 0.21317
USYes        -0.157557    0.148923    -1.06 0.29073
Income:Advertising 0.000751    0.000278     2.70 0.00729 **
Price:Age      0.000107    0.000133     0.80 0.42381
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

Non-linearities


Example: Auto dataset.



A scatterplot between a predictor and the response may reveal a non-linear relationship.

Non-linearities

Example: Auto dataset.




A scatterplot between a predictor and the response may reveal a non-linear relationship.

Solution: include polynomial terms in the model.

$$\text{MPG} = \beta_0 + \beta_1 \times \text{horsepower} + \varepsilon$$

Non-linearities

Example: Auto dataset.




A scatterplot between a predictor and the response may reveal a non-linear relationship.

Solution: include polynomial terms in the model.

$$\text{MPG} = \beta_0 + \beta_1 \times \text{horsepower} + \beta_2 \times \text{horsepower}^2 + \varepsilon$$

Non-linearities

Example: Auto dataset.




A scatterplot between a predictor and the response may reveal a non-linear relationship.

Solution: include polynomial terms in the model.

$$\begin{aligned} \text{MPG} = & \beta_0 + \beta_1 \times \text{horsepower} \\ & + \beta_2 \times \text{horsepower}^2 \\ & + \beta_3 \times \text{horsepower}^3 + \varepsilon \end{aligned}$$

Non-linearities

Example: Auto dataset.



A scatterplot between a predictor and the response may reveal a non-linear relationship.

Solution: include polynomial terms in the model.

$$\begin{aligned}\text{MPG} = & \beta_0 + \beta_1 \times \text{horsepower} \\ & + \beta_2 \times \text{horsepower}^2 \\ & + \beta_3 \times \text{horsepower}^3 \\ & + \dots + \varepsilon\end{aligned}$$



Non-linearities

In 2 or 3 dimensions, this is easy to visualize. What do we do when we have too many predictors?

Non-linearities

In 2 or 3 dimensions, this is easy to visualize. What do we do when we have too many predictors?

Plot the residuals against the *response* and look for a pattern:



Correlation of error terms

We assumed that the errors for each sample are independent:

$$y_i = f(x_i) + \varepsilon_i \quad ; \quad \varepsilon_i \sim \mathcal{N}(0, \sigma) \text{ i.i.d.}$$

What if this breaks down?

Correlation of error terms

We assumed that the errors for each sample are independent:

$$y_i = f(x_i) + \varepsilon_i \quad ; \quad \varepsilon_i \sim \mathcal{N}(0, \sigma) \text{ i.i.d.}$$

What if this breaks down?

The main effect is that this invalidates any assertions about Standard Errors, confidence intervals, and hypothesis tests:

Example: Suppose that by accident, we double the data (we use each sample twice).

Correlation of error terms

We assumed that the errors for each sample are independent:

$$y_i = f(x_i) + \varepsilon_i \quad ; \quad \varepsilon_i \sim \mathcal{N}(0, \sigma) \text{ i.i.d.}$$

What if this breaks down?

The main effect is that this invalidates any assertions about Standard Errors, confidence intervals, and hypothesis tests:

Example: Suppose that by accident, we double the data (we use each sample twice). Then, the standard errors would be artificially smaller by a factor of $\sqrt{2}$.

Correlation of error terms

When could this happen in real life:

- ▶ **Time series:** Each sample corresponds to a different point in time. The errors for samples that are close in time are correlated.

Correlation of error terms

When could this happen in real life:

- ▶ **Time series:** Each sample corresponds to a different point in time. The errors for samples that are close in time are correlated.
- ▶ **Spatial data:** Each sample corresponds to a different location in space.

Correlation of error terms

When could this happen in real life:

- ▶ **Time series:** Each sample corresponds to a different point in time. The errors for samples that are close in time are correlated.
- ▶ **Spatial data:** Each sample corresponds to a different location in space.
- ▶ Study on predicting height from weight at birth.


Correlation of error terms

When could this happen in real life:

- ▶ **Time series:** Each sample corresponds to a different point in time. The errors for samples that are close in time are correlated.
- ▶ **Spatial data:** Each sample corresponds to a different location in space.
- ▶ Study on predicting height from weight at birth. Suppose some of the subjects in the study are in the same family, their shared environment could make them deviate from $f(x)$ in similar ways.

Correlation of error terms

Simulations of time series with increasing correlations between ε_i .




Non-constant variance of error (heteroskedasticity)

The variance of the error depends on the input.

Non-constant variance of error (heteroskedasticity)

The variance of the error depends on the input.


To diagnose this, we can plot residuals vs. fitted values:



Non-constant variance of error (heteroskedasticity)

The variance of the error depends on the input.

To diagnose this, we can plot residuals vs. fitted values:



Solution: If the trend in variance is relatively simple, we can transform the response using a logarithm, for example.