

# Lecture 18: GAMs

Reading: Sections 7.7

STATS 202: Data mining and analysis

Jonathan Taylor

November 7, 2017

Slide credits: Sergio Bacallado

## Generalized Additive Models (GAMs)

Extension of non-linear models to multiple predictors:

$$\text{wage} = \beta_0 + \beta_1 \times \text{year} + \beta_2 \times \text{age} + \beta_3 \times \text{education} + \epsilon$$

$$\longrightarrow \text{wage} = \beta_0 + f_1(\text{year}) + f_2(\text{age}) + f_3(\text{education}) + \epsilon$$

## Generalized Additive Models (GAMs)

Extension of non-linear models to multiple predictors:

$$\text{wage} = \beta_0 + \beta_1 \times \text{year} + \beta_2 \times \text{age} + \beta_3 \times \text{education} + \epsilon$$

$$\longrightarrow \text{wage} = \beta_0 + f_1(\text{year}) + f_2(\text{age}) + f_3(\text{education}) + \epsilon$$

The functions  $f_1, \dots, f_p$  can be polynomials, natural splines, smoothing splines, local regressions...

## Fitting a GAM

- ▶ If the functions  $f_1$  have a basis representation, we can simply use least squares:
  - ▶ Natural cubic splines
  - ▶ Polynomials
  - ▶ Step functions

$$\text{wage} = \beta_0 + f_1(\text{year}) + f_2(\text{age}) + f_3(\text{education}) + \epsilon$$

## Fitting a GAM

- ▶ Otherwise, we can use **backfitting**:

## Fitting a GAM

► Otherwise, we can use **backfitting**:

1. Keep  $f_2, \dots, f_p$  fixed, and fit  $f_1$  using the partial residuals:

$$y_i - \beta_0 - f_2(x_{i2}) - \dots - f_p(x_{ip}),$$

as the response.

## Fitting a GAM

► Otherwise, we can use **backfitting**:

1. Keep  $f_2, \dots, f_p$  fixed, and fit  $f_1$  using the partial residuals:

$$y_i - \beta_0 - f_2(x_{i2}) - \dots - f_p(x_{ip}),$$

as the response.

2. Keep  $f_1, f_3, \dots, f_p$  fixed, and fit  $f_2$  using the partial residuals:

$$y_i - \beta_0 - f_1(x_{i1}) - f_3(x_{i3}) - \dots - f_p(x_{ip}),$$

as the response.

## Fitting a GAM

- Otherwise, we can use **backfitting**:

1. Keep  $f_2, \dots, f_p$  fixed, and fit  $f_1$  using the partial residuals:

$$y_i - \beta_0 - f_2(x_{i2}) - \dots - f_p(x_{ip}),$$

as the response.

2. Keep  $f_1, f_3, \dots, f_p$  fixed, and fit  $f_2$  using the partial residuals:

$$y_i - \beta_0 - f_1(x_{i1}) - f_3(x_{i3}) - \dots - f_p(x_{ip}),$$

as the response.

3. ...



# Fitting a GAM

- Otherwise, we can use **backfitting**:

1. Keep  $f_2, \dots, f_p$  fixed, and fit  $f_1$  using the partial residuals:

$$y_i - \beta_0 - f_2(x_{i2}) - \dots - f_p(x_{ip}),$$

as the response.

2. Keep  $f_1, f_3, \dots, f_p$  fixed, and fit  $f_2$  using the partial residuals:

$$y_i - \beta_0 - f_1(x_{i1}) - f_3(x_{i3}) - \dots - f_p(x_{ip}),$$

as the response.

3. ...

4. Iterate

## Fitting a GAM

- ▶ Otherwise, we can use **backfitting**:

1. Keep  $f_2, \dots, f_p$  fixed, and fit  $f_1$  using the partial residuals:

$$y_i - \beta_0 - f_2(x_{i2}) - \dots - f_p(x_{ip}),$$

as the response.

2. Keep  $f_1, f_3, \dots, f_p$  fixed, and fit  $f_2$  using the partial residuals:

$$y_i - \beta_0 - f_1(x_{i1}) - f_3(x_{i3}) - \dots - f_p(x_{ip}),$$

as the response.

3. ...

4. Iterate

- ▶ This works for smoothing splines and local regression. **For smoothing splines this is a descent method, descending on convex loss ...**

## Backfitting: coordinate descent

- ▶ Also works for linear regression...

## Backfitting: coordinate descent

- ▶ Also works for linear regression...
  1. Initialize  $\hat{\beta}^{(0)} = 0$  and, (say).

## Backfitting: coordinate descent

► Also works for linear regression...

1. Initialize  $\hat{\beta}^{(0)} = 0$  and, (say).
2. Given  $\hat{\beta}^{(T-1)}$ , choose a coordinate  $0 \leq k(T) \leq p$  and find

$$\begin{aligned}\hat{\alpha}(T) &= \operatorname{argmin}_{\alpha} \sum_{i=1}^n (Y_i - \hat{\beta}_0^{(T-1)} - \sum_{j:j \neq k(T)} X_{ij} \hat{\beta}_j^{(T-1)} - \alpha X_{ik(T)})^2 \\ &= \frac{\sum_{i=1}^n X_{ik(T)} (Y_i - \hat{\beta}_0^{(T-1)} - \sum_{j:j \neq k(T)} X_{ij} \hat{\beta}_j^{(T-1)})}{\sum_{i=1}^n X_{ik(T)}^2}\end{aligned}$$

## Backfitting: coordinate descent

► Also works for linear regression...

1. Initialize  $\hat{\beta}^{(0)} = 0$  and, (say).
2. Given  $\hat{\beta}^{(T-1)}$ , choose a coordinate  $0 \leq k(T) \leq p$  and find

$$\begin{aligned}\hat{\alpha}(T) &= \operatorname{argmin}_{\alpha} \sum_{i=1}^n (Y_i - \hat{\beta}_0^{(T-1)} - \sum_{j:j \neq k(T)} X_{ij} \hat{\beta}_j^{(T-1)} - \alpha X_{ik(T)})^2 \\ &= \frac{\sum_{i=1}^n X_{ik(T)} (Y_i - \hat{\beta}_0^{(T-1)} - \sum_{j:j \neq k(T)} X_{ij} \hat{\beta}_j^{(T-1)})}{\sum_{i=1}^n X_{ik(T)}^2}\end{aligned}$$

3. Set  $\hat{\beta}^{(T)} = \hat{\beta}^{(T-1)}$  except  $k(T)$  entry which we set to  $\hat{\alpha}(T)$ .

## Backfitting: coordinate descent

- ▶ Also works for linear regression...

1. Initialize  $\hat{\beta}^{(0)} = 0$  and, (say).
2. Given  $\hat{\beta}^{(T-1)}$ , choose a coordinate  $0 \leq k(T) \leq p$  and find

$$\begin{aligned}\hat{\alpha}(T) &= \operatorname{argmin}_{\alpha} \sum_{i=1}^n (Y_i - \hat{\beta}_0^{(T-1)} - \sum_{j:j \neq k(T)} X_{ij} \hat{\beta}_j^{(T-1)} - \alpha X_{ik(T)})^2 \\ &= \frac{\sum_{i=1}^n X_{ik(T)} (Y_i - \hat{\beta}_0^{(T-1)} - \sum_{j:j \neq k(T)} X_{ij} \hat{\beta}_j^{(T-1)})}{\sum_{i=1}^n X_{ik(T)}^2}\end{aligned}$$

3. Set  $\hat{\beta}^{(T)} = \hat{\beta}^{(T-1)}$  except  $k(T)$  entry which we set to  $\hat{\alpha}(T)$ .
4. Iterate

## Backfitting: coordinate descent and LASSO

- ▶ Also works for LASSO...



## Backfitting: coordinate descent and LASSO

- ▶ Also works for LASSO...
  1. Initialize  $\hat{\beta}^{(0)} = 0$  and, (say).

## Backfitting: coordinate descent and LASSO

► Also works for LASSO...

1. Initialize  $\hat{\beta}^{(0)} = 0$  and, (say).
2. Given  $\hat{\beta}^{(T-1)}$ , choose a coordinate  $0 \leq k(T) \leq p$  and find

$$\hat{\alpha}_\lambda(T) = \operatorname{argmin}_\alpha \sum_{i=1}^n (r_{ik(T)}^{(T-1)} - \alpha X_{ik(T)})^2 + \lambda \sum_{j:j \neq k(T)} |\hat{\beta}_j^{(T-1)}| + \lambda |\alpha|$$

with  $r_j^{(T-1)}$  the  $j$ -th partial residual at iteration  $T$

$$r_j^{(T-1)} = Y - \hat{\beta}_0^{(T-1)} - \sum_{l:l \neq j} X_l \hat{\beta}_l^{(T-1)}.$$

Solution is a simple soft-thresholded version of previous  $\hat{\alpha}(T)$  –  
**Very fast! Used in glmnet**

## Backfitting: coordinate descent and LASSO

► Also works for LASSO...

1. Initialize  $\hat{\beta}^{(0)} = 0$  and, (say).
2. Given  $\hat{\beta}^{(T-1)}$ , choose a coordinate  $0 \leq k(T) \leq p$  and find

$$\hat{\alpha}_{\lambda}(T) = \underset{\alpha}{\operatorname{argmin}} \sum_{i=1}^n (r_{ik(T)}^{(T-1)} - \alpha X_{ik(T)})^2 + \lambda \sum_{j:j \neq k(T)} |\hat{\beta}_j^{(T-1)}| + \lambda |\alpha|$$

with  $r_j^{(T-1)}$  the  $j$ -th partial residual at iteration  $T$

$$r_j^{(T-1)} = Y - \hat{\beta}_0^{(T-1)} - \sum_{l:l \neq j} X_l \hat{\beta}_l^{(T-1)}.$$

Solution is a simple soft-thresholded version of previous  $\hat{\alpha}(T)$  –  
**Very fast! Used in glmnet**

# Backfitting: coordinate descent and LASSO

► Also works for LASSO...

1. Initialize  $\hat{\beta}^{(0)} = 0$  and, (say).
2. Given  $\hat{\beta}^{(T-1)}$ , choose a coordinate  $0 \leq k(T) \leq p$  and find

$$\hat{\alpha}_{\lambda}(T) = \underset{\alpha}{\operatorname{argmin}} \sum_{i=1}^n (r_{ik(T)}^{(T-1)} - \alpha X_{ik(T)})^2 + \lambda \sum_{j:j \neq k(T)} |\hat{\beta}_j^{(T-1)}| + \lambda |\alpha|$$

with  $r_j^{(T-1)}$  the  $j$ -th partial residual at iteration  $T$

$$r_j^{(T-1)} = Y - \hat{\beta}_0^{(T-1)} - \sum_{l:l \neq j} X_l \hat{\beta}_l^{(T-1)}.$$

Solution is a simple soft-thresholded version of previous  $\hat{\alpha}(T)$  –  
**Very fast! Used in glmnet**

3. Set  $\hat{\beta}^{(T)} = \hat{\beta}^{(T-1)}$  except  $k(T)$  entry which we set to  $\hat{\alpha}_{\lambda}(T)$ .

# Backfitting: coordinate descent and LASSO

► Also works for LASSO...

1. Initialize  $\hat{\beta}^{(0)} = 0$  and, (say).
2. Given  $\hat{\beta}^{(T-1)}$ , choose a coordinate  $0 \leq k(T) \leq p$  and find

$$\hat{\alpha}_{\lambda}(T) = \underset{\alpha}{\operatorname{argmin}} \sum_{i=1}^n (r_{ik(T)}^{(T-1)} - \alpha X_{ik(T)})^2 + \lambda \sum_{j:j \neq k(T)} |\hat{\beta}_j^{(T-1)}| + \lambda |\alpha|$$

with  $r_j^{(T-1)}$  the  $j$ -th partial residual at iteration  $T$

$$r_j^{(T-1)} = Y - \hat{\beta}_0^{(T-1)} - \sum_{l:l \neq j} X_l \hat{\beta}_l^{(T-1)}.$$

Solution is a simple soft-thresholded version of previous  $\hat{\alpha}(T)$  –  
**Very fast! Used in glmnet**

3. Set  $\hat{\beta}^{(T)} = \hat{\beta}^{(T-1)}$  except  $k(T)$  entry which we set to  $\hat{\alpha}_{\lambda}(T)$ .
4. Iterate...

## Backfitting: GAM

- ▶ Let's look at basis functions

## Backfitting: GAM

- ▶ Let's look at basis functions
  1. Initialize  $\hat{\beta}^{(0)} = 0$  and, (say).

## Backfitting: GAM

► Let's look at basis functions

1. Initialize  $\hat{\beta}^{(0)} = 0$  and, (say).
2. Given  $\hat{\beta}^{(T-1)}$ , choose a coordinate  $0 \leq k(T) \leq p$  and find

$$\begin{aligned}\hat{\alpha}_\lambda(T) = \operatorname{argmin}_{\alpha \in \mathbb{R}^{n_{k(T)}}} & \\ & \sum_{i=1}^n (Y_i - \hat{\beta}_0^{(T-1)} - \sum_{j:j \neq k(T)} \sum_{l=1}^{n_j} f_{lj}(X_{ij}) \hat{\beta}_{lj}^{(T-1)}) \\ & - \sum_{l=1}^{n_{k(T)}} \alpha_l f_{lk(T)}(X_{ik(T)})^2\end{aligned}$$



## Backfitting: GAM

► Let's look at basis functions

1. Initialize  $\hat{\beta}^{(0)} = 0$  and, (say).
2. Given  $\hat{\beta}^{(T-1)}$ , choose a coordinate  $0 \leq k(T) \leq p$  and find

$$\begin{aligned}\hat{\alpha}_\lambda(T) = \operatorname{argmin}_{\alpha \in \mathbb{R}^{n_{k(T)}}} & \\ & \sum_{i=1}^n (Y_i - \hat{\beta}_0^{(T-1)} - \sum_{j:j \neq k(T)} \sum_{l=1}^{n_j} f_{lj}(X_{ij}) \hat{\beta}_{lj}^{(T-1)} \\ & - \sum_{l=1}^{n_{k(T)}} \alpha_l f_{lk(T)}(X_{ik(T)}))^2\end{aligned}$$

## Backfitting: GAM

► Let's look at basis functions

1. Initialize  $\hat{\beta}^{(0)} = 0$  and, (say).
2. Given  $\hat{\beta}^{(T-1)}$ , choose a coordinate  $0 \leq k(T) \leq p$  and find

$$\begin{aligned}\hat{\alpha}_\lambda(T) = \operatorname{argmin}_{\alpha \in \mathbb{R}^{n_{k(T)}}} & \\ & \sum_{i=1}^n (Y_i - \hat{\beta}_0^{(T-1)} - \sum_{j:j \neq k(T)} \sum_{l=1}^{n_j} f_{lj}(X_{ij}) \hat{\beta}_{lj}^{(T-1)} \\ & - \sum_{l=1}^{n_{k(T)}} \alpha_l f_{lk(T)}(X_{ik(T)}))^2\end{aligned}$$

3. Set  $\hat{\beta}^{(T)} = \hat{\beta}^{(T-1)}$  except  $k(T)$  entries which we set to  $\hat{\alpha}_\lambda(T)$ .  
**Blockwise coordinate descent!**

# Backfitting: GAM

► Let's look at basis functions

1. Initialize  $\hat{\beta}^{(0)} = 0$  and, (say).
2. Given  $\hat{\beta}^{(T-1)}$ , choose a coordinate  $0 \leq k(T) \leq p$  and find

$$\begin{aligned}\hat{\alpha}_\lambda(T) = \operatorname{argmin}_{\alpha \in \mathbb{R}^{n_{k(T)}}} & \\ & \sum_{i=1}^n (Y_i - \hat{\beta}_0^{(T-1)} - \sum_{j:j \neq k(T)} \sum_{l=1}^{n_j} f_{lj}(X_{ij}) \hat{\beta}_{lj}^{(T-1)} \\ & - \sum_{l=1}^{n_{k(T)}} \alpha_l f_{lk(T)}(X_{ik(T)}))^2\end{aligned}$$

3. Set  $\hat{\beta}^{(T)} = \hat{\beta}^{(T-1)}$  except  $k(T)$  entries which we set to  $\hat{\alpha}_\lambda(T)$ .  
**Blockwise coordinate descent!**
4. Iterate...

## Properties of GAMs

- ▶ GAMs are a step from linear regression toward a fully nonparametric method.

## Properties of GAMs

- ▶ GAMs are a step from linear regression toward a fully nonparametric method.
- ▶ The only constraint is additivity. This can be partially addressed by adding key interaction variables  $X_i X_j$  (or tensor product of basis functions – e.g. polynomials of two variables).


## Properties of GAMs

- ▶ GAMs are a step from linear regression toward a fully nonparametric method.
- ▶ The only constraint is additivity. This can be partially addressed by adding key interaction variables  $X_i X_j$  (or tensor product of basis functions – e.g. polynomials of two variables).
- ▶ We can report degrees of freedom for many non-linear functions.

## Properties of GAMs

- ▶ GAMs are a step from linear regression toward a fully nonparametric method.
- ▶ The only constraint is additivity. This can be partially addressed by adding key interaction variables  $X_i X_j$  (or tensor product of basis functions – e.g. polynomials of two variables).
- ▶ We can report degrees of freedom for many non-linear functions.
- ▶ As in linear regression, we can examine the significance of each of the variables.

## Example: Regression for Wage




year: natural spline with  $df=4$ .

age: natural spline with  $df=5$ .

education: factor.



## Example: Regression for Wage



year: smoothing spline with  $df=4$ .

age: smoothing spline with  $df=5$ .

education: step function.

## GAMs for classification

We can model the log-odds in a classification problem using a GAM:

$$\log \frac{P(Y = 1 \mid X)}{P(Y = 0 \mid X)} = \beta_0 + f_1(X_1) + \cdots + f_p(X_p).$$

Again fit by backfitting ...

## Backfitting: GAM with logistic loss

- ▶ Also works for logistic...

## Backfitting: GAM with logistic loss

- ▶ Also works for logistic...
  1. Initialize  $\hat{\beta}^{(0)} = 0$  and, (say).

## Backfitting: GAM with logistic loss

► Also works for logistic...

1. Initialize  $\hat{\beta}^{(0)} = 0$  and, (say).
2. Given  $\hat{\beta}^{(T-1)}$ , choose a coordinate  $0 \leq k(T) \leq p$  with  $\ell$  logistic loss, find

$$\begin{aligned}\hat{\alpha}_\lambda(T) = \operatorname{argmin}_{\alpha \in \mathbb{R}^{n_{k(T)}}} & \\ & \sum_{i=1}^n \ell \left( Y_i, \hat{\beta}_0^{(T-1)} + \sum_{j:j \neq k(T)} \sum_{l=1}^{n_j} f_{lj}(X_{ij}) \hat{\beta}_{lj}^{(T-1)} \right. \\ & \left. + \sum_{l=1}^{n_{k(T)}} \alpha_l f_{lk(T)}(X_{ik(T)}) \right)\end{aligned}$$

## Backfitting: GAM with logistic loss

► Also works for logistic...


1. Initialize  $\hat{\beta}^{(0)} = 0$  and, (say).
2. Given  $\hat{\beta}^{(T-1)}$ , choose a coordinate  $0 \leq k(T) \leq p$  with  $\ell$  logistic loss, find

$$\begin{aligned}\hat{\alpha}_\lambda(T) = \operatorname{argmin}_{\alpha \in \mathbb{R}^{n_{k(T)}}} & \\ & \sum_{i=1}^n \ell \left( Y_i, \hat{\beta}_0^{(T-1)} + \sum_{j:j \neq k(T)} \sum_{l=1}^{n_j} f_{lj}(X_{ij}) \hat{\beta}_{lj}^{(T-1)} \right. \\ & \left. + \sum_{l=1}^{n_{k(T)}} \alpha_l f_{lk(T)}(X_{ik(T)}) \right)\end{aligned}$$

3. Works for losses that have a *linear predictor*. For GAMs, the linear predictor is

$$\beta_0 + f_1(X_1) + \cdots + f_p(X_p)$$

## Example: Classification for Wage>250




year: linear.

age: smoothing spline with  $df=5$ .

education: step function.

## Example: Classification for Wage>250



year: linear.

age: smoothing spline with  $df=5$ .

education: step function.

Exclude samples with education < HS.