

1. (a) **Solution:**

The number of users should be a poisson distribution which is

$X \sim Poi(\lambda)$  where  $\lambda = 5.5$

The probability that more than 7 users will sign-up for the social networking site in the next minute is

$$P(E) = 1 - P(E^c) = 1 - P\{X \leq 7\} = 1 - \sum_{i=0}^7 \frac{e^{-5.5} 5.5^i}{i!}$$

**Answer:**

We can also use normal distribution to approximate the probability

$$P\{X > 7.5\} = 1 - \Phi\left(\frac{7.5-5.5}{\sqrt{5.5}}\right) = 1 - 0.8032 = 0.1968$$

(b) **Answer:**

The probability that more than 13 users will sign-up for the social networking site in the next 2 minute is

$$P\{X > 13.5\} = 1 - P\{X \leq 13.5\} = 1 - \Phi\left(\frac{13.5-5.5 \times 2}{\sqrt{5.5 \times 2}}\right) = 1 - 0.7734 = 0.2266$$

(c) **Answer:**

The probability that more than 15 users will sign-up for the social networking site in the next 3 minute is

$$P\{X > 15.5\} = 1 - P\{X \leq 15.5\} = 1 - \Phi\left(\frac{15.5-5.5 \times 3}{\sqrt{5.5 \times 3}}\right) = \Phi(0.2462) = 0.5987$$

2. (a) **Solution:**

$$\int_0^1 \int_0^x c \frac{y}{x} dy dx = \int_0^1 \frac{1}{2} c \frac{x^2}{x} dx = \frac{1}{4} c x^2 \Big|_0^1 = \frac{1}{4} c = 1$$

**Answer:**

$$c = 4$$

(b) **Solution:**

$$f_X(x) = \int_0^x 4 \frac{y}{x} dy = 2c \frac{y^2}{x} \Big|_0^x = 2x$$

$$f_Y(y) = \int_y^1 4 \frac{y}{x} dx = cy \ln(x) \Big|_y^1 = -4y \ln(y)$$

$$f_{XY}(x, y) \neq f_X(x) f_Y(y)$$

**Answer:**

X and Y are not independent

(c) **Answer:**

$$f_X(x) = 2x$$

(d) **Answer:**

$$f_Y(y) = -4y \ln(y)$$

(e) **Answer:**

$$E[X] = \int_0^1 2x^2 dx = \frac{2}{3} x^3 \Big|_0^1 = \frac{2}{3}$$

(f) **Answer:**

$$E[Y] = \int_0^1 -4y^2 \ln(y) dy = -4(\ln(y) \int_0^1 y^2 dy - \int_0^1 (\frac{1}{y} \frac{1}{3} y^3) dy)$$

$$E[Y] = -4(\ln(y) \frac{1}{3} y^3 - \frac{1}{9} y^3) \Big|_0^1 = \frac{4}{9}$$

3. (a) **Answer:**

$$P\{X_1 + X_2 \geq 5000\} = P\left\{\frac{X_1 + X_2 - 2200 \times 2}{\sqrt{52900 \times 2}} \geq \frac{5000 - 2200 \times 2}{\sqrt{52900 \times 2}}\right\} = 1 - \Phi(1.84) = 0.033$$

To be more accurate, we should use  $> 4999.5$  instead of  $\geq 5000$ . The result in this particular case has no significant difference.

(b) **Solution:**

$$P\{X \geq 2000\} = 1 - \Phi\left(\frac{2000 - 2200}{\sqrt{52900}}\right) = \Phi(0.87) = 0.808$$

**Answer:**

The probability that the weekly number of visitors exceeds 2000 in at least 2 of the next 3 weeks is

$$\binom{3}{2}p^2(p-1) + \binom{3}{1}p^3 = 0.904$$

4. (a) **Answer:**

$$A \sim N(\mu_1 + \mu_2, \sigma_1^2 + \sigma_2^2)$$

(b) **Answer:**

$$B \sim N(5\mu_1 + 2, 5\sigma_1^2)$$

(c) **Solution:**

$$C_a = aX$$

$$C_b = -bY$$

$$C_c = c^2Z$$

$$C_a \sim N(a\mu_1, a\sigma_1^2)$$

$$C_b \sim N(-b\mu_1, b\sigma_2^2)$$

$$C_c \sim N(c^2\mu_1, c^2\sigma_3^2)$$

**Answer:**

$$C \sim N(a\mu_1 - b\mu_2 + c^2\mu_3, a\sigma_1^2 + b\sigma_2^2 + c^2\sigma_3^2)$$

5. (a) **Answer:**

$$f_{XY}(x, y) = \begin{cases} \frac{1}{6} \frac{1}{x} & x \in \{1, 2, 3, 4, 5, 6\}, y \leq x \\ 0 & \text{Otherwise} \end{cases}$$

(b) **Answer:**

$$P(X = j | Y = i) = \begin{cases} \frac{1}{6} \frac{1}{j} & i \leq j \\ 0 & i > j \end{cases}$$

(c) **Solution:**

$$f_X(x) = \begin{cases} \frac{1}{6} & x \in \{1, 2, 3, 4, 5, 6\} \\ 0 & x \notin \{1, 2, 3, 4, 5, 6\} \end{cases}$$

$$f_Y(y) = \begin{cases} \frac{1}{6} \sum_{i=y}^6 \frac{1}{i} & y \in \{1, 2, 3, 4, 5, 6\} \\ 0 & y \notin \{1, 2, 3, 4, 5, 6\} \end{cases}$$

$$f_{XY}(x, y) \neq f_X(x)f_Y(y)$$

**Answer:**

X and Y are NOT independent

6. **Solution:**

Since the expectation of the distance are symmetrical when the package is dropped off in the 4 different quadrants. Let  $D'$  = the distance the robot travels to get to the package in first quadrants.

$$f'_D(d) = \frac{1}{4} \left(\frac{1}{5}\right)^2$$

$$E[D'] = \frac{1}{100} \int_0^5 \int_0^5 (x+y) dx dy = \frac{1}{100} \int_0^5 \frac{25}{2} + 5y dy = \frac{125}{100}$$

**Answer:**

$$E[D] = 4E[D'] = 4 \frac{125}{100} = 5$$

7. **Solution:**

The probability that "max update" is executed in each loop  $p_i = P(E_i) = \frac{1}{i+1}$ , because the probability that all of the previous independent numbers is

$$\frac{i!}{(i+1)!} = \frac{1}{i+1}$$

Let

$$I_i = \begin{cases} 1 & E_i \text{ occurs} \\ 0 & \text{otherwise} \end{cases}$$

**Answer:**

The expected number of times that "max update" is executed is

$$E[X] = \sum_{i=0}^{n-1} E[I_i] = \sum_{i=0}^{n-1} p_i = \sum_{i=1}^n \frac{1}{i}$$

8. (a) **Solution:**

$$E[X] = \frac{a}{a+b} = 0.5$$

$$Var[X] = \frac{ab}{(a+b)^2(a+b+1)}$$

**Answer:**

$$a = b = 4$$

(b) **Solution:**

$X|13 \text{ flips resulting in 8 heads and 5 tails} \sim \text{Beta}(4+8, 4+5)$

**Answer:**

$$f(x) = \begin{cases} \frac{1}{B(12,9)} x^{11} (1-x)^8 & \text{if } 0 < x < 1 \\ 0 & \text{otherwise} \end{cases}$$

(c) **Solution:**

$$a' = 4 + 8 = 12$$

$$b' = 4 + 4 = 8$$

**Answer:**

$$E[X|12 \text{ flips resulting in 8 heads and 4 tails}] = \frac{a'}{a'+b'} = 0.6$$

(d) **Answer:**

$$Var(X|12 \text{ flips resulting in 8 heads and 4 tails}) = \frac{a'b'}{(a'+b')^2(a'+b'+1)} = 0.0114$$

9. **Solution:**

By post processing the 2 data file we can find

For randomFlips1.txt

$$P_1\{\text{Flip} = \text{Head}\} = 0.49$$

$$P_1\{\text{Flip} = \text{Tail}\} = 0.51$$

$$P_1\{\text{Flip} = \text{Tail} \mid \text{previous flip} = \text{head}\} = 0.694$$

$$P_1\{\text{Flip} = \text{Tail} \mid \text{previous flip} = \text{Tail}\} = 0.329$$

$$P_1\{\text{Flip} = \text{Head} \mid \text{previous flip} = \text{head}\} = 0.306$$

$$P_1\{\text{Flip} = \text{Head} \mid \text{previous flip} = \text{Tail}\} = 0.671$$

From the previous results, we can find that the flips are not independent to the previous flip result.

For example

Let

$P_1(H') = P_1\{\text{previous flip} = \text{head}\}$   
 $P_1(HH') = P_1(H|H')P(H') = 0.306 \times 0.49 = 0.15$   
 $P_1(H)p_1(H') = 0.51 \times 0.51 = 0.26 \neq P_1(HH')$   
 For randomFlips2.txt  
 $P\{\text{Flip} = \text{Head}\} = 0.503$   
 $P\{\text{Flip} = \text{Tail}\} = 0.497$   
 $P\{\text{Flip} = \text{Tail} \mid \text{previous flip} = \text{head}\} = 0.51$   
 $P\{\text{Flip} = \text{Tail} \mid \text{previous flip} = \text{Tail}\} = 0.5$   
 $P\{\text{Flip} = \text{Head} \mid \text{previous flip} = \text{head}\} = 0.49$   
 $P\{\text{Flip} = \text{Head} \mid \text{previous flip} = \text{Tail}\} = 0.5$   
 $P_2(HH') = P_2(H|H')P(H') = 0.25$   
 $P_2(H)p_2(H') = P_2(HH')$

For the 2nd file, we can find that the flips are independent to the previous flip result.

**Answer:**

Sequence 2 is a better random generator, because of the independence between flip results in row.

10. (a) **Answer:**

$$E[X] = 7.41$$

$$E[Y] = 8.03$$

(b) **Answer:**

$$E[X^2] = 58.84$$

$$E[Y^2] = 68.17$$

(c) **Solution:**

$$\mu_A = 7.41, \sigma_A^2 = E[X^2] - E[X]^2 = 3.98$$

$$\mu_B = 8.03, \sigma_B^2 = E[Y^2] - E[Y]^2 = 3.68$$

**Answer:**

$$X \sim N(7.41, 3.98)$$

$$Y \sim N(8.03, 3.68)$$

(d) **Solution:**

Given writer is A, for each keystroke in email, the probability that the duration equals to the timing information is  $P(X_i = x_i|A) = P(E_i|A) = \frac{1}{\sqrt{2\pi}\sigma_A} e^{-(x_i - \mu_A)^2 / 2\sigma_A^2}$

$$\ln(P(E_i|A)) = \ln\left(\frac{1}{\sqrt{2\pi}\sigma_A}\right) - (x_i - \mu_A)^2 / 2\sigma_A^2$$

So, the probability that the email is written given writer is A is

$$\ln(P(E|A)) = \ln\left(\prod_{i=1}^n P(E_i|A)\right) = \sum_{i=1}^n \ln(P(E_i|A))$$

$$\ln(P(E|A)) = n \ln\left(\frac{1}{\sqrt{2\pi}\sigma_A}\right) - \sum_{i=1}^n (x_i - \mu_A)^2 / 2\sigma_A^2 = -248.57$$

The probability that the email is written given writer is B is

$$\ln(P(E|B)) = n \ln\left(\frac{1}{\sqrt{2\pi}\sigma_B}\right) - \sum_{i=1}^n (x_i - \mu_B)^2 / 2\sigma_B^2 = -243.80$$

The probability that the writer is A given the email has been written is

$$P(A|E) = \frac{P(AE)}{P(E)} = \frac{P(E|A)P(A)}{P(E)}$$

The probability that the writer is B given the email has been written is

$$P(B|E) = \frac{P(E|B)P(B)}{P(E)}$$

$$\frac{P(A|E)}{P(B|E)} = \frac{0.5P(E|A)}{0.5P(E|B)} = \frac{P(E|A)}{P(E|B)}$$

$$\frac{P(A|E)}{P(B|E)} = e^{\ln(P(E|A)) - \ln(P(E|B))}$$

**Answer:**

$$\frac{P(A|E)}{P(B|E)} = e^{-4.77} = 0.0085$$

11. (a) **Answer:**

$$P\{X > 7.5\} = 1 - P\{X \leq 7.5\} = 1 - \sum_{i=0}^7 P\{X = i\}$$

$$P\{X > 7.5\} = 0.06$$

(b) **Solution:**

$$\int_{7.5}^{30} \frac{K}{x} dx = 1$$

$$K \ln(x) \Big|_{7.5}^{30} = 0.06$$

$$\int_{7.5}^{30} \frac{C}{x^3} dx = 1$$

$$-C \frac{1}{2x^2} \Big|_{7.5}^{30} = 0.06$$

**Answer:**

$$K = 0.043$$

$$C = 7.2$$

(c) **Answer:**

$$P_1\{X > 10\} = 0.043 \ln(x) \Big|_{10}^{30} = 0.047$$

$$P_2\{X > 10\} = -7.2 \frac{1}{2x^2} \Big|_{10}^{30} = 0.032$$

(d) **Solution:**

$$E_1[X] = \sum_{i=0}^7 P\{X = i\}i + \int_{7.5}^{30} \frac{K}{x} x dx$$

$$E_1[X] = 3.02 + 0.043x \Big|_{7.5}^{30} = 3.99$$

$$E_2[X] = \sum_{i=0}^7 P\{X = i\}i + \int_{7.5}^{30} \frac{C}{x^3} x dx$$

$$E_2[X] = 3.02 - 7.2 \frac{1}{x} \Big|_{7.5}^{30} = 3.74$$

**Answer:**

$$E_1[S] = 3.99$$

$$E_2[S] = 3.74$$

(e) **Solution:**

$$E_1[X^2] = \sum_{i=0}^7 P\{X = i\}i^2 + \int_{7.5}^{30} \frac{K}{x} x^2 dx$$

$$E_1[X^2] = 12.06 + \frac{0.043}{2} x^2 \Big|_{7.5}^{30} = 30.2$$

$$E_2[X^2] = \sum_{i=0}^7 P\{X = i\}i^2 + \int_{7.5}^{30} \frac{C}{x^3} x^2 dx$$

$$E_2[X^2] = 12.06 + 7.2 \ln(x) \Big|_{7.5}^{30} = 22.04$$

**Answer:**

$$E_1[S^2] = 30.2$$

$$E_2[S^2] = 22.04$$