

Refining Quantum Phase Estimation Precision Conditions on Unitaries for Many-Electron Systems

1st Jérémie Messud

TotalEnergies

2, place Jean Millier

92078 Paris La Défense Cedex, France

jeremie.messud@totalenergies.com

2nd Wassil Sennane

QUANTSOC

9 Rue des Colonnnes

75002 Paris, France

direction@qcsocrd.eu

Abstract—Beyond ground state energy estimation, quantum phase estimation (QPE) applied to many-electron systems has the potential to output an approximation of the ground state, enabling in a second step an evaluation of observables other than the energy. We here focus on the impact of approximate controlled-unitaries implementations on QPE precision. After recalling the role of the QPE free parameters, we derive first-order and unified conditions on the unitaries that are necessary to control the QPE energy estimation precision together with the QPE output state precision, important in case we want to leverage the full potential of QPE. We apply these conditions to a Trotterization case, leading to tighter or more general bounds than in previous works. The main results in this article are formal. First numerical illustrations on the H_2 molecule provide useful insights.

Index Terms—Quantum phase estimation, many-electron systems, unitaries precision, Trotterization, state projection.

I. INTRODUCTION

We consider the computation of bound states of many-electron systems, governed by the stationary Schrödinger equation, with a primary focus on the ground state. Finding the exact solution of this equation with classical computing leads to a computational time that grows exponentially with the system size N_S , the latter being represented by the number of spin-orbitals when the state is expanded on an orbital basis [1]. Tractable classical approaches with polynomial scaling in N_S , such as truncated configuration interaction (CI) [2], density functional theory (DFT) [3], and tensor-network methods [4], [5], provide practical approximations. However, these methods can fail to achieve quantitative accuracy for strongly correlated many-electron systems, limiting their predictive capacity in computational chemistry and materials science [6]–[8].

Quantum computing offers an alternative framework to address this challenge. In particular, quantum phase estimation (QPE) could enable the computation of ground state features and potentially achieve exponentially superior scaling in N_S than exact classical schemes [9]–[11]. After early work established the application of QPE to electronic structure problems [12], [13], various subsequent developments refined its application to molecular Hamiltonians, e.g. [14]–[23]. In addition to the estimation of ground state energy (denoted by ground energy in the following to lighten the notations), QPE has the potential to output a ground state approximation [13]

(called QPE ‘projection’ capability), enabling in a second step an evaluation of observables other than the energy.

The practical performance of QPE critically depends on several free parameters. These include the number of phase qubits N , the number of required shots, the overlap of the initial state $|\psi_{\text{init}}\rangle$ with the exact ground state, and the implementation of controlled-unitaries that involve an exponentiation of the system Hamiltonian H together with a free parameter t . In practice, these latter implementation often involves a Trotterization approximation, introducing an additional free parameter called the number of Trotter steps n which provides control on the trade-off between precision (or error) and circuit depth [24]–[27]. A deep understanding of these parameters and their impact is important for an efficient use of QPE in computational chemistry and material science. Reference [28] presents a review of the QPE free parameters with an emphasis on aspects related to the success probability. Complementarily, precision aspects related to an approximate implementation of unitaries remain to be explored further. Indeed, existing precision conditions are known to be quite loose [29], which can lead to an overestimation of the required QPE resources, and specifically related to energy estimation [30].

After recalling the QPE free parameters that are significant for our considerations, we build on the works of [29], [30] and detail a first-order formalism as a unified basis to develop precision conditions on unitaries. We then propose new conditions that contribute to control energy estimation precision together with output state precision, important to leverage the projection potential of QPE. We finally apply these conditions to a Trotterization case and show that they lead to bounds that are tighter than those obtained in [29] and more general than those obtained in [30]. Our main results are formal. A first numerical test on the H_2 molecule provides useful insights.

II. QPE OVERVIEW, CONDITIONS AND PROBLEM

We consider the stationary Schrödinger equation:

$$H |\psi_j\rangle = E_j |\psi_j\rangle, \quad (1)$$

indexed by a positive integer j . E_0 and $|\psi_0\rangle$ denote the ground energy and ground state, respectively.

QPE uses two registers [9], [11]. The first one is an N -qubit phase register $|l\rangle$ whose measure encodes an estimation of an

eigen-energy of the system. The second one is an N_S -qubit system register $|\psi_{\text{init}}\rangle$, related to a spin-orbital expansion. It can be expressed using the basis of eigenstates of H as:

$$|\psi_{\text{init}}\rangle = \sum_{j \geq 0} c_j |\psi_j\rangle \quad , \quad \sum_{j \geq 0} |c_j|^2 = 1. \quad (2)$$

In practice, $|\psi_{\text{init}}\rangle$ is obtained from a prior computation with polynomial cost, e.g. truncated CI [2], tensor networks [4], [5], or parameterized quantum circuits [31], [32]. We define the following unitary operator that acts on the system register:

$$U = e^{-i2\pi t H}, \quad (3)$$

where the usage of Hartree (Ha) units for energy and atomic units for time t is implicit. QPE leverages a series of unitaries U^{2^q} each controlled by the phase qubit number $q \in \{0, \dots, N-1\}$. A phase register measure that gives a value $l \in \{0, \dots, 2^N-1\}$ projects the system register on the following output state:

$$|\psi_{\text{out}}^{(l)}\rangle = \sum_{j \geq 0} c_j^{(l)} |\psi_j\rangle \quad , \quad c_j^{(l)} = c_j \frac{f(\theta_j - \frac{l}{2^N})}{\sqrt{P(l)}}, \quad (4)$$

where the probability associated to the measure is given by:

$$P(l) = \sum_{j \geq 0} |c_j|^2 \left| f\left(\theta_j - \frac{l}{2^N}\right) \right|^2, \quad (5)$$

and $f(\theta_j - \frac{l}{2^N}) = \frac{1}{2^N} \frac{\sin(\pi 2^N (\theta_j - \frac{l}{2^N}))}{\sin(\pi (\theta_j - \frac{l}{2^N}))} e^{i\pi(2^N-1)(\theta_j - \frac{l}{2^N})}$ represents a ‘blurring function’ related to discretization effects, due to the fact that QPE approximates a continuous phase $\theta_j \in [0, 1]$ by a discrete quantity $\frac{l}{2^N}$ [13], [28], [33].

QPE can allow us to approximate E_0 from the most probable phase measurements l^* in the sense of $P(l)$, equation (5), provided an initial state with a sufficiently large $|c_0|^2$ is given and discretization effects are mitigated [13], [28]. We suppose that we are in such a situation in the following. The ground energy is then estimated by ($\lceil \cdot \rceil$ is the ceiling function):

$$E_0 \approx -\frac{1}{t} \frac{l^*}{2^N} + \frac{1}{t} \lceil E_0 t \rceil, \quad (6)$$

which requires an a priori knowledge of $\lceil E_0 t \rceil$ (this will be discussed later). After a l^* measurement, the output state $|\psi_{\text{out}}^{(l^*)}\rangle$ can provide a projection on the ground state $|\psi_0\rangle$ in the situations discussed in [13], [28], i.e. when $|c_0^{(l^*)}|^2 \gg |c_{j>0}^{(l^*)}|^2$.

QPE’s efficiency relies on various free parameters [28]. We remind conditions on the parameters that are important for the developments here, related to unitaries implementation.

(t condition.) The time t choice is the first parameter to set as it is crucial to allow to recover the ground energy E_0 from $\frac{l^*}{2^N}$, i.e. to constrain *a priori* the $\lceil E_0 t \rceil$ value in (6). Given that the problem Hamiltonian is naturally expressed as a Linear Combination of Unitaries (LCU) [34]–[36],

$$H = \sum_{\beta=1}^M \gamma_{\beta} H_{\beta} \quad , \quad \gamma_{\beta} \in \mathbb{R}, \quad (7)$$

where H_{β} are Hermitian unitaries, it is common to choose $t = \frac{\alpha}{\sum_{\beta} |\gamma_{\beta}|}$ where $\alpha \in [\frac{1}{2}, 1]$, which implies $\lceil E_0 t \rceil = 0$. Two

other cases leveraging prior information and leading to larger t values were derived in [28].

(N condition.) Once t defined and the ambiguity on $\lceil E_0 t \rceil$ removed, we can set accordingly the number of phase qubits. Reaching chemical precision $\varepsilon_{\text{ch}} = 1.6 \times 10^{-3}$ Ha on energy estimation requires:

$$N \geq N_{\min}(t) = \left\lceil \log_2 \left(\frac{1}{t \varepsilon_{\text{ch}}} \right) \right\rceil - 1. \quad (8)$$

Interestingly, the minimum number of phase qubits $N_{\min}(t)$ is directly related to the choice made for t , a smaller t leading to a larger $N_{\min}(t)$. Adding more phase qubits, i.e. taking $N = N_{\min}(t) + a$ where a is a positive integer, can make sense to mitigate discretization effects or increase the confidence level, as detailed in [28].

(U^{2^q} condition.) The practical and approximate implementation of U^{2^q} is denoted by $\mathcal{S}(U^{2^q})$. The following condition constrains achieving a precision driven by ε on the approximate unitaries ($\|\cdot\|_2$ denotes the spectral norm):

$$\|U^{2^q} - \mathcal{S}(U^{2^q})\|_2 \lesssim 2\pi t 2^q \varepsilon = \frac{2^q}{2^{N_{\min}(t)}} \pi \text{ if } \varepsilon = \varepsilon_{\text{ch}}, \quad (9)$$

the equality being obtained using (8). As we will demonstrate, choosing $\varepsilon = \varepsilon_{\text{ch}}$ can be related to first-order to the ability of the approximate unitaries to achieve chemical precision on energy, but smaller ε choices can make sense to also control output state precision. In the case of order- p Trotterization for $\mathcal{S}(U^{2^q})$, an upper bound on the minimum number $n_{\min}^{(p)}(q, t)$ of Trotter steps to reach precision ε in (9) is given by [28]–[30]:

$$n_{\min}^{(p)}(q, t) \approx \left\lceil 2\pi 2^q t (C_p / \varepsilon)^{\frac{1}{p}} \right\rceil, \quad (10)$$

where forms for C_p will be given later. Then, if $\varepsilon = \varepsilon_{\text{ch}}$ is constrained, we have $n_{\min}^{(p)}(q, t) \approx \left\lceil \frac{2^q}{2^{N_{\min}(t)}} \pi \left(\frac{C_p}{\varepsilon^{p+1}} \right)^{\frac{1}{p}} \right\rceil$, and a minimum number of Trotter steps over the whole QPE circuit,

$$n_{\min\text{-tot}}^{(p)} \approx \lceil 2^a \mathcal{C}_p \rceil, \quad \mathcal{C}_p = \pi (C_p / \varepsilon^{p+1})^{\frac{1}{p}}, \quad (11)$$

that tends to be independent of $N_{\min}(t)$ or t , and to depend only on the number a of phase qubits beyond $N_{\min}(t)$. The corresponding QPE complexity equals $\lceil 2^a \mathcal{C}_p \rceil \times \mathcal{N}_p(N_S)$, where $\mathcal{N}_p(N_S)$ represents one Trotter step complexity.

However, challenges related to the precision of approximate implementations of unitaries remain. The work in [29] leads to a specific C_p form which overestimates $n_{\min}^{(p)}$ and thus the required QPE resources [30]. The work in [30] leads to a tighter bound but it is energy specific. To our knowledge, conditions to control the output state precision remain to be studied, which is important in case we want to leverage the projection potential of QPE. We here build on the works of [29], [30] and detail a unified first-order formalism that allows us to propose new conditions that contribute to control QPE energy estimation and output state precisions together.

III. UNITARY OPERATOR PRECISION CONDITION

The ‘approximate’ $\mathcal{S}(U^{2^q})$ must be unitary to be implementable in a quantum framework. We consider it is defined through an ‘effective QPE Hamiltonian’ $H^{\mathcal{S}}(\lambda)$ (Hermitian):

$$\mathcal{S}(U^{2^q}) = e^{-i2\pi t 2^q H^{\mathcal{S}}(\lambda)} \quad , \quad \lambda \in \mathbb{R}^+, \quad (12)$$

where the λ parameter controls the ‘quality’ of the effective Hamiltonian, $H^S(\lambda)$ getting closer to H as λ gets smaller. In other terms, $H^S(\lambda)$ is defined by the Hermitian logarithm of the approximate unitary (a Trotterization case will be detailed below). We denote by $E_i^S(\lambda)$ the eigen-energies of $H^S(\lambda)$ and by $|\psi_i^S(\lambda)\rangle$ its eigen-states, which represent the ‘effective QPE energies and states’ (resp.). We do not explicit the λ dependency of $\mathcal{S}(U^{2^q})$ to keep the notations light.

As H and $H_S(\lambda)$ are Hermitian, corollary A.5 of [29] gives:

$$\|U^{2^q} - \mathcal{S}(U^{2^q})\|_2 \leq 2\pi t 2^q \|H - H^S(\lambda)\|_2, \quad (13)$$

which is not directly interpretable as a precision condition on energy $E_i^S(\lambda)$ or state $|\psi_i^S(\lambda)\rangle$. To obtain such an interpretation, we consider in the following a first-order development of $H^S(\lambda)$ in λ , accurate for sufficiently small λ values (we will come back to the pertinency of this assumption later):

$$H^S(\lambda) = H + \lambda \delta H + \mathcal{O}(\lambda^2), \quad \delta H = \delta H^\dagger. \quad (14)$$

From (13)-(14), we have:

$$\|U^{2^q} - \mathcal{S}(U^{2^q})\|_2 \leq 2\pi t 2^q \|\lambda \delta H\|_2 + \mathcal{O}(\lambda^2), \quad (15)$$

which leads to the following first-order condition:

$$\boxed{\frac{1}{2\pi t 2^q} \|U^{2^q} - \mathcal{S}(U^{2^q})\|_2 \lesssim \|\lambda \delta H\|_2} \quad (16)$$

(unitary condition to 1st-order in λ),

that upper-bounds the unitaries difference. All $\approx$ (instead of $=$) and $\lesssim$ (instead of $\leq$) in this article highlight that a first-order approximation has been considered to obtain the result.

IV. ENERGY PRECISION CONDITION

We aim at controlling the eigen-energy precision related to the effective QPE Hamiltonian, important for an usage of QPE for energy estimation. From now on, we consider perturbation theory for a non-degenerate state i . Note that the other states $j \neq i$ can be degenerate without affecting any of our results (we thus do not make potential degeneracy explicit, which lightens the notations). In the end, we will focus on $i = 0$ meaning that we will suppose a non-degenerate ground state, which is preferable in the context of an usage of QPE for projection [28].

Perturbation theory gives $E_i^S(\lambda) = E_i + \langle \psi_i | \lambda \delta H | \psi_i \rangle + \mathcal{O}(\lambda^2)$ (chapter XI of [37]), which leads to:

$$|E_i^S(\lambda) - E_i| = |\langle \psi_i | \lambda \delta H | \psi_i \rangle| + \mathcal{O}(\lambda^2). \quad (17)$$

To bound the energy difference, we establish the relation:

$$|\langle \psi_i | \delta H | \psi_i \rangle| \leq \sqrt{\langle \psi_i | (\delta H)^2 | \psi_i \rangle} \leq \|\delta H\|_2, \quad (18)$$

where the first inequality is obvious and the second inequality is related to the definition of the spectral norm for a Hermitian operator, i.e. $\|\delta H\|_2 = \max_{\|\psi\|_2=1} \sqrt{\langle \psi | (\delta H)^2 | \psi \rangle}$.

From (17) and (18), we deduce:

$$\boxed{|E_i^S(\lambda) - E_i| \lesssim \sqrt{\langle \psi_i | (\lambda \delta H)^2 | \psi_i \rangle} \leq \|\lambda \delta H\|_2} \quad (19)$$

(energy condition to 1st-order in λ).

This gives options to upper-bound the energy difference.

V. STATE PRECISION CONDITION

We aim at controlling the precision of the eigen-state $|\psi_i^S(\lambda)\rangle$ related to the effective QPE Hamiltonian, necessary in the context of a usage of QPE for projection. A reasoning directly on states is impractical due to their dimensionality. Scalar measures of distinguishability between states like the fidelity or trace distance [11] are commonly used.

Following [38], [39], perturbation theory leads to ¹:

$$\begin{aligned} |\langle \psi_i | \psi_i^S(\lambda) \rangle| &= 1 - \frac{\lambda^2}{2} \sum_{k \neq i} \frac{|\langle \psi_k | \delta H | \psi_i \rangle|^2}{(E_k - E_i)^2} + \mathcal{O}(\lambda^4) \\ \Rightarrow |\langle \psi_i | \psi_i^S(\lambda) \rangle|^2 &= 1 - \lambda^2 \sum_{k \neq i} \frac{|\langle \psi_k | \delta H | \psi_i \rangle|^2}{(E_k - E_i)^2} + \mathcal{O}(\lambda^4). \end{aligned}$$

This allows us to evaluate among others the trace distance, equal in our case to the square root of one minus the fidelity:

$$\begin{aligned} T(|\psi_i\rangle, |\psi_i^S(\lambda)\rangle) &= \sqrt{1 - |\langle \psi_i | \psi_i^S(\lambda) \rangle|^2} \\ &= \lambda \sqrt{\sum_{k \neq i} \frac{|\langle \psi_k | \delta H | \psi_i \rangle|^2}{(E_k - E_i)^2}} + \mathcal{O}(\lambda^2). \end{aligned} \quad (20)$$

Note that, contrariwise to the energy, having λ small is not sufficient to ensure a small first-order term for trace distance. We must additionally have: $\forall k \neq i : |E_k - E_i| \not\ll \lambda |\langle \psi_k | \delta H | \psi_i \rangle|$, making state precision conditions usually more stringent than energy conditions.

To bound the trace distance, we establish the relation:

$$\begin{aligned} \sqrt{\sum_{k \neq i} \frac{|\langle \psi_k | \delta H | \psi_i \rangle|^2}{(E_k - E_i)^2}} &\leq \frac{1}{\Delta E_i} \sqrt{\sum_{k \neq i} |\langle \psi_k | \delta H | \psi_i \rangle|^2} \\ &= \frac{1}{\Delta E_i} \sqrt{\langle \psi_i | (\delta H)^2 | \psi_i \rangle - \langle \psi_i | \delta H | \psi_i \rangle^2} \\ &= \frac{A_i}{\Delta E_i} \sqrt{\langle \psi_i | (\delta H)^2 | \psi_i \rangle} \leq \frac{A_i}{\Delta E_i} \|\delta H\|_2, \end{aligned} \quad (21)$$

where we considered the i^{th} -state energy gap for the first inequality:

$$\Delta E_i = \min_{k \neq i} |E_k - E_i|, \quad (22)$$

the reasoning in chapter XI of [37] for the standard-deviation equality, the following quantity for the second equality (defined for $\langle \psi_i | (\delta H)^2 | \psi_i \rangle \neq 0$ which is not constraining ²):

$$A_i = \sqrt{1 - \frac{\langle \psi_i | \delta H | \psi_i \rangle^2}{\langle \psi_i | (\delta H)^2 | \psi_i \rangle}} \in [0, 1], \quad (23)$$

and the relation in (18) for the last inequality.

¹Standard perturbation theory leads to states $|\psi_i^S(\lambda)\rangle = |\psi_i\rangle - \lambda \sum_{k \neq i} \frac{\langle \psi_k | \delta H | \psi_i \rangle}{E_k - E_i} |\psi_k\rangle + \mathcal{O}(\lambda^2)$ that are normalized to first-order in λ , i.e. $\langle \psi_i^S(\lambda) | \psi_i^S(\lambda) \rangle = 1 + \mathcal{O}(\lambda^2)$ and $\langle \psi_i | \psi_i^S(\lambda) \rangle = 1$. This is not suitable for our problematic. The solution is to consider strict normalization perturbation theory [38], [39]. This affects state expansion but not energy expansion, i.e. all results derived in section IV remain the same, which is coherent.

²Having $\langle \psi_i | (\delta H)^2 | \psi_i \rangle = \|\delta H | \psi_i \rangle\|_2^2 = 0$ implies $\delta H | \psi_i \rangle = 0$, thus $\forall n \geq 1 : (\delta H)^n | \psi_i \rangle = 0$, leading to perturbation theory irrelevance for state i .

From (20) and (21), we deduce:

$$\boxed{\frac{\Delta E_i}{A_i} \times T(|\psi_i\rangle, |\psi_i^S(\lambda)\rangle) \lesssim \sqrt{\langle \psi_i | (\lambda \delta H)^2 | \psi_i \rangle} \leq \|\lambda \delta H\|_2} \quad (24)$$

(state condition to 1st-order in λ).

This gives options to upper-bound the state precision.

VI. CONSTRAINING ENERGY PRECISION AND STATE PRECISION TOGETHER

From (16), (19) and (24), we can deduce that constraining:

$$\boxed{\|\lambda \delta H\|_2 \leq \varepsilon \quad (\text{unitary \& energy \& state}),} \quad (25)$$

allows us to upper-bound in one go the precisions on unitaries, energy difference and trace distance (for any state i).

Then, we can deduce using (19) and (24) that both chemical precision ε_{ch} on energy difference and target precision α_{tar} on trace distance (for state i) can be controlled taking:

$$\boxed{\varepsilon = \min \left(\varepsilon_{\text{ch}}, \frac{\Delta E_i}{A_i} \alpha_{\text{tar}} \right) \quad (\text{energy \& state}).} \quad (26)$$

Note that doing the choice $\varepsilon = \varepsilon_{\text{ch}}$ instead of (26) is equivalent to constrain the target precision of state i to a specific precision value α_{ch_i} given by:

$$\alpha_{\text{tar}} \rightarrow \alpha_{\text{ch}_i} = \frac{A_i}{\Delta E_i} \varepsilon_{\text{ch}}. \quad (27)$$

In the context of an usage of QPE for projection in addition to energy estimation, this choice yields a reasonable α_{ch_i} when $\frac{A_i}{\Delta E_i} \leq 1 \text{ Ha}^{-1}$, since then $\alpha_{\text{ch}_i} \lesssim 10^{-3}$, corresponding to a good trace-distance precision. By contrast, if $\frac{A_i}{\Delta E_i} \geq 10 \text{ Ha}^{-1}$, then $\alpha_{\text{ch}_i} > 10^{-2}$, leading to a much poorer precision. In such cases, the usage of the condition in (26) is required to give control on both energy and trace distance precisions.

We highlight that the upper-bounds related to (25) can be loose. Interestingly, we could define from (19), (24) and (26) a tighter upper-bound that can still allow us to control in one go the precisions on energy difference and trace distance:

$$\boxed{\sqrt{\langle \psi_i | (\lambda \delta H)^2 | \psi_i \rangle} \leq \varepsilon \quad (\text{energy \& state, tighter}),} \quad (28)$$

but not anymore the precision on unitaries difference as $\sqrt{\langle \psi_i | (\lambda \delta H)^2 | \psi_i \rangle}$ does not appear in (16). Equation (28) leads to a tighter bound than (25) when the eigen-state i is far from the ‘maximum state’ that defines the spectral norm.

The bound related to (28) can still be loose. Indeed, the trace distance condition demonstration considered ΔE_i instead of the larger $E_{k \neq i} - E_i$ values, remind (21), which can hardly be changed for practical reasons. But, for the energy difference, (17) may lead us to consider (if the 1st-order term is not zero):

$$\boxed{|\langle \psi_i | \lambda \delta H | \psi_i \rangle| \lesssim \varepsilon \quad (\text{energy only, even tighter}),} \quad (29)$$

which does not anymore constrain a precision on unitaries and trace distance, as $|\langle \psi_i | \lambda \delta H | \psi_i \rangle|$ cannot be directly related to

these two quantities. This is why it is sufficient to consider only the chemical precision $\varepsilon = \varepsilon_{\text{ch}}$ in (29), instead of the more constraining form (26). Note that [30] proposed a condition similar to (29) in the context of order-2 Trotterization.

All the conditions developed above are valid for any state i . From now on we consider the ground state $i = 0$, which is usually the main target of QPE applied many-electron systems. ΔE_0 , equation (22), then represents the spectral gap. It is to be noted from (26) that adding the state precision condition can usually be more constraining than using the energy precision condition only, since the spectral gap ΔE_0 can often be small for realistic many-electron systems. Equation (26) raises the question of estimating *a priori* a reasonable value for $\frac{\Delta E_0}{A_0}$ (e.g., from the system giving the QPE initial state or statistical analysis). Similarly, the potential usage of conditions (28) or (29) raises the question of estimating a reasonable approximation of $|\psi_0\rangle$ [30]. Note that the spectral norm condition (25) can be efficiently approximated numerically using stochastic estimation when δH is defined by a LCU [40], which might represent an advantage. An extensive discussion of these aspects goes beyond the scope of this article.

This achieves to detail a unified formalism that allowed us to justify and develop various first-order precision conditions related to approximate implementation of unitaries. Especially, the conditions (24)-(26) and (28) are new to our knowledge. We now explicit clear forms for $\lambda \delta H$ on Trotterization cases.

VII. TROTTERIZATION CASES

Given that H naturally takes a LCU form (7), we have:

$$U^{2^q} = e^{-i2\pi t 2^q H} = e^{-i2\pi t 2^q \sum_{\beta=1}^M \gamma_{\beta} H_{\beta}}, \quad (30)$$

which encourages the usage of order- p Trotterization approximation, denoted by $\mathcal{S}(U^{2^q}, p)$, a common implementation where a parameter $n \in \mathbb{N}^*$ called the ‘number of Trotter steps’ must be set [11], [24], [25]. From the Baker-Campbell-Hausdorff (BCH) formula [11], we have [29], [30]:

$$\mathcal{S}(U^{2^q}, p) = e^{-i2\pi t 2^q H_p^S(\lambda_p)}, \quad \lambda_p = \left(2\pi \frac{t}{n} 2^q \right)^p, \quad (31)$$

where $H_p^S(\lambda_p) = H + \lambda_p \delta H_p + \mathcal{O}(\lambda_p^2)$ represents the effective order- p Trotter Hamiltonian (not to be confused with the order of our previous developments in λ). Equations (25), (28) or (29) can each be rewritten in the form $\lambda_p C_p \lesssim \varepsilon$, where:

$$\boxed{C_p \rightarrow \|\delta H_p\|_2 \quad \text{or} \quad \sqrt{\langle \psi_0 | (\delta H_p)^2 | \psi_0 \rangle} \quad (\text{smaller})} \quad (32)$$

or $|\langle \psi_0 | \delta H_p | \psi_0 \rangle|$ (energy only, even smaller).

Thus, after some manipulations using (31), we can deduce that ensuring the precision conditions (25), (28) or (29) implies:

$$n \gtrsim n_{\min}^{(p)}(q, t), \quad (33)$$

where $n_{\min}^{(p)}(q, t)$ is defined by (10) and C_p can take the explicit forms in (32) in the cases that have been justified here. In other terms, using at least $n_{\min}^{(p)}(q, t)$ Trotter steps computed

from one of the C_p in (32) ensures that the conditions in (16), (19) or (24) are satisfied, depending on the chosen option ³.

Note that the work in [29] can be considered as giving a different and always larger C_p value, that we denote by C'_p , obtained from considerations directly on $\|U^{2^q} - \mathcal{S}(U^{2^q}, p)\|_2$ without involving first-order approximation in λ_p . Our first-order reasoning leads to tighter upper-bounds, which might theoretically have a more limited range of validity but tend to behave better in practice (a numerical illustration will be given in next section). Let us make that explicit using the δH_p form given by order-1 Trotter approximation [29], [30]:

$$\mathcal{S}(U^{2^q}, 1)^{\frac{1}{n}} = \prod_{\beta=1}^M e^{-iH_\beta \gamma_\beta \frac{2\pi t 2^q}{n}}, \quad \lambda_1 = 2\pi \frac{t}{n} 2^q \quad (34)$$

$$\delta H_1 = -\frac{i}{2} \sum_{\alpha=1}^{M-1} \sum_{\beta=\alpha+1}^M [\gamma_\beta H_\beta, \gamma_\alpha H_\alpha].$$

Whatever choice made in (32), we have:

$$C_1 \leq \|\delta H_1\|_2 = \frac{1}{2} \left\| \sum_{\alpha=1}^{M-1} \sum_{\beta=\alpha+1}^M [\gamma_\beta H_\beta, \gamma_\alpha H_\alpha] \right\|_2, \quad (35)$$

whereas the upper-bound obtained in [29] leads to :

$$C'_1 = \frac{1}{2} \sum_{\alpha=1}^{M-1} \left\| \sum_{\beta=\alpha+1}^M [\gamma_\beta H_\beta, \gamma_\alpha H_\alpha] \right\|_2. \quad (36)$$

By triangular inequality, we always have $C_1 \leq C'_1$. A similar conclusion holds for any order- p Trotterization, i.e. $C_p \leq C'_p$ where C'_p is the form obtained in [29] ⁴. We now illustrate numerically various upper-bounds discussed in this article.

VIII. ORDER-1 TROTTERIZATION AND H₂ MOLECULE

We consider the H₂ molecule in the configuration summarized in Table I. All calculations were performed on a Quantum Learning Machine (QLM) from Bull, which enables manipulation of molecular Hamiltonians as well as emulations of quantum processing units using the myQLM package. Exact eigen-energies E_j and eigen-states $|\psi_j\rangle$ were obtained by a full diagonalization of the Hamiltonian. This allowed us to calculate ΔE_0 , A_0 , etc. Spectral norms are computed with the SciPy package [41]. Order-1 Trotterization ($p = 1$) is used, as in (34). QPE is initialized with the Hartree-Fock (HF) state, $|\psi_{\text{init}}\rangle = |\psi_{\text{HF}}\rangle$ (with energy E_{HF}), good enough to ensure QPE's success w.r.t. energy estimation and state projection.

Fig. 1 illustrates the quantities derived above, all rescaled on an energy unit (Ha) to ease comparison, for $q = 0$ (thus

³It is possible in (8) to consider the ε of (26) instead of ε_{ch} . This constrains $N_{\text{min}}(t)$ to be larger than necessary to reach chemical precision on energy estimation, but the qubit overhead remains small (log scaling). The conceptual interest is that the property in (11) then always holds, i.e. a minimum number of Trotter steps over the whole circuit that tends to depend only on a .

⁴Order-2 Trotter approximation implies $\lambda_2 = (2\pi \frac{t}{n} 2^q)^2$ and $\delta H_2 = -\frac{1}{3} \sum_{\alpha=1}^{2M-1} \sum_{\beta=\alpha+1}^{2M} \sum_{\nu=\beta}^{2M} \left(1 - \frac{\delta_{\nu,\beta}}{2}\right) [\gamma_\nu H_\nu, [\gamma_\beta H_\beta, \gamma_\alpha H_\alpha]]$ with $H_{M+i} = H_{M+1-i}$ for $i = 1, \dots, M$. We can demonstrate that the C'_2 formulation in [29] differs from $\|\delta H_2\|_2$ by the 'position' of $\sum_{\nu=\beta}^{2M-1}$ (outside the norm in C'_2), leading by triangular inequality to $C_2 \leq C'_2$, etc.

TABLE I
DATA FOR H₂ IN HA AND ATOMIC UNITS WITH BOND LENGTH 0.5 Å. THE STO-3G BASIS SET IS USED: EACH H IS REPRESENTED WITH A 1S ORBITAL, LEADING TO A SYSTEM WITH $N_S = 4$ QUBITS.

	Parameter	Value
Initial system	E_{init}	-1.042996
Exact system	E_0	-1.055160
	ΔE_0	0.702985
QPE parameters	$t = 1/(2 \sum_{\beta} \gamma_{\beta})$	0.215149
	$\lceil E_0 t \rceil = \lceil E_{\text{init}} t \rceil$	0
	$N_{\text{min}}(t)$	11
Order-1 Trotter. features	A_0	1
	$\ \delta H_1\ _2 = \sqrt{ \langle \psi_0 (\delta H_1)^2 \psi_0 \rangle }$	0.052420
	C'_1	0.196930
	$n_{\text{min}}^{(1)}(0, t)$	30
	$n_{\text{min-tot}}^{(1)}(a = 0)$	6×10^4




Fig. 1. QPE results for H₂ ground state with $t = 1/(2 \sum_{\beta} |\gamma_{\beta}|)$. Plot of the quantities related to (13), (16), (19), (24), (32) and (36) (see caption), as a function of the Trotter steps number n . Trace distance and unitaries difference are rescaled on an energy unit, with factors $\Delta E_0/A_0$ and $1/(2\pi t)$ (resp.). The chemical precision area appears in grey; it can be related to trace distance precision $\alpha_{\text{ch}_0} \leq 2.3 \times 10^{-3}$ (unitless) through (27).

$\lambda = 2\pi \frac{t}{n}$). It can first be seen that $|E_0^S(\lambda) - E_0|$ does not evolve in the same way than other quantities. Indeed, we have in this specific H₂ case $\langle \psi_0 | \delta H_1 | \psi_0 \rangle = 0$ which implies $A_0 = 1$ and a cancellation of the first-order energy term in (17), that invalidates the use of the option (29) (note that [30] used (29) for H₂ in the context of order-2 Trotterization, which is valid as we then have $|\langle \psi_0 | \delta H_2 | \psi_0 \rangle| \neq 0$). This leads for the energy difference to a second-order behavior in $\lambda = 2\pi \frac{t}{n}$, i.e. $\propto (\frac{1}{n})^2$, as observed. All other quantities evolve as a first-order in λ thus in $\frac{1}{n}$. The curves associated to $\|H^S(\lambda) - H\|_2$ and its first-order approximation $\|\delta H_1\|_2$ show that sticking with $\|\delta H_1\|_2$ is largely sufficient as soon as $n \geq 5$. Adding higher-order terms would be needed for small values of n (thus large values of $\lambda = 2\pi \frac{t}{n}$) but these values are not pertinent as they are far from allowing us to reach the chemical precision. The rescaled unitary difference $\|U - \mathcal{S}(U, 1)\|_2$ is of the same order of magnitude than




Fig. 2. QPE results for H_2 ground state with $t = 1/(2\sum_{\beta}|\gamma_{\beta}|)$. Plots (a), (b), (c) are function of the phase qubits number N and correspond to various Trotter step numbers $n(q, t)$ (n should be considered as a function of q and t in coherence with (33)). The evolution of energy differences (red) and trace distance (blue, rescaled on an energy using $\Delta E_0/A_0$) is shown. Straight lines correspond to initial (HF) ground state quantities. The chemical precision area appears in grey; it can be related to trace distance precision $\alpha_{\text{ch}_0} \leq 2.3 \times 10^{-3}$ (unitless) through (27).

$\|\delta H_1\|_2$, which is satisfying. We notice two significant gaps in the curves. The first gap is related to the difference between C'_1 , equation (36), and $C_1 \rightarrow \|\delta H_1\|_2$, equation (35), which reduces by almost an order of magnitude the required number of Trotter steps n to reach a desired precision and can be useful for more accurate resource estimation. Indeed, $C'_1 \approx 4\|\delta H_1\|_2$, which illustrates that the ‘triangular inequality effect’ is non-negligible, especially since this effect is likely to increase with the system size N_S (implying in general more LCU terms). Note that the usually tighter bound in (28) satisfies in our H_2 case the uncommon relation $\sqrt{\langle \psi_0 | (\delta H_1)^2 | \psi_0 \rangle} = \|\delta H_1\|_2$ (illustrations on molecules for which the equality does not hold will be presented in future publications). The second gap lies in the difference between the rescaled trace distance and the energy term, leading to more than an order of magnitude difference on the required n to reach a similar precision. The main reason, already explained above, is that the energy difference behaves as a second-order. More generally, trace distance may tend to converge more slowly than energy difference because of spectral gap ΔE_0 effects.

Fig. 2 illustrates the QPE behavior for various numbers of phase qubits and Trotter steps. We can observe how these parameters allow us to improve the initial state features and reach a desired precision, coherently with our previous considerations. However, the difference between the QPE estimated ground energy and the exact ground energy does not evolve as strongly as in Fig. 1 w.r.t. Trotter step numbers. This is mostly due to our limited N range, which gives a lower-bound on the attainable energy precision, and thus limits the energy difference evolution between panels (b) and (c). If we added more phase qubits, the energy difference would continue to decrease coherently with Fig. 1. On the other hand, the trace distance between the QPE projected state (after

a l^* measure) and the exact ground state seems to reach the Trotterization limit of Fig. 1. Indeed, if the N choice affects directly the energy estimation, it affects less directly the projected state. This is because projection is driven by the $c_j \times f(\theta_j - \frac{l^*}{2N})/\sqrt{P(l^*)}$ terms in (4), which are mostly impacted by N changes through the discretization effects depicted in [13], [28]. The slightly erratic behavior of the trace distance w.r.t. N is related to these effects (they remain limited as the initial state is here good).

IX. CONCLUSION

Overall, this work contributes to providing a clearer picture of the limitations and opportunities of QPE applied to many-electron systems. We focused on a study of the precision related to approximate implementations of the unitaries. We developed various first-order conditions that contribute to control the ground energy estimation precision together with the output state precision, important to leverage the full potential of QPE. The role of the spectral gap ΔE_0 and of A_0 on state precision tends to make requirements on unitaries more stringent for projection than for energy estimation. Applied to Trotterization, our first-order upper bounds are tighter than those obtained in [29] or more general than those obtained in [30]. Our first numerical test on H_2 illustrated these behaviors.

Numerical studies on various many-electron systems are being carried out, among others to estimate the statistical variations of the $A_0/\Delta E_0$ term and evaluate the impact of the options (25), (28) and (29). Numerical methods to efficiently approximate the involved quantities are being evaluated.

ACKNOWLEDGMENT

We warmly thank Y. Chatterjee, H. Calandra, J.-P. Mascomère and O. Ladhari for insightful comments. We thank TotalEnergies for the permission to publish this work.

REFERENCES

- [1] H. Gao, S. Imamura, A. Kasagi, and E. Yoshida, “Distributed implementation of full configuration interaction for one trillion determinants,” *Journal of Chemical Theory and Computation*, vol. 20, no. 3, pp. 1185–1192, 2024.
- [2] D. Zgid, E. Gull, and G. K.-L. Chan, “Truncated configuration interaction expansions as solvers for correlated quantum impurity models and dynamical mean-field theory,” *Physical Review B—Condensed Matter and Materials Physics*, vol. 86, no. 16, p. 165128, 2012.
- [3] W. Kohn and L. J. Sham, “Self-consistent equations including exchange and correlation effects,” *Physical review*, vol. 140, no. 4A, p. A1133, 1965.
- [4] B. Anselme Martin, T. Ayral, F. Jamet, M. J. Rančić, and P. Simon, “Combining matrix product states and noisy quantum computers for quantum simulation,” *Physical Review A*, vol. 109, no. 6, p. 062437, 2024.
- [5] F. Jamet, L. P. Lindoy, Y. Rath, C. Lenihan, A. Agarwal, E. Fontana, F. Simkovic, B. A. Martin, and I. Rungger, “Anderson impurity solver integrating tensor network methods with quantum computing,” *APL Quantum*, vol. 2, no. 1, 2025.
- [6] A. d. Freitas Goncalves, E. Parazzi Lyra, S. Ramdas Chavan, P. L. Llewellyn, L. F. Mercier Franco, and Y. Magnin, “Fundamental of CO₂ adsorption and diffusion in subnanoporous materials: Application to CaF₂,” *The Journal of Physical Chemistry C*, vol. 129, no. 40, pp. 18190–18199, 2025.
- [7] E. P. Lyra and L. F. Franco, “Deriving force fields with a multiscale approach: From ab initio calculations to molecular-based equations of state,” *The Journal of Chemical Physics*, vol. 157, no. 11, 2022.
- [8] G. Greene-Diniz, D. Z. Manrique, W. Sennane, Y. Magnin, E. Shishenina, P. Cordier, P. Llewellyn, M. Krompiec, M. J. Rančić, and D. M. Ramo, “Modelling carbon capture on metal-organic frameworks with quantum computing,” *EPJ Quantum Technology*, vol. 9, no. 1, p. 37, 2022.
- [9] P. W. Shor, “Algorithms for quantum computation: discrete logarithms and factoring,” in *Proceedings 35th annual symposium on foundations of computer science*, pp. 124–134, Ieee, 1994.
- [10] A. Y. Kitaev, “Quantum measurements and the abelian stabilizer problem,” 1995.
- [11] M. A. Nielsen and I. L. Chuang, *Quantum computation and quantum information*. Cambridge university press, 2010.
- [12] D. S. Abrams and S. Lloyd, “Quantum algorithm providing exponential speed increase for finding eigenvalues and eigenvectors,” *Phys. Rev. Lett.*, vol. 83, pp. 5162–5165, Dec 1999.
- [13] B. C. Travaglione and G. J. Milburn, “Generation of eigenstates using the phase-estimation algorithm,” *Physical Review A*, vol. 63, Feb. 2001.
- [14] N. Nusran, *Application of phase estimation algorithms to improve diamond spin magnetometry*. PhD thesis, University of Pittsburgh, 2014.
- [15] P. M. Q. Cruz, G. Catarina, R. Gautier, and J. Fernandez-Rossier, “Optimizing quantum phase estimation for the simulation of hamiltonian eigenstates,” *Quantum Science & Technology*, vol. 5, 2019.
- [16] T. E. O’Brien, B. Tarasinski, and B. M. Terhal, “Quantum phase estimation of multiple eigenvalues for small-scale (noisy) experiments,” *New Journal of Physics*, vol. 21, no. 2, p. 023022, 2019.
- [17] L. Pezzè and A. Smerzi, “Quantum phase estimation algorithm with gaussian spin states,” *PRX Quantum*, 2020.
- [18] L. Lin and Y. Tong, “Heisenberg-limited ground-state energy estimation for early fault-tolerant quantum computers,” *PRX quantum*, vol. 3, no. 1, p. 010318, 2022.
- [19] K. Sugisaki, “Projective measurement-based quantum phase difference estimation algorithm for the direct computation of eigenenergy differences on a quantum computer,” *Journal of Chemical Theory and Computation*, vol. 19, pp. 7617 – 7625, 2023.
- [20] N. J. C. Papadopoulos, J. T. Reilly, J. D. Wilson, and M. J. Holland, “Reductive quantum phase estimation,” *Phys. Rev. Res.*, vol. 6, no. 3, p. 033051, 2024.
- [21] M. Barbieri, I. Gianani, A. Z. Goldberg, and L. L. Sánchez-Soto, “Quantum multiphase estimation,” *Contemporary Physics*, vol. 65, p. 112–124, Apr. 2024.
- [22] J. M. A. Hualde, M. Kowalik, L. Remme, F. E. Wolff, J. van Velzen, W. Killick, R. Bottcher, C. Weimer, J. Krauser, and E. Marsili, “Quantum computing in corrosion modeling: Bridging research and industry,” *arXiv preprint arXiv:2412.07933*, 2024.
- [23] A. Shukla and P. Vedula, “Towards practical quantum phase estimation: A modular, scalable, and adaptive approach,” 2025.
- [24] M. Suzuki, “General theory of fractal path integrals with applications to many-body theories and statistical physics,” *Journal of Mathematical Physics*, vol. 32, pp. 400–407, Feb. 1991.
- [25] A. K. Rajagopal and C. Tsallis, “Generalization of the Lie-Trotter Product Formula for q-Exponential Operators,” *Physics Letters A*, vol. 257, pp. 283–287, July 1999. arXiv:cond-mat/9903106.
- [26] S. Chehade, A. Delgado, S. Wang, and Z. Wang, “Error estimates and higher order trotter product formulas in jordan-banach algebras,” *Linear Algebra and its Applications*, vol. 730, pp. 430–449, 2026.
- [27] C. Ronfaut, R. Ollive, and S. Louise, “Numerical error extraction by quantum measurement algorithm,” arXiv: 2602.01927, 2026.
- [28] W. Sennane and J. Messud, “On the robustness of quantum phase estimation to compute ground properties of many-electron systems,” arXiv:2601.05788, submitted for publication, 2026.
- [29] A. M. Childs, Y. Su, M. C. Tran, N. Wiebe, and S. Zhu, “A Theory of Trotter Error,” *Physical Review X*, vol. 11, p. 011020, Feb. 2021. arXiv:1912.08854 [quant-ph].
- [30] S. G. Mehendale, L. A. Martínez-Martínez, P. D. Kamath, and A. F. Izmaylov, “Estimating Trotter Approximation Errors to Optimize Hamiltonian Partitioning for Lower Eigenvalue Errors,” Aug. 2025. arXiv:2312.13282 [physics].
- [31] S. Fomichev, K. Hejazi, M. S. Zini, M. Kiser, J. Fraxanet, P. A. M. Casares, A. Delgado, J. Huh, A.-C. Voigt, J. E. Mueller, et al., “Initial state preparation for quantum chemistry on quantum computers,” *PRX Quantum*, vol. 5, no. 4, p. 040339, 2024.
- [32] D. Halder, S. P. V., V. Agarwal, and R. Maitra, “Digital quantum simulation of strong correlation effects with iterative quantum phase estimation over the variational quantum eigensolver algorithm: H₄ on a circle as a case study,” 2021.
- [33] N. Bauer and G. Siopsis, “Post-variational ground state estimation via qpe-based quantum imaginary time evolution,” 2025.
- [34] M. A. Nielsen et al., “The fermionic canonical commutation relations and the jordan-wigner transform,” *School of Physical Sciences The University of Queensland*, vol. 59, p. 75, 2005.
- [35] A. M. Childs and N. Wiebe, “Hamiltonian simulation using linear combinations of unitary operations,” arXiv:1202.5822, 2012.
- [36] I. Loaiza, A. M. Khah, N. Wiebe, and A. F. Izmaylov, “Reducing molecular electronic hamiltonian simulation cost for linear combination of unitaries approaches,” *Quantum Science and Technology*, vol. 8, no. 3, p. 035019, 2023.
- [37] C. Cohen-Tannoudji, B. Diu, and F. Laloe, *Mécanique Quantique - Tome 2: Nouvelle édition*. EDP sciences, 2018.
- [38] S.-J. GU, “Fidelity approach to quantum phase transitions,” *International Journal of Modern Physics B*, vol. 24, p. 4371–4458, Sept. 2010.
- [39] D. Rossini and E. Vicari, “Ground-state fidelity at first-order quantum transitions,” *Physical Review E*, vol. 98, Dec. 2018.
- [40] N. S. Blunt, A. V. Ivanov, and A. J. Bay-Smidt, “A Monte Carlo approach to bound Trotter error,” Oct. 2025. arXiv:2510.11621 [quant-ph].
- [41] P. Virtanen, R. Gommers, T. E. Oliphant, M. Haberland, T. Reddy, D. Cournapeau, E. Burovski, P. Peterson, W. Weckesser, J. Bright, S. J. van der Walt, M. Brett, J. Wilson, K. J. Millman, et al, and SciPy 1.0 Contributors, “SciPy 1.0: Fundamental Algorithms for Scientific Computing in Python,” *Nature Methods*, vol. 17, pp. 261–272, 2020.