

Constrained Counterdiabatic Quantum Approximate Optimization Algorithm for Portfolio Optimization

Jose Falla

*Department of Physics and Astronomy
University of Delaware
Newark, DE, USA
jfalla@udel.edu*

Ilya Saфро

*Department of Computer and Information Sciences
Department of Physics and Astronomy
University of Delaware
Newark, DE, USA
isafro@udel.edu*

Abstract—We introduce a counterdiabatic (CD) extension of the Quantum Approximate Optimization Algorithm (QAOA) for constrained portfolio optimization. By incorporating approximate adiabatic gauge potentials generated from nested commutators of the Ising-type portfolio problem Hamiltonian and the Hamming weight-preserving XY mixer Hamiltonian into our variational ansatz, the resulting Constrained Counterdiabatic QAOA (CCD-QAOA) achieves improved optimization performance under realistic budget and risk constraints. Benchmarking against standard XY-mixer QAOA, Grover-mixer QAOA, and penalty-based QAOA formulations, our numerical simulations demonstrate that, for a fixed QAOA depth, our CCD-QAOA approach consistently results in better approximation ratios. **Reproducibility:** Link and data will be available upon acceptance.

Index Terms—Constrained Optimization, Counterdiabatic Driving, Quantum Approximate Optimization Algorithm, Portfolio Optimization

I. INTRODUCTION

Variational quantum algorithms are among the most promising approaches for leveraging near-term and fault-tolerant quantum hardware to address classically challenging optimization problems. The Quantum Approximate Optimization Algorithm (QAOA) [1] has emerged as a particularly versatile framework, providing a controllable interpolation between digitized adiabatic evolution and shallow variational circuits. QAOA has been applied in such areas as finance [2], network science [3], and transportation [4], to mention just a few. Despite its conceptual simplicity and broad applicability, QAOA performance often degrades for constrained combinatorial optimization problems, where competing energy scales and small spectral gaps hinder efficient state preparation.

Many practically relevant optimization tasks are intrinsically constrained. Portfolio optimization, in particular, provides a natural and structured example: risk-return tradeoffs yield dense quadratic interactions, while budget constraints restrict feasible configurations to a narrow solution subspace. Standard penalty-based encodings introduce large energy penalties that distort the spectrum and can suppress effective gaps [5]–[7], while constraint-preserving mixers require problem-specific circuit constructions [8]–[10]. In both cases, the variational landscape can become difficult to navigate (in particular, at low circuit depth).

From the perspective of adiabatic quantum computation, these challenges can be traced to diabatic transitions generated during interpolation between mixing and cost Hamiltonians [11]. Counterdiabatic (CD) driving offers a principled mechanism for suppressing such transitions by augmenting the Hamiltonian with an adiabatic gauge potential (AGP). While exact counterdiabatic terms are generally nonlocal and experimentally inaccessible, variational approximations provide a systematic and hardware-compatible route to incorporating CD corrections into digitized algorithms [12]–[14].

Our contribution: In this work, we develop a counterdiabatic extension of QAOA tailored to constrained optimization problems and demonstrate its performance in the context of portfolio optimization. Our approach constructs approximate gauge potentials variationally within a structured operator algebra generated by the cost and constraint-preserving mixer Hamiltonians. This framework provides (i) a systematic method for generating physically relevant counterdiabatic operators, (ii) a controlled truncation strategy, and (iii) insight into how constraint structure influences ansatz complexity in CD-QAOA.

We show that incorporating counterdiabatic generators into QAOA substantially improves approximation ratios at fixed circuit depth for instances of the portfolio optimization problem using Hamming weight-preserving XY mixers as generators of adiabatic gauge potentials. Beyond the specific application to portfolio optimization, our results illustrate a broader principle: constrained combinatorial optimization problems possess structured operator algebras that can be leveraged to construct efficient counterdiabatic ansätze. This connection between adiabatic gauge potentials and variational quantum circuits provides a unifying framework for enhancing QAOA performance in structured optimization landscapes.

II. BACKGROUND

A. The Quantum Approximate Optimization Algorithm

The Quantum Approximate Optimization Algorithm provides a variational framework for approximately preparing the ground state of a problem Hamiltonian encoding a combinatorial optimization task [1]. Starting from an easy-to-prepare reference state, QAOA constructs a parametrized quantum

circuit consisting of alternating applications of cost and mixing unitaries.

At its core, QAOA translates a classical cost function into a Hermitian operator known as the *problem Hamiltonian* H_C , whose lowest-energy state encodes the desired solution. The algorithm proceeds by alternating unitary evolutions generated by H_C and a complementary *mixer Hamiltonian* H_M . The role of H_M is to drive transitions between computational basis states, thereby enabling exploration of the solution space and helping to avoid entrapment in suboptimal configurations. Despite its elegant formulation, QAOA encounters several practical limitations. Chief among these is the difficulty of optimizing its variational parameters using classical routines, particularly as the circuit depth p increases. The resulting optimization landscape is typically highly nonconvex, leading to significant computational overhead that can erode any potential quantum advantage [15]. A well-known manifestation of this challenge is the presence of *barren plateaus*, where gradients vanish exponentially with system size, rendering training ineffective [16], [17].

Specifically, for a circuit of depth p , the QAOA state is defined as

$$|\psi(\gamma, \beta)\rangle = \prod_{k=1}^p e^{-i\beta_k H_M} e^{-i\gamma_k H_C} |\psi_0\rangle, \quad (1)$$

where H_C is the cost Hamiltonian encoding the solution to the combinatorial optimization problem, H_M is a mixing Hamiltonian, and $|\psi_0\rangle$ is the initial state, usually taken to be the uniform superposition of the computational basis state on N qubits. The parameters $\gamma = (\gamma_1, \dots, \gamma_p)$ and $\beta = (\beta_1, \dots, \beta_p)$ are optimized classically to minimize the expectation value

$$\langle \psi(\gamma, \beta) | H_C | \psi(\gamma, \beta) \rangle. \quad (2)$$

The standard choice of mixer is the transverse-field Hamiltonian

$$H_M = \sum_{i=1}^N X_i, \quad (3)$$

where X_i is the Pauli- X operator acting on qubit i . This operator generates transitions between computational basis states and enables exploration of the Hilbert space during the variational optimization.

The corresponding initial state is typically chosen as the uniform superposition

$$|\psi_0\rangle = |+\rangle^{\otimes N}, \quad (4)$$

which is the ground state of H_M . In the limit $p \rightarrow \infty$ with small steps, QAOA reproduces continuous adiabatic evolution [1].

QAOA can be viewed as a Trotterized approximation to adiabatic evolution between the mixer and cost Hamiltonians. Consider the time-dependent Hamiltonian

$$H(s) = (1-s)H_M + sH_C, \quad (5)$$

where $s \in [0, 1]$ parametrizes the interpolation. In the limit $p \rightarrow \infty$, QAOA approaches a digitized version of adiabatic evolution under $H(s)$, provided the parameters (γ_k, β_k) approximate the continuous schedule.

For shallow circuits, however, the evolution is necessarily nonadiabatic, and diabatic transitions can significantly reduce the overlap between the prepared state and the ground state of H_C .

QAOA performance has an evident dependence on the circuit depth and, as previously mentioned, it is observed that increasing depth improves the quality of the possible approximation (at the cost of increasing the parameter search space). Furthermore, finding appropriate solutions has a strong dependence on the ratio of a problem's constraints to variables (problem density). Hence, QAOA exhibits strong dependence on a problems density, and, for any fixed ansatz, there exists problem instances of high-density that appear not to be accessible. This feature persists as a fundamental limitation exhibited by QAOA [18].

Additionally, constrained optimization problems present several challenges for standard QAOA implementations. First, when constraints are implemented through penalty terms, the cost Hamiltonian contains energy scales associated with both the optimization objective and constraint enforcement [19]. Large penalty coefficients distort the spectrum of H_C and can produce small effective gaps along the interpolation path, making it difficult for low-depth circuits to approximate the ground state.

Second, the standard transverse-field mixer does not preserve constraints such as fixed Hamming weight [9]. As a result, the QAOA state may explore infeasible regions of the Hilbert space during the optimization. Although penalty terms suppress such states energetically, this mechanism can lead to inefficient exploration and slower convergence of the classical optimizer.

Finally, the dense interaction structure arising from covariance matrices in portfolio optimization generates a complex energy landscape with many competing configurations. In such settings, low-depth QAOA circuits may struggle to capture the correlations required to approximate the optimal portfolio [20].

B. Counterdiabatic Corrections

From the perspective of adiabatic dynamics, the limitations above can be understood as consequences of diabatic transitions induced by the finite-depth evolution. When the interpolation between H_M and H_C encounters small spectral gaps, transitions out of the instantaneous ground state become likely, reducing the final state fidelity.

Counterdiabatic (CD) driving provides a principled mechanism for suppressing such transitions by augmenting the Hamiltonian with additional terms that cancel diabatic excitations [12]. While the exact counterdiabatic Hamiltonian is generally nonlocal, variational approximations based on the

adiabatic gauge potential offer a tractable route for incorporating these corrections into variational quantum circuits [21].

Consider a family of Hamiltonians $H(\lambda)$ parametrized by a control parameter λ . In adiabatic quantum evolution, the system is evolved according to $H(\lambda(t))$ with λ slowly varying in time. Addition of a counterdiabatic Hamiltonian term to the adiabatic component results in

$$H(\lambda) = (1 - \lambda(t))H_i + \lambda(t)H_f + f(\lambda)H_{CD}, \quad (6)$$

where H_i and H_f are the initial and problem Hamiltonians ($H_{AD} = (1 - \lambda(t))H_i + \lambda(t)H_f$), respectively, and H_{CD} is the counterdiabatic term with scheduling $f(\lambda)$, which vanishes at the beginning and end of the protocol [14].

Inclusion of the exact CD term allows following the instantaneous ground state of the original Hamiltonian at all times during the evolution. Therefore, the CD Hamiltonian term can be defined as,

$$H_{CD} = \dot{\lambda}A_\lambda, \quad (7)$$

where A_λ is known as the adiabatic gauge potential (AGP). When the evolution is very fast, H_{CD} increases dramatically, and in the adiabatic limit this term vanishes. The adiabatic gauge potential is defined as

$$A_\lambda = i \sum_m (1 - |m(\lambda)\rangle \langle m(\lambda)|) |\partial m(\lambda)\rangle \langle m(\lambda)|, \quad (8)$$

corresponding to the nonadiabatic transitions between eigenstates $|m(\lambda)\rangle$ [22]. The AGP satisfies the condition

$$[i\partial H_{AD} - [A_\lambda, H_{AD}], H_{AD}] = 0. \quad (9)$$

Obtaining the exact form for A_λ is challenging since, in many cases, it contains exponentially many terms with nonlocal many-body interactions. In many cases, it suffices to employ the approximate form of the adiabatic gauge potential, which can be obtained from a nested commutator

$$A_\lambda^{(l)} = i \sum_{k=1}^l \alpha_k(t) \underbrace{[H_{ad}, [H_{ad}, \dots [H_{ad}, \partial_\lambda H_{ad}]]}_{2k-1}, \quad (10)$$

where $\alpha_k(t)$ is the counterdiabatic coefficient. This coefficient is obtained by minimizing the operator distance between the exact gauge potential and the approximate gauge potential. In some sense, we construct the approximate gauge potential by restricting A_λ to a variational operator ansatz

$$A_\lambda^* = \sum_k \alpha_k(\lambda) O_k, \quad (11)$$

where $\{O_k\}$ is a chosen operator basis and α_k are variational coefficients.

Following the variational principle introduced in the context of adiabatic gauge potentials, the optimal coefficients are obtained by minimizing the Frobenius norm of the Hilbert-Schmidt operator

$$G(A_\lambda) = \partial_\lambda H_{AD} + i[A_\lambda, H_{AD}], \quad (12)$$

which corresponds to minimizing the action

$$S(A_\lambda) = \text{Tr} [G(A_\lambda)^2]. \quad (13)$$

This yields a linear system of equations for the coefficients α_k . The resulting operator A_λ^* provides the best approximation to the exact gauge potential within the chosen operator subspace.

Once an approximate gauge potential has been obtained, to incorporate these corrections into a variational circuit, evolution is digitized in a manner analogous to QAOA. Specifically, the counterdiabatic QAOA ansatz augments the standard alternating structure with unitaries generated by the approximate gauge potential operators.

For circuit depth p , the resulting variational state is

$$|\psi_{CD}\rangle = \prod_{k=1}^p \prod_j e^{-i\eta_{k,j} O_j} e^{-i\beta_k H_M} e^{-i\gamma_k H_C} |\psi_0\rangle, \quad (14)$$

where $\eta_{k,j}$ are additional variational parameters associated with the counterdiabatic generators O_j .

These additional operators compensate for diabatic transitions that occur in shallow QAOA circuits, effectively increasing the expressive power of the variational ansatz without requiring a proportional increase in circuit depth.

A crucial ingredient in the above construction is the choice of operator basis $\{O_k\}$. In principle, the exact gauge potential can involve highly nonlocal Pauli strings, making a naive operator expansion impractical. However, the structure of the cost Hamiltonian H_C strongly constrains the form of operators that appear in the nested commutator expansion of the gauge potential.

C. Constrained Portfolio Optimization as a Quantum Hamiltonian

Portfolio optimization is a canonical problem in quantitative finance, originating from the Markowitz mean-variance framework [7]. The goal is to construct a portfolio that balances expected return against risk, where risk is typically quantified by the covariance of asset returns.

Consider a universe of N assets with expected return vector $\boldsymbol{\mu} = (\mu_1, \dots, \mu_N)$ and covariance matrix Σ . In the classical mean-variance formulation, the objective is to minimize the risk-adjusted cost function

$$C(\mathbf{x}) = \lambda \sum_{i,j=1}^N \Sigma_{ij} x_i x_j - \sum_{i=1}^N \mu_i x_i, \quad (15)$$

where $x_i \in \{0, 1\}$ denotes whether asset i is included in the portfolio and λ controls the tradeoff between return and risk. The first term penalizes correlated selections that increase portfolio variance, while the second term favors assets with high expected returns.

In many practical settings, the portfolio must satisfy a budget constraint specifying the number of selected assets. This can be expressed as

$$\sum_{i=1}^N x_i = B, \quad (16)$$

where B is the desired portfolio size.

One common approach for satisfying the budget is to enforce this constraint through a quadratic penalty term, yielding the unconstrained objective

$$C_P(\mathbf{x}) = \lambda \sum_{i,j} \Sigma_{ij} x_i x_j - \sum_i \mu_i x_i + \alpha \left(\sum_i x_i - B \right)^2, \quad (17)$$

where α is a penalty coefficient chosen large enough to ensure feasible solutions satisfy the constraint.

Alternatively, one may restrict the quantum evolution to the feasible subspace defined by $\sum_i x_i = B$, for example by designing mixers that preserve Hamming weight [23].

To encode the optimization problem in a quantum algorithm, we map the binary variables x_i to spin-1/2 degrees of freedom using the standard transformation

$$x_i = \frac{1 - Z_i}{2}, \quad (18)$$

where Z_i is the Pauli- Z operator acting on qubit i .

Substituting this relation into the cost function produces a Hamiltonian whose ground state corresponds to the optimal portfolio. Up to a constant energy shift, the resulting cost Hamiltonian takes the form

$$H_C = \sum_{i < j} J_{ij} Z_i Z_j + \sum_i h_i Z_i, \quad (19)$$

where the coupling coefficients J_{ij} arise from the covariance matrix and constraint penalties, and the local fields h_i encode contributions from expected returns and linear terms generated by the constraint expansion [5], [24].

The Hamiltonian H_C therefore has the structure of a fully connected Ising model with pairwise ZZ interactions and local Z fields. Explicitly, the interaction coefficients are determined by

$$J_{ij} = \frac{\lambda}{2} \Sigma_{ij}, \quad (20)$$

for $i < j$, up to convention and penalty terms, while the local fields depend on both the return vector and the constraint parameters.

This form is directly compatible with QAOA and related variational algorithms, where the cost unitary is generated by H_C . Importantly, the dense interaction graph induced by the covariance matrix and the additional constraint penalties produces a complex energy landscape, making this class of problems a useful testbed for studying improved variational ansätze, such as counterdiabatic QAOA.

III. RELATED WORK

Several well-known challenges limit the practical performance of QAOA, including the cost of repeated classical optimization, the rapid growth of the variational parameter space with circuit depth, and deteriorating scalability on large constrained instances [25]. A number of improvements have therefore been proposed to address these issues. Parameter transferability is one of the most effective acceleration mechanisms, as optimized angles can often be reused across related instances, depths, or problem sizes, reducing and sometimes largely avoiding expensive classical optimization [26]–[28]. Other works exploit problem symmetries and invariant subspaces to identify redundant parameters and reduce the effective dimensionality of the QAOA ansatz [29]–[32]. Initialization plays a similarly important role: warm-start strategies (e.g., problem-informed initial states, and structured parameter seeding) [33], [34], and initialization schemes based on flexible distributions such as the beta-distribution-based [35] can significantly improve optimization quality and trainability. To extend QAOA to larger instances, several hybrid methods combine it with decomposition frameworks that partition a large optimization problem into smaller subproblems and then coordinate their solutions [36], [37].

For optimization problems with hard constraints, the standard transverse-field mixer is generally insufficient because it explores infeasible regions of the Hilbert space. To address this limitation, Hadfield *et al.* introduced the Quantum Alternating Operator Ansatz (QAOA⁺), which generalizes QAOA by allowing problem-specific mixers that preserve feasibility [38]. Among these, the XY mixer has become one of the most important constructions for Hamming-weight constrained problems because it preserves particle number (or budget) exactly. Analytical and numerical studies of XY mixers were further developed by Wang *et al.*, who demonstrated their effectiveness for constrained optimization and one-hot encodings [10]. Subsequent work has refined the theory and compilation of constrained mixers, including efficient Trotterized implementations and generalizations beyond the standard ring and complete-graph topologies [9].

These constrained mixer ideas have been particularly impactful in finance applications, especially portfolio optimization, where cardinality and budget constraints naturally define a fixed Hamming-weight subspace. Recent benchmarking studies have shown that XY -mixer QAOA provides significant advantages over unconstrained mixers for portfolio instances [8], [20]. In particular, the importance of initializing in the ground state of the XY mixer (Dicke state) has been linked to improved trainability and better alignment with the adiabatic limit [20].

Parallel to the development of QAOA, the quantum control and adiabatic computing communities have established the framework of *counterdiabatic driving* (CD) and *shortcuts to adiabaticity*, in which additional Hamiltonian terms suppress diabatic transitions during finite-time evolution [12]. A central object in this theory is the adiabatic gauge potential (AGP),

which generates the exact counterdiabatic correction. Because exact AGPs are typically intractable, Sels and Polkovnikov introduced a variational principle for constructing approximate local AGPs by minimizing an action based on the operator commutator structure [39]. This variational AGP framework has since become a standard route for practical counterdiabatic approximations in many-body systems.

The connection between QAOA and counterdiabatic driving has motivated several recent attempts to augment variational quantum optimization with AGP-inspired terms. These works generally show that incorporating approximate counterdiabatic corrections can improve optimization performance at low circuit depth by emulating faster adiabatic transport.

IV. METHODS

For the portfolio optimization problem, we wish to enforce the budget constraint $\sum_i x_i = B$ exactly. To this end, we restrict the dynamics to the fixed-Hamming-weight subspace using an XY mixer. The mixer Hamiltonian is defined as

$$H_M^{XY} = \sum_{i < j} (X_i X_j + Y_i Y_j), \quad (21)$$

which preserves the total excitation number. As a result, the evolution generated by H_M^{XY} remains entirely within the feasible subspace of valid portfolios.

The QAOA circuit is initialized in a Dicke state of Hamming weight B ,

$$|D_B^N\rangle = \binom{N}{B}^{-1/2} \sum_{\substack{z \in \{0,1\}^N \\ |z|=B}} |z\rangle, \quad (22)$$

which is an equal superposition over all feasible configurations. This state can be prepared efficiently using known state preparation protocols and serves as natural eigenstate of H_M^{XY} within the fixed-weight subspace [40].

We consider the interpolating Hamiltonian

$$H(\lambda) = (1 - \lambda)H_M^{XY} + \lambda H_C, \quad (23)$$

where, for the portfolio optimization problem, H_C has the form of Eq. 19. The commutator

$$[H_C, H_M^{XY}] \quad (24)$$

generates operators of the form

$$X_i Y_j Z_k, \quad Y_i X_j Z_k, \quad (25)$$

which correspond to three-body Pauli strings. Motivated by this structure, we truncate the gauge potential to the form

$$A_\lambda^* = \sum_{i < j} \alpha_{ij} (X_i Y_j - Y_i X_j) + \sum_{i < j < k} \beta_{ijk} O_{ijk}, \quad (26)$$

where O_{ijk} denotes three-body Pauli strings of the form

$$O_{ijk} \in \{X_i Y_j Z_k - Y_i X_j Z_k, \text{ permutations}\}. \quad (27)$$

The first term corresponds to two-body generators already present in the XY mixer algebra, while the second captures leading-order corrections arising from interactions with the cost Hamiltonian.

To illustrate how higher-body operators arise in the gauge potential, we consider a representative commutator between a term in the cost Hamiltonian and a term in the XY mixer.

Let $H_C \supset J_{ij} Z_i Z_j$ and $H_M^{XY} \supset X_i X_k + Y_i Y_k$, where i, j, k are distinct. Using Pauli commutation relations, we compute

$$[Z_i Z_j, X_i X_k] = Z_j [Z_i, X_i] X_k \quad (28)$$

$$= Z_j (2i Y_i) X_k \quad (29)$$

$$= 2i Y_i X_k Z_j. \quad (30)$$

Similarly,

$$[Z_i Z_j, Y_i Y_k] = Z_j [Z_i, Y_i] Y_k \quad (31)$$

$$= Z_j (-2i X_i) Y_k \quad (32)$$

$$= -2i X_i Y_k Z_j. \quad (33)$$

Combining these contributions, we obtain

$$[Z_i Z_j, X_i X_k + Y_i Y_k] = 2i (Y_i X_k - X_i Y_k) Z_j, \quad (34)$$

which is a three-body Pauli operator. This example demonstrates that commutators between the cost Hamiltonian and the XY mixer naturally generate operators of the form $X_i Y_k Z_j - Y_i X_k Z_j$, motivating the inclusion of three-body terms in the gauge potential ansatz.

The coefficients α_{ij} and β_{ijk} are obtained by minimizing the action

$$\mathcal{S} = \text{Tr} \left[(\partial_\lambda H + i[A_\lambda, H])^2 \right]. \quad (35)$$

Substituting the ansatz for A_λ^* into this expression yields a system of linear equations of the form

$$\sum_l M_{kl} c_l = v_k, \quad (36)$$

where $c_l \in \{\alpha_{ij}, \beta_{ijk}\}$, and

$$M_{kl} = -\text{Tr}([O_k, H][O_l, H]), \quad (37)$$

$$v_k = i \text{Tr}(O_k[H, \partial_\lambda H]). \quad (38)$$

This linear system determines the optimal coefficients within the chosen operator subspace.

For the portfolio Hamiltonian considered here, the structure of H_C implies that the coefficients α_{ij} depend primarily on local fields and pairwise couplings involving qubits i and j , while the three-body coefficients β_{ijk} encode higher-order correlations induced by the covariance matrix.

To implement this evolution in a variational circuit, we digitize the time evolution into p layers, yielding the ansatz




Fig. 1. Schematic of Counterdiabatic QAOA. For the portfolio optimization problem, the basis computational state is initialized to the fixed-Hamming-weight Dicke state. Subsequently, p layers of the cost, mixer, and counterdiabatic unitaries are applied. A final measurement in the computational basis is made, and the variational parameters are optimized and updated classically.

$$|\psi_{CD}\rangle = \prod_{k=1}^p e^{-i\eta_k H_{CD}} e^{-i\beta_k H_M^{XY}} e^{-i\gamma_k H_C} |D_B^N\rangle, \quad (39)$$

where standard QAOA is augmented by three-body counterdiabatic generators. While the non-counterdiabatic XY-mixer approach preserves Hamming weight, adding these counterdiabatic contributions will result in leakage into the infeasible subspace. This is further analyzed in the following section.

The additional parameters η_k allow the circuit to approximate counterdiabatic evolution while retaining the flexibility of a variational ansatz. In practice, we find that truncating the operator expansion to include only up to three-body terms provides a favorable tradeoff between expressibility and circuit complexity.

V. NUMERICAL RESULTS

We now benchmark the performance of counterdiabatic QAOA for constrained portfolio optimization and compare it against standard QAOA implemented with the same XY mixer and fixed-Hamming-weight initialization (Dicke state).

Unless otherwise stated, all numerical simulations are performed on randomly generated portfolio instances, using Qiskit Finance, with $N = 12$ assets, expected return vector μ sampled from a uniform distribution, and covariance matrices Σ generated as dense positive semidefinite matrices. The portfolio size is fixed to Hamming weight $B = 4$, and all simulations are restricted to this feasible subspace.

Furthermore, we employ the conditional value-at-risk (CVAR) technique, which deviates from the traditional sampling of the mean in measurement outcomes. We use a CVAR approach since problems with a diagonal Hamiltonian, such as

combinatorial optimization problems, the sample mean may be a poor choice because when H is diagonal, there exists a ground state which is a basis state [41].

All numerical simulations were performed by augmenting *OpenQuantumComputing*'s implementation of QAOA [42] with a counterdiabatic routine.

We first study the approximation ratio as a function of QAOA depth p . For each instance, we define

$$r = \frac{\langle H_C \rangle - E_{\max}}{E_{\min} - E_{\max}}, \quad (40)$$

where $E_{\min}$ and $E_{\max}$ denote the minimum and maximum energies within the feasible subspace.

Figure 2 shows a comparison in approximation ratio for different approaches to constrained portfolio optimization. The methods presented are standard XY-mixer QAOA, XY-mixer CD QAOA (our approach), Grover mixer QAOA [43], and penalty-based QAOA. For the XY mixers, we employ both chain and ring structures [10]. Furthermore, this figure shows approximation ratios for different conditional values at risk. The figure shows that CD-QAOA consistently achieves higher approximation ratios at fixed p . For all methods, a smaller CVAR value results in a better approximation ratio.

To quantify the ability of the variational circuits to prepare the optimal portfolio state, we compute the success probability

$$P_{GS} = |\langle \psi_{\text{opt}} | \psi \rangle|^2, \quad (41)$$

where $|\psi_{\text{opt}}\rangle$ is the exact ground state of the portfolio Hamiltonian in the feasible subspace.

A key aspect of Fig. 3 is that it points to a limitation of including counterdiabatic driving to XY-mixer QAOA. By constructing the CD Hamiltonian with three-body terms, we are




Fig. 2. Comparison of approximation ratios for XY mixers (with and without CD contributions), penalty-based constrained QAOA, and Grover mixer QAOA for different conditional values at risk.




Fig. 3. Comparison of success probabilities for XY mixers (with and without CD contributions), penalty-based constrained QAOA, and Grover mixer QAOA for different conditional values at risk.

effectively leaking into the infeasible subspace because these terms are not Hamming-weight preserving. By comparison, an XY-mixer approach that does not include a CD term always results in a success probability of 1, as all sampled bit strings are feasible solutions. Nevertheless, we see that, in general, our approach produces better success probabilities compared to a penalty-based approach, which is also Hamming-weight non-preserving.

Including three-body CD correction terms to our XY-mixer architecture increases the complexity of the ansatz. This is evidenced in an increased optimization runtime as well as an increase in the transpiled circuit depth and number of CNOT gates.

Figures 4 and 5 illustrate the scaling of runtime, CNOT




Fig. 4. Incremental and cumulative runtime for all constrained optimization methods at a fixed CVAR value of 1.

gate count, total two-qubit gate count, and transpiled circuit depth for increasing QAOA depth. As expected, the runtime in the counterdiabatic regime exceeds runtime for any other method; not only is there a quantum overhead in the number of necessary gates, but also an additional variational parameter to be optimized, leading to classical overhead as well. This scalability issue could be addressed by, for example, warm starting the QAOA instance [44], [45].

On the other hand, while we see an increase in the number of CNOT and total two-qubit gates, and in the transpiled circuit depth, the CD approach remains scalable when compared to Grover-mixer QAOA. A possible solution to reduce the total number of two-qubit gates in the CD regime is to adaptively select a subset of the three-body CD operator pool, akin to ADAPT-QAOA [46].

VI. DISCUSSION

The numerical results presented above demonstrate that incorporating counterdiabatic corrections into constrained QAOA provides systematic performance improvements across portfolio instances, particularly in the shallow-circuit regime. From a dynamical perspective, this improvement can be understood as a direct consequence of suppressing diabatic transitions along the interpolation path between the XY mixer and the portfolio cost Hamiltonian.




Fig. 5. Scaling of CNOT gate count, total two-qubit gate count, and transpiled circuit depth as a function of QAOA depth for all constrained optimization methods at a fixed CVAR value of 1.

The use of an XY mixer restricts the dynamics to the fixed-Hamming-weight subspace, ensuring exact preservation of the budget constraint. And while adding a CD term causes leakage to the infeasible subspace, the success probability is improved over a penalty-based method. Within this constrained manifold, the interpolation encounters nontrivial spectral structure generated by the dense covariance matrix and the risk-return tradeoff terms. In particular, near-degenerate excited states frequently arise from competing portfolio configurations with similar variance profiles. At low circuit depth, standard QAOA lacks sufficient expressive power to resolve these avoided crossings, leading to diabatic population transfer into suboptimal feasible states.

The variationally constructed adiabatic gauge potential compensates for these transitions by introducing operators that locally rotate the instantaneous eigenbasis of the interpolating Hamiltonian. The commutator-based derivation makes this mechanism explicit: the leading corrections generated by the

XY mixer and the pairwise Ising cost terms naturally produce three-body Pauli strings. These terms encode correlated excitation transport conditioned on the state of neighboring qubits, allowing the circuit to capture higher-order dependencies in the covariance graph.

This observation provides important physical intuition for why the three-body truncation performs particularly well. While the original portfolio Hamiltonian contains only pairwise ZZ couplings, the instantaneous eigenbasis rotates in a correlated many-body manner under the XY dynamics. The three-body counterdiabatic terms represent the lowest-order operators capable of capturing this correlated basis rotation, explaining their disproportionate impact on shallow-depth performance.

More broadly, our results suggest that constrained combinatorial optimization problems possess a natural operator hierarchy. Rather than relying on heuristic ansatz design, the counterdiabatic framework provides a systematic route to identifying the most physically relevant generators needed to suppress diabatic errors. This perspective may be particularly valuable for other fixed-subspace optimization problems, including scheduling, routing, and cardinality-constrained quadratic programs.

An important practical implication concerns circuit complexity. Although the full gauge potential generally contains operators of increasing Pauli weight, our results indicate that truncation at three-body terms already captures the dominant corrections for the portfolio instances studied here. This suggests a favorable expressibility-to-complexity tradeoff in which significant gains can be achieved without the prohibitive overhead associated with deeper standard QAOA circuits.

Finally, the observed improvements in classical optimizer convergence suggest that counterdiabatic corrections reshape the variational landscape in addition to improving circuit expressibility. A more detailed analysis of trainability, gradient concentration, and spectral geometry in constrained CD-QAOA remains an important direction for future work.

VII. CONCLUSION

In this work, we introduced a counterdiabatic extension of the Quantum Approximate Optimization Algorithm for constrained portfolio optimization. By combining an XY mixer, Dicke-state initialization, and a variational approximation to the adiabatic gauge potential, we constructed an ansatz that systematically improves upon standard QAOA.

A key result of this work is the explicit construction of the gauge-potential operator pool generated by the portfolio cost Hamiltonian and the XY mixer. This analysis shows that the leading counterdiabatic corrections naturally include both two-body and three-body Pauli operators, with the latter playing a particularly important role in capturing correlated diabatic effects within the feasible subspace. Our numerical simulations demonstrate that these counterdiabatic terms lead to improved approximation ratios and superior shallow-depth scaling across portfolio instances with dense covariance structure.

Beyond the specific application to portfolio optimization, our results establish a general framework for enhancing constrained variational quantum algorithms through systematically derived counterdiabatic corrections. The connection between adiabatic gauge potentials and structured optimization Hamiltonians provides a broadly applicable methodology for constructing physically motivated ansätze in constrained Hilbert spaces.

Future work includes extending this framework to larger financial optimization models using classically driven decomposition [47], operator truncation strategies with adaptive [48] and generative learning [49] methods, and hardware-efficient decompositions of the three-body counterdiabatic terms. More generally, the present results suggest that combining constraint-preserving mixers with truncated gauge-potential expansions may offer a scalable path toward improved quantum optimization on near-term devices.

ACKNOWLEDGMENT

This work was supported in part by NSF award #2427042.

REFERENCES

- [1] Edward Farhi, Jeffrey Goldstone, and Sam Gutmann. A Quantum Approximate Optimization Algorithm, November 2014. arXiv:1411.4028 [quant-ph].
- [2] Dylan Herman, Cody Googin, Xiaoyuan Liu, Yue Sun, Alexey Galda, Ilya Safro, Marco Pistoia, and Yuri Alexeev. Quantum computing for finance. *Nature Reviews Physics*, 5(8):450–465, 2023.
- [3] Ruslan Shaydulin, Hayato Ushijima-Mwesigwa, Ilya Safro, Susan Mniszewski, and Yuri Alexeev. Network community detection on small quantum computers. *Advanced Quantum Technologies*, 2(9):1900029, 2019.
- [4] Utkarsh Azad, Bikash K Behera, Emad A Ahmed, Prasanta K Panigrahi, and Ahmed Farouk. Solving vehicle routing problem using quantum approximate optimization algorithm. *IEEE Transactions on Intelligent Transportation Systems*, 2022.
- [5] Zhijie Tang, Alex Lu Dou, and Arit Kumar Bishwas. Comparative analysis of diverse methodologies for portfolio optimization leveraging quantum annealing techniques, July 2024. arXiv:2403.02599 [quant-ph].
- [6] Zain H. Saleem, Teague Tomesh, Bilal Tariq, and Martin Suchara. Approaches to Constrained Quantum Approximate Optimization. *SN COMPUT. SCI.*, 4(2):183, January 2023.
- [7] Iordanis Kerenidis, Anupam Prakash, and Dániel Szilágyi. Quantum Algorithms for Portfolio Optimization. In *Proceedings of the 1st ACM Conference on Advances in Financial Technologies*, pages 147–155, Zurich Switzerland, October 2019. ACM.
- [8] Sebastian Brandhofer, Daniel Braun, Vanessa Dehn, Gerhard Hellstern, Matthias Hüls, Yanjun Ji, Iliia Polian, Amandeep Singh Bhatia, and Thomas Wellens. Benchmarking the performance of portfolio optimization with QAOA. *Quantum Inf Process*, 22(1):25, December 2022.
- [9] Franz G. Fuchs, Kjetil Olsen Lye, Halvor Møll Nilsen, Alexander J. Stasik, and Giorgio Sartor. Constrained mixers for the quantum approximate optimization algorithm. *Algorithms*, 15(6):202, June 2022. arXiv:2203.06095 [quant-ph].
- [10] Zhihui Wang, Nicholas C. Rubin, Jason M. Dominy, and Eleanor G. Rieffel. X Y mixers: Analytical and numerical results for the quantum alternating operator ansatz. *Phys. Rev. A*, 101(1):012320, January 2020.
- [11] M. H. S. Amin. Consistency of the Adiabatic Theorem. *Phys. Rev. Lett.*, 102(22):220401, June 2009.
- [12] D. Guéry-Odelin, A. Ruschhaupt, A. Kiely, E. Torrontegui, S. Martínez-Garaot, and J.G. Muga. Shortcuts to adiabaticity: Concepts, methods, and applications. *Rev. Mod. Phys.*, 91(4):045001, October 2019.
- [13] Ieva Cepaitė, Anatoli Polkovnikov, Andrew J. Daley, and Callum W. Duncan. Counterdiabatic Optimized Local Driving. *PRX Quantum*, 4(1):010312, January 2023.
- [14] Narendra N. Hegade, Xi Chen, and Enrique Solano. Digitized counterdiabatic quantum optimization. *Phys. Rev. Research*, 4(4):L042030, November 2022.
- [15] L. Zhou, S. Wang, S. Choi, P. Hanne, and M. Lukin. Quantum approximate optimization algorithm: Performance, mechanism, and implementation on near-term devices. *Phys. Rev. X*, 10, 2020.
- [16] Jarrod R McClean, Sergio Boixo, Vadim N Smelyanskiy, Ryan Babbush, and Hartmut Neven. Barren plateaus in quantum neural network training landscapes. *Nature communications*, 9(1):1–6, 2018.
- [17] M. Larocca, S. Thanasilp, S. Wang, K. Sharma, J. Biamonte, P. J. Coles, L. Cincio, J. R. McClean, and Z. Holmes. Barren plateaus in variational quantum computing. *Nat. Rev. Phys.*, pages 1–16, 2025.
- [18] V. Akshay, H. Philathong, M.E.S. Morales, and J.D. Biamonte. Reachability Deficits in Quantum Approximate Optimization. *Phys. Rev. Lett.*, 124(9):090504, March 2020.
- [19] David Bucher, Daniel Porawski, Maximilian Janetschek, Jonas Stein, Corey O’Meara, Giorgio Cortiana, and Claudia Linnhoff-Popien. Efficient QAOA Architecture for Solving Multi-Constrained Optimization Problems, June 2025. arXiv:2506.03115 [quant-ph].
- [20] Zichang He, Ruslan Shaydulin, Shouvanik Chakrabarti, Dylan Herman, Changhao Li, Yue Sun, and Marco Pistoia. Alignment between initial state and mixer improves QAOA performance for constrained optimization. *npj Quantum Inf*, 9(1):121, November 2023.
- [21] Christopher Jarzynski. Generating shortcuts to adiabaticity in quantum and classical dynamics. *Phys. Rev. A*, 88(4):040101, October 2013.
- [22] Michael Kolodrubetz, Dries Sels, Pankaj Mehta, and Anatoli Polkovnikov. Geometry and non-adiabatic response in quantum and classical systems. *Physics Reports*, 697:1–87, June 2017.
- [23] Yue Ruan, Pengyue Chen, Qi Li, Ling Yang, Zhiqiang Yuan, Xiling Xue, Xi Li, and Zhihao Liu. XY-mixer ansatz assisted by counterdiabatic driving for combinatorial optimization. *Phys. Rev. Research*, 7(1):013243, March 2025.
- [24] Patrick Reberstrost and Seth Lloyd. Quantum Computational Finance: Quantum Algorithm for Portfolio Optimization. *Künstl Intell*, August 2024.
- [25] Amira Abbas, Andris Ambainis, Brandon Augustino, Andreas Bäertschi, Harry Buhman, Carleton Coffrin, Giorgio Cortiana, Vedran Dunjko, Daniel J Egger, Bruce G Elmegreen, et al. Challenges and opportunities in quantum optimization. *Nature Reviews Physics*, pages 1–18, 2024.
- [26] Alexey Galda, Eesh Gupta, Jose Falla, Xiaoyuan Liu, Danylo Lykov, Yuri Alexeev, and Ilya Safro. Similarity-based parameter transferability in the quantum approximate optimization algorithm. *Frontiers in Quantum Science and Technology*, 2, 2023.
- [27] Jose Falla, Quinn Langfitt, Yuri Alexeev, and Ilya Safro. Graph representation learning for parameter transferability in quantum approximate optimization algorithm. *Quantum Machine Intelligence*, 6(2):46, 2024.
- [28] Kien X Nguyen, Bao Bach, and Ilya Safro. Cross-problem parameter transfer in quantum approximate optimization algorithm: A machine learning approach. In *2025 IEEE International Conference on Quantum Computing and Engineering (QCE)*, pages 1–10. IEEE, 2025.
- [29] Boris Tselikhovskiy, Bao Bach, Jose Falla, and Ilya Safro. Reductions of qaoa induced by classical symmetries: Theoretical insights and practical implications. *arXiv preprint arXiv:2602.16141*, 2026.
- [30] Boris Tselikhovskiy, Ilya Safro, and Yuri Alexeev. Equivariant qaoa and the duel of the mixers. *arXiv preprint arXiv:2405.07211*, 2024.
- [31] Boris Tselikhovskiy, Ilya Safro, and Yuri Alexeev. Symmetries and dimension reduction in quantum approximate optimization algorithm. *arXiv preprint arXiv:2309.13787*, 2026.
- [32] Ruslan Shaydulin, Stuart Hadfield, Tad Hogg, and Ilya Safro. Classical symmetries and the quantum approximate optimization algorithm. *Quantum Information Processing*, 20(11):1–28, 2021.
- [33] Reuben Tate, Jai Moondra, Bryan Gard, Greg Mohler, and Swati Gupta. Warm-started qaoa with custom mixers provably converges and computationally beats goemans-williamson’s max-cut at low circuit depths. *Quantum*, 7:1121, 2023.
- [34] Daniel J Egger, Jakub Mareček, and Stefan Woerner. Warm-starting quantum optimization. *Quantum*, 5:479, 2021.
- [35] Ankit Kulshrestha and Ilya Safro. Beinit: Avoiding barren plateaus in variational quantum algorithms. *2022 IEEE International Conference on Quantum Computing and Engineering (QCE)*, pages 197–203, 2022.
- [36] Filip B Maciejewski, Bao G Bach, Maxime Dupont, P Aaron Lott, Bhuvanesh Sundar, David E Bernal Neira, Ilya Safro, and Davide Venturelli. A multilevel approach for solving large-scale qubo problems with noisy hybrid quantum approximate optimization. In *2024 IEEE High Performance Extreme Computing Conference (HPEC)*, pages 1–10. IEEE, 2024.

- [37] Bao Bach, Jose Falla, and Ilya Safro. Mlqaoa: Graph learning accelerated hybrid quantum-classical multilevel qaoa. In *2024 IEEE International Conference on Quantum Computing and Engineering (QCE)*, volume 01, pages 1–12, 2024.
- [38] Stuart Hadfield, Zhihui Wang, Bryan O’Gorman, Eleanor G. Rieffel, Davide Venturelli, and Rupak Biswas. From the Quantum Approximate Optimization Algorithm to a Quantum Alternating Operator Ansatz. *Algorithms*, 12(2):34, February 2019. arXiv:1709.03489 [quant-ph].
- [39] Dries Sels and Anatoli Polkovnikov. Minimizing irreversible losses in quantum systems by local counterdiabatic driving. *Proc. Natl. Acad. Sci. U.S.A.*, 114(20), May 2017.
- [40] Yu Li, Guojing Tian, Xiaoyu He, and Xiaoming Sun. Preparation of Hamming-Weight-Preserving Quantum States with Log-Depth Quantum Circuits, August 2025. arXiv:2508.14470 [quant-ph].
- [41] Panagiotis KI Barkoutsos, Giacomo Nannicini, Anton Robert, Ivano Tavernelli, and Stefan Woerner. Improving Variational Quantum Optimization using CVaR. *Quantum*, 4:256, April 2020. arXiv:1907.04769 [quant-ph].
- [42] Franz Georg Fuchs. QAOA: A modular python library for the quantum approximate optimization algorithm, 2024. Version 2.0.0.
- [43] Andreas Bartschi and Stephan Eidenbenz. Grover Mixers for QAOA: Shifting Complexity from Mixer Design to State Preparation. In *2020 IEEE International Conference on Quantum Computing and Engineering (QCE)*, pages 72–82, Denver, CO, USA, October 2020. IEEE.
- [44] Steven Kordonowy and Hannes Leipold. The Lie Algebra of XY-mixer Topologies and Warm Starting QAOA for Constrained Optimization, September 2025. arXiv:2505.18396 [quant-ph].
- [45] David Bucher, Maximilian Janetschek, Michael Poppel, Jonas Stein, Claudia Linnhoff-Popien, and Sebastian Feld. Constrained Quantum Optimization via Iterative Warm-Start XY-Mixers, April 2026. arXiv:2604.02083 [quant-ph].
- [46] Linghua Zhu, Ho Lun Tang, George S. Barron, F. A. Calderon-Vargas, Nicholas J. Mayhall, Edwin Barnes, and Sophia E. Economou. Adaptive quantum approximate optimization algorithm for solving combinatorial problems on a quantum computer. *Phys. Rev. Research*, 4(3):033029, July 2022.
- [47] Hayato Ushijima-Mwesigwa, Ruslan Shaydulin, Christian FA Negre, Susan M Mniszewski, Yuri Alexeev, and Ilya Safro. Multilevel combinatorial optimization across quantum architectures. *ACM Transactions on Quantum Computing*, 2(1):1–29, 2021.
- [48] Linghua Zhu, Ho Lun Tang, George S Barron, FA Calderon-Vargas, Nicholas J Mayhall, Edwin Barnes, and Sophia E Economou. Adaptive quantum approximate optimization algorithm for solving combinatorial problems on a quantum computer. *Physical Review Research*, 4(3):033029, 2022.
- [49] Ilya Tyagin, Marwa H Farag, Kyle Sherbert, Karunya Shirali, Yuri Alexeev, and Ilya Safro. QAOA-GPT: Efficient generation of adaptive and regular quantum approximate optimization algorithm circuits. In *2025 IEEE International Conference on Quantum Computing and Engineering (QCE)*, volume 1, pages 1505–1515. IEEE, 2025.