

RBM 2hidden

thou

August 2018

1 Introduction

consider a RBM with 2 hidden unit h_1, h_2 , the input is σ , which the dimension is N , the inverse temperature is β , so the joint probability is

$$P(\sigma, h_1, h_2) = \frac{1}{Z} \exp \frac{\beta}{\sqrt{N}} (h_1 \sum_{i=1}^N \xi_i^1 \sigma_i + h_2 \sum_{i=1}^N \xi_i^2 \sigma_i) \quad (1)$$

sum over the hidden unit, we can get,

$$P(\sigma) = \frac{1}{Z} [2 \cosh \frac{\beta}{\sqrt{N}} \sum_{i=1}^N (\xi_i^1 + \xi_i^2) \sigma_i + 2 \cosh \frac{\beta}{\sqrt{N}} \sum_{i=1}^N (\xi_i^1 - \xi_i^2) \sigma_i] \quad (2)$$

so we can get the partition function as

$$Z = \sum_{\sigma_i} [2 \cosh \frac{\beta}{\sqrt{N}} \sum_{i=1}^N (\xi_i^1 + \xi_i^2) \sigma_i + 2 \cosh \frac{\beta}{\sqrt{N}} \sum_{i=1}^N (\xi_i^1 - \xi_i^2) \sigma_i] \quad (3)$$

the final results is

$$Z = 2 \prod_{i=1}^N [2 \cosh \frac{\beta}{\sqrt{N}} (\xi_i^1 + \xi_i^2)] + 2 \prod_{i=1}^N [2 \cosh \frac{\beta}{\sqrt{N}} (\xi_i^1 - \xi_i^2)] \quad (4)$$

by using the equation for small x

$$\cosh x = \exp(\frac{x^2}{2}) \quad (5)$$

we simplify the paration function as:

$$Z = 2^{N+1} e^{\beta^2} (2 \cosh \frac{\beta^2}{N} \sum_{i=1}^N \xi_i^1 \xi_i^2) \quad (6)$$

the message passing for this system is:

$$P_{i \rightarrow a}(\xi_i^1, \xi_i^2) \propto \prod_{b \in \partial i/a} \hat{P}_{b \rightarrow i}(\xi_i^1, \xi_i^2) \quad (7)$$

and

$$\hat{P}_{b \rightarrow i}(\xi_i^1, \xi_i^2) \propto Z^{-1} \cosh \beta X \cosh \beta Y \prod_{j \in \partial b/i} P_{j \rightarrow b}(\xi_j^1, \xi_j^2) \quad (8)$$

where

$$X = \frac{1}{\sqrt{N}} \sum_{i=1}^N \xi_i^1 \sigma_i^a \quad (9)$$

and

$$Y = \frac{1}{\sqrt{N}} \sum_{i=1}^N \xi_i^2 \sigma_i^a \quad (10)$$

by use

$$\omega_1 = \sum_{j \in \partial b/i} \frac{1}{\sqrt{N}} \xi_j^1 \sigma_j^b \quad (11)$$

$$\omega_2 = \sum_{j \in \partial b/i} \frac{1}{\sqrt{N}} \xi_j^2 \sigma_j^b \quad (12)$$

by relaxed BP,

$$m_{b \rightarrow i}^1 = \sum_{j \in \partial b/i} \frac{1}{\sqrt{N}} \sigma_j^n m_{j \rightarrow b}^1 \quad (13)$$

$$m_{b \rightarrow i}^2 = \sum_{j \in \partial b/i} \frac{1}{\sqrt{N}} \sigma_j^n m_{j \rightarrow b}^2 \quad (14)$$

and

$$\hat{\sigma}_{b \rightarrow i}^1 = 1 - \frac{1}{N} \sum_{j \in \partial b/i} (m_{j \rightarrow b}^1)^2 \quad (15)$$

$$\hat{\sigma}_{b \rightarrow i}^2 = 1 - \frac{1}{N} \sum_{j \in \partial b/i} (m_{j \rightarrow b}^2)^2 \quad (16)$$

and the correlation is :

$$\rho_{b \rightarrow i} = \frac{\frac{1}{N} \sum_{j \in \partial b/i} (Q_{j \rightarrow b} - m_{j \rightarrow b}^1 m_{j \rightarrow b}^2)}{\hat{\sigma}_{b \rightarrow i}^1 \hat{\sigma}_{b \rightarrow i}^2} \quad (17)$$

where

$$Q_{j \rightarrow b} = \frac{\prod_{a \in \partial j/b} \hat{P}_{a \rightarrow j}^{++} + \prod_{a \in \partial j/b} \hat{P}_{a \rightarrow j}^{--} - \prod_{a \in \partial j/b} \hat{P}_{a \rightarrow j}^{+-} - \prod_{a \in \partial j/b} \hat{P}_{a \rightarrow j}^{-+}}{\prod_{a \in \partial j/b} \hat{P}_{a \rightarrow j}^{++} + \prod_{a \in \partial j/b} \hat{P}_{a \rightarrow j}^{--} + \prod_{a \in \partial j/b} \hat{P}_{a \rightarrow j}^{+-} + \prod_{a \in \partial j/b} \hat{P}_{a \rightarrow j}^{-+}} \quad (18)$$

we can get

$$\hat{P}_{b \rightarrow i} = \cosh^{-1} \left(\frac{\beta^2}{N} \sum_j Q_j \right) e^{\frac{1}{2} \beta^2 (\hat{\sigma}_{b \rightarrow i}^1 + \hat{\sigma}_{b \rightarrow i}^2)} [e^{\beta^2 \rho_{b \rightarrow i} \hat{\sigma}_{b \rightarrow i}^1 \hat{\sigma}_{b \rightarrow i}^2} \cosh \beta X + e^{-\beta^2 \rho_{b \rightarrow i} \hat{\sigma}_{b \rightarrow i}^1 \hat{\sigma}_{b \rightarrow i}^2} \cosh \beta Y] \quad (19)$$

where

$$X = \frac{1}{\sqrt{N}} (\xi_i^1 + \xi_i^2) \sigma_i^b + m_{b \rightarrow i}^1 + m_{b \rightarrow i}^2 \quad (20)$$

$$Y = \frac{1}{\sqrt{N}} (\xi_i^1 - \xi_i^2) \sigma_i^b + m_{b \rightarrow i}^1 - m_{b \rightarrow i}^2 \quad (21)$$

and

$$Q_j = \frac{\prod_{a \in \partial j} \hat{P}_{a \rightarrow j}^{++} + \prod_{a \in \partial j} \hat{P}_{a \rightarrow j}^{--} - \prod_{a \in \partial j} \hat{P}_{a \rightarrow j}^{+-} - \prod_{a \in \partial j/b} \hat{P}_{a \rightarrow j}^{-+}}{\prod_{a \in \partial j} \hat{P}_{a \rightarrow j}^{++} + \prod_{a \in \partial j} \hat{P}_{a \rightarrow j}^{--} + \prod_{a \in \partial j} \hat{P}_{a \rightarrow j}^{+-} + \prod_{a \in \partial j} \hat{P}_{a \rightarrow j}^{-+}} \quad (22)$$

$$m_{j \rightarrow b}^1 = \frac{\prod_{a \in \partial j/b} \hat{P}_{a \rightarrow j}^{++} + \prod_{a \in \partial j/b} \hat{P}_{a \rightarrow j}^{+-} - \prod_{a \in \partial j/b} \hat{P}_{a \rightarrow j}^{-+} - \prod_{a \in \partial j/b} \hat{P}_{a \rightarrow j}^{--}}{\prod_{a \in \partial j/b} \hat{P}_{a \rightarrow j}^{++} + \prod_{a \in \partial j/b} \hat{P}_{a \rightarrow j}^{--} + \prod_{a \in \partial j/b} \hat{P}_{a \rightarrow j}^{+-} + \prod_{a \in \partial j/b} \hat{P}_{a \rightarrow j}^{-+}} \quad (23)$$

$$m_{j \rightarrow b}^2 = \frac{\prod_{a \in \partial j/b} \hat{P}_{a \rightarrow j}^{++} + \prod_{a \in \partial j/b} \hat{P}_{a \rightarrow j}^{+-} - \prod_{a \in \partial j/b} \hat{P}_{a \rightarrow j}^{--} - \prod_{a \in \partial j/b} \hat{P}_{a \rightarrow j}^{-+}}{\prod_{a \in \partial j/b} \hat{P}_{a \rightarrow j}^{++} + \prod_{a \in \partial j/b} \hat{P}_{a \rightarrow j}^{--} + \prod_{a \in \partial j/b} \hat{P}_{a \rightarrow j}^{+-} + \prod_{a \in \partial j/b} \hat{P}_{a \rightarrow j}^{-+}} \quad (24)$$