

GOTTA Prototype: Known-Asteroid Extraction and Statistical Performance Evaluation

Yun-Ao Xiao^{1,2}, Hu Zou¹, Feiyang Tian^{3,4}, Zhaoxing Liu^{3,4}, Jian Gao^{3,4}, Shuai Feng⁵,
Bo Zhang⁵, Yuyi Zhuang^{3,4} and The GOTTA Collaboration

¹ National Astronomical Observatories, Chinese Academy of Sciences, Beijing 100101, China;
xiaoyunao@nao.cas.cn

² School of Astronomy and Space Science, University of Chinese Academy of Sciences, Beijing
101408, China

³ Department of Astronomy, Beijing Normal University, Beijing 100875, China

⁴ Institute of Frontier Sciences in Astronomy and Astrophysics, Beijing Normal University,
Beijing 100875, China

⁵ Shandong Key Laboratory of Space Environment and Exploration Technology, School of Space
Science and Technology, Institute of Space Sciences, Shandong University, Weihai, Shandong,
264209, China

Received 20xx month day; accepted 20xx month day

Abstract The GOTTA Prototype is a wide-field, high-cadence pathfinder for the full Global Open Transient Telescope Array. Its repeated survey imaging naturally records large numbers of Solar System small bodies, providing an opportunity to validate moving-object processing before the full array is deployed. We present a known-object association workflow for recovering cataloged asteroids from GOTTA Prototype photometric catalogs. For each exposure, the workflow determines the CCD footprint and mid-exposure epoch, predicts known-object ephemerides, filters candidates within the detector footprint, associates predicted positions with calibrated catalog detections, and augments the accepted matches with orbital and observing-geometry information from external small-body databases. The resulting recovered sample contains 56,230 detections of 16,053 unique known asteroids over 73 UTC nights, including 77 near-Earth objects and 19 potentially hazardous asteroids. The sample is dominated by main-belt asteroids. The observed-minus-predicted positional residual has a median separation of 0.220 arcsec and a two-dimensional RMS of 0.355 arcsec. We characterize the recovered sample in sky position, orbital class, aperture photometry, astrometric residual, angular rate, observing geometry, and repeated temporal sampling. These results demonstrate that the GOTTA Prototype has established a working known-object recovery layer and provide a practical baseline for

future moving-object association, NEO follow-up support, and asteroid light-curve studies with the full GOTTA array.

Key words: surveys — minor planets, asteroids: general — techniques: photometric — techniques: astrometric — methods: data analysis

1 INTRODUCTION

Near-Earth objects (NEOs) and potentially hazardous asteroids (PHAs) have long motivated systematic searches for small bodies in the Solar System. The Spaceguard Survey emphasized that the first step toward mitigating impact hazards is a comprehensive search for Earth-crossing asteroids and comets, followed by reliable orbit determination and long-term tracking (Morrison 1992). During the past three decades, wide-field CCD surveys have transformed this task from targeted searches into survey-scale operations. Programs such as Spacewatch, LINEAR, the Catalina Sky Survey, Pan-STARRS, and ATLAS have greatly increased the inventory of known minor planets and have established the basic observing and processing framework for modern moving-object surveys (e.g. Stokes et al. 2000; Larsen 2007; Larson et al. 2003; Denneau et al. 2013; Tonry et al. 2018).

Moving-object science in time-domain surveys depends not only on single-image depth, but also on cadence, sky coverage, astrometric precision, and the ability to associate detections across time. A discovery-oriented pipeline generally starts from transient or non-stationary detections, forms same-night tracklets, and links tracklets obtained on different nights to determine preliminary orbits (Kubica et al. 2007; Denneau et al. 2013; Vereš & Chesley 2017). The same data stream also supports attribution and precovery of known objects, which are essential for orbit refinement, survey validation, and physical characterization. This distinction is particularly important for prototype surveys: even before a full discovery pipeline is finalized, the recovery of cataloged asteroids can be used to evaluate astrometric performance, photometric behavior, source extraction quality, and temporal sampling under real observing conditions.

Several recent and forthcoming surveys illustrate the range of strategies relevant to Solar System time-domain work. ATLAS was designed as a high-cadence all-sky system optimized for the discovery of dangerous near-Earth asteroids and other rapidly varying sources (Tonry et al. 2018). ZTF combines a 47 deg² camera with near-real-time reduction and alert distribution for transients, variables, and moving objects (Bellm et al. 2019; Patterson et al. 2019). The Rubin Observatory Legacy Survey of Space and Time will provide much deeper multi-epoch imaging and is expected to substantially expand the number of Solar System detections (Ivezić et al. 2019; Jones et al. 2018; Schwamb et al. 2023). In parallel, the NEO Surveyor mission will search in the infrared from space and is optimized for discovering and characterizing NEOs, especially objects difficult to observe from the ground (Mainzer et al. 2023). These facilities show that cadence, depth, wavelength, and sky coverage are complementary rather than interchangeable survey properties.

High-cadence survey data are also valuable for asteroid physical characterization. Time-series photometry constrains rotation periods and amplitudes, and multi-geometry light curves can be used to infer spin states and shapes through light-curve inversion (Kaasalainen & Torppa 2001; Kaasalainen et al. 2001; Warner et al. 2009). Sparse survey photometry further enlarges the statistical sample of asteroid light curves when sufficient temporal and phase-angle coverage is available. Accurate photometry and observing geometry are therefore necessary not only for detection and orbit association, but also for studying asteroid rotation, shape, surface properties, and activity. Phase-angle effects must also be treated explicitly when survey photometry is interpreted physically, for example through the H, G_1, G_2 phase-function formalism (Muinonen et al. 2010).

The SiTian/GOTTA program is designed to provide high-cadence, wide-field optical monitoring with a distributed array of telescopes (Liu et al. 2021). Its pathfinder, the Mini-SiTian Array at Xinglong Observatory, has already demonstrated the scientific value of high-cadence, multi-band observations for time-domain astronomy. The Mini-SiTian Array consists of three 30 cm telescopes, has operated at Xinglong since 2022, and provides large-field photometric monitoring that informs both observation strategy and data processing for the full SiTian/GOTTA system (Han et al. 2025). These pathfinder data and procedures are directly relevant to the development of Solar System analysis tools for future high-cadence survey operations.

For asteroid science specifically, Liu et al. (2025) developed an asteroid photometry and period-analysis pipeline using Mini-SiTian data from the f02 region. They derived rotation periods for 22 asteroids, including 14 objects without periods listed in ALCDEF, and found agreement with existing results for several objects with sufficiently dense photometric sampling. This work demonstrates the feasibility of asteroid photometry with SiTian pathfinder observations and provides an important motivation for building the corresponding known-object association and validation layer for the GOTTA Prototype.

The GOTTA Prototype extends this pathfinder context to a larger-aperture wide-field system. Mini-SiTian has shown that small pathfinder facilities can deliver useful asteroid light curves, while the GOTTA Prototype provides a testbed for survey-scale source catalogs, exposure geometry, and small-body database augmentation. The present work does not attempt to report unknown asteroid discoveries, measure formal completeness, or operate a full moving-object linking system. Instead, it focuses on known-object recovery, a necessary precursor to tracklet construction, moving-object alert filtering, orbit refinement, and light-curve preparation.

In this paper we present a known-asteroid extraction pipeline for GOTTA Prototype photometric catalogs. We use cataloged minor planets as a controlled sample to validate the end-to-end association between survey catalogs and known-object ephemerides. This approach allows us to evaluate astrometric residuals, photometric measurement behavior, angular-rate coverage, orbital-class distribution, and repeated-observation statistics. The recovered sample contains 56,230 matched detections of 16,053 unique known asteroids, including a small but useful subset of NEOs and PHAs.

The paper is organized as follows. Section 2 describes the GOTTA Prototype data and the recovered known-asteroid sample. Section 3 presents the ephemeris-based extraction and augmentation method. Section 4 analyzes the recovered sample and evaluates the association performance. Section 5 discusses the implications for the future GOTTA moving-object pipeline and summarizes the main conclusions.

2 OBSERVATIONS AND CATALOG DATA

2.1 GOTTA Prototype Survey Data

The observations analyzed in this work were obtained with the GOTTA Prototype during the original four-CMOS test campaign from 2025 January to June. The prototype is a 1 m-class wide-field Ring Schmidt system located at Xinglong Observatory and was operated during this stage with a single SDSS-like g filter. The optical design, detector layout, and observing configuration are described in detail by Zhengyang Li et al. (in preparation); here we only summarize the aspects required for known-asteroid recovery.

The present analysis uses calibrated source catalogs produced by the GOTTA Prototype basic data-reduction pipeline. That pipeline performs calibration-frame construction, image preprocessing, astrometric calibration, photometric calibration, and source extraction, as described by Niu Li et al. (in preparation). The astrometric solution is tied to Gaia DR3 (Gaia Collaboration et al. 2023), and the photometric catalog provides source positions, aperture photometry, Kron-like photometry, photometric uncertainties, and quality flags for each exposure. Source extraction and astrometric calibration use established wide-field catalog-processing tools in the same general class as SExtractor, SCAMP, and astrometry.net (Bertin & Arnouts 1996; Bertin 2006; Lang et al. 2010); related CMOS wide-field reduction strategies have also been developed for Mini-SiTian (Xiao et al. 2025). For known-asteroid recovery, the most important quantities are the WCS, the exposure time and observing epoch, the measured source coordinates, and the aperture-photometry measurements.

The companion reduction analysis shows that the calibrated catalogs are sufficiently accurate for sub-arcsecond source association. Stellar astrometric residuals are much smaller than the matching scale used here, while the photometric catalog includes aperture-corrected measurements in a sequence of circular apertures. Asteroid detections are generally fainter than the stars used for calibration and may be affected by motion during the exposure. Their observed-minus-predicted residuals therefore include additional contributions from centroiding uncertainty, ephemeris uncertainty, timing uncertainty, and possible trailing.

2.2 Recovered Known-Asteroid Sample

The recovered sample consists of accepted associations between predicted positions of cataloged asteroids and calibrated GOTTA source detections. Each association records the measured source position and photometry, the predicted ephemeris, the observing epoch, and auxiliary orbital and observing-geometry quantities. The final sample contains 56,230 detections of 16,053 unique known




Fig. 1: Representative known-asteroid detections and image cutouts from the pilot-survey material. Green labels and circles identify recovered known asteroids. The figure illustrates the type of image-level detections represented by the accepted associations analyzed in this work.

asteroids over 73 UTC nights. It is dominated by main-belt asteroids, with smaller contributions from Mars-crossing asteroids, Jupiter Trojans, NEOs, and PHAs.

Figure 1 shows representative image cutouts of recovered asteroids. These examples illustrate the type of source detections included in the statistical analysis; all quantitative results below are derived from the full recovered sample.

3 KNOWN-ASTEROID EXTRACTION METHOD

3.1 Known-Object Association Strategy

The goal of the pipeline is to identify detections of cataloged asteroids in calibrated GOTTA Prototype source catalogs. The procedure is designed as a known-object association workflow rather than a discovery-oriented linking pipeline. For each exposure, the pipeline reads the calibrated photometric catalog and the corresponding FITS header. The WCS solution is used to determine the sky footprint of the detector, while the exposure midpoint is computed from the start time and exposure duration. Known-asteroid candidates are predicted for this epoch and for the Xinglong observer location. Objects outside the padded field region, outside the CCD footprint, or fainter than the adopted ephemeris-magnitude limit are removed before cross-matching. The remaining predictions are associated with catalog detections using an angular nearest-neighbor criterion. Accepted associations are then merged with the source photometry and augmented with orbit and observing-geometry quantities from external small-body databases.

Figure 2 summarizes the workflow. The present analysis uses only accepted associations. It is therefore appropriate for evaluating the quality and content of the recovered known-object sample, but not for measuring a formal detection efficiency, which would require the full set of predicted but unrecovered objects.

3.2 Ephemeris Prediction and Exposure Geometry

For a cataloged asteroid, the starting point of the ephemeris calculation is its osculating orbit at a reference epoch. The orbit can be expressed by the standard elements $(a, e, i, \Omega, \omega, M)$, where a is the semi-major axis, e is the eccentricity, i is the inclination, Ω is the longitude of the ascending




Fig. 2: Workflow of the known-asteroid extraction procedure. The pipeline starts from calibrated per-exposure photometric catalogs and exposure metadata, computes the exposure geometry and observing epoch, predicts known-object positions, applies field and footprint filters, matches surviving predictions to catalog detections, and augments accepted associations with external small-body information.

node, ω is the argument of perihelion, and M is the mean anomaly. In the two-body limit, the mean anomaly evolves as

$$M(t) = M_0 + n(t - t_0), \quad (1)$$

where n is the mean motion. Kepler's equation,

$$M = E - e \sin E, \quad (2)$$

is solved for the eccentric anomaly E , from which the heliocentric position is obtained. For survey applications, the apparent topocentric direction also depends on the observer location, time scale,

light-time correction, and perturbations from Solar System bodies. The pipeline therefore uses a fast ephemeris-screening step to identify plausible in-field candidates and then augments the accepted matches with refined information from the JPL Small-Body Database¹ and Horizons² systems. The Minor Planet Center³ provides the broader astrometric-observation infrastructure and reporting context for Solar System small-body work. The Lowell Observatory astorb database⁴ is used as an orbit-catalog resource for fast screening.

The exposure footprint is computed directly from the WCS of each calibrated catalog. The detector corners are projected from pixel coordinates to the sky, and the search radius is initialized from the maximum angular distance between the field center and the projected corners, with an additional padding term to avoid edge losses. The exposure midpoint is

$$t_{\text{mid}} = t_{\text{start}} + \frac{1}{2}t_{\text{exp}}. \quad (3)$$

This midpoint is used for ephemeris evaluation because even modest sky-plane motion can lead to measurable offsets if the start time is used instead. After the initial sky-coordinate query, predicted positions are transformed back to detector coordinates, and only candidates falling within the valid pixel range are retained.

3.3 Catalog Matching and Small-Body Augmentation

The filtered predictions are associated with the GOTTA source catalog using their angular separation from detected sources. The residual between a measured source position and a predicted ephemeris is defined as

$$\Delta\theta = [(\Delta\alpha \cos \delta)^2 + (\Delta\delta)^2]^{1/2}, \quad (4)$$

where $\Delta\alpha$ and $\Delta\delta$ are the right ascension and declination differences. A match is accepted only when the nearest-neighbor separation is below the adopted arcsecond-level threshold. The threshold is chosen to be larger than the expected astrometric and centroiding uncertainties for faint moving sources, but sufficiently small to reduce chance associations in crowded fields.

For each accepted association, the measured catalog quantities are merged with the predicted ephemeris. The photometric measurements include aperture magnitudes and their uncertainties; Kron-like measurements are retained for comparison but are not used as the primary photometric statistic in the final analysis. The predicted asteroid magnitude is used only as a consistency diagnostic, because it is not necessarily defined in the same photometric system as the GOTTA *g*-band measurement and is also affected by asteroid color, phase behavior, and rotational variability.

The matched detections are further associated with small-body database entries to obtain object identifiers, orbit classes, orbital elements, and observer-dependent geometry. The latter includes heliocentric distance, topocentric distance, phase angle, and sky-plane angular rates. These quantities are used in the statistical analysis below to interpret the recovered sample in terms of orbital population, observing geometry, and apparent motion.

¹ https://ssd.jpl.nasa.gov/tools/sbdb_lookup.html

² <https://ssd.jpl.nasa.gov/horizons/>

³ <https://minorplanetcenter.net/>

⁴ <https://asteroid.lowell.edu/main/astorb/>

Table 1: Statistics for the recovered GOTTA known-asteroid sample.

Quantity	Value	Note
Accepted asteroid-source associations	56,230	Accepted source-ephemeris matches
Distinct known asteroids	16,053	Unique minor-planet identifiers
Exposure catalogs with accepted associations	13,253	Per-exposure products
UTC nights represented	73	Recovered sample date coverage
Field groups represented	588	Grouped by pointing metadata
NEO-class objects	77	Identified from small-body metadata
PHA-class objects	19	Identified from small-body metadata
Median ephemeris magnitude	18.2215	Predicted brightness
Median g_{aper}	18.1110	Adopted aperture magnitude
Median $\sigma(g_{\text{aper}})$	0.1239 mag	Adopted aperture uncertainty
Median aperture S/N proxy	8.7621	Flux/error in adopted aperture
Median observed-minus-predicted separation	0.220 arcsec	Accepted associations
84th-percentile observed-minus-predicted separation	0.484 arcsec	Accepted associations
2D residual RMS	0.355 arcsec	$\sqrt{\text{mean}(\text{sep}^2)}$
RMS in $\Delta\alpha \cos \delta$	0.250 arcsec	Coordinate residual
RMS in $\Delta\delta$	0.252 arcsec	Coordinate residual
Median angular rate	29.276 arcsec hr ⁻¹	Derived from augmented geometry
Median phase angle	13.245 deg	Derived from augmented geometry
Associations without geometry augmentation	107	Excluded from geometry statistics

The statistical calculations and figure generation use the standard Python scientific stack and Astropy for astronomical table and coordinate handling ([Astropy Collaboration et al. 2013, 2018, 2022](#)). These software citations are included for reproducibility of the analysis products.

4 RESULTS

4.1 Recovered Sample and Orbital Populations

The recovery procedure yields 56,230 accepted associations corresponding to 16,053 distinct known asteroids. These detections span 73 UTC nights during the 2025 January–June prototype campaign. The size of this sample is sufficient to characterize the dominant orbital populations recovered by the survey and to examine how the association residuals depend on brightness and apparent motion. Table 1 summarizes the main statistics.

The orbit-class composition is shown in Table 2. Main-belt asteroids account for 51,058 detections and 14,674 unique objects. Outer main-belt asteroids, Jupiter Trojans, inner main-belt asteroids, and Mars-crossing asteroids make up most of the remaining sample. The NEO classes are represented by Amor, Apollo, and Aten objects. The dominance of main-belt objects is expected because the survey footprint and limiting depth preferentially sample objects near ordinary

Table 2: Orbit-class composition of the recovered known-asteroid sample. Separations are in arcsec.

Code	Orbit class	Detections	Unique objects	Median g_{aper}	Median separation
MBA	Main-belt Asteroid	51,058	14,674	18.123	0.221
OMB	Outer Main-belt Asteroid	1,970	522	17.893	0.187
TJN	Jupiter Trojan	1,362	317	18.005	0.207
IMB	Inner Main-belt Asteroid	897	290	18.315	0.257
MCA	Mars-crossing Asteroid	656	166	17.847	0.228
AMO	Amor	139	31	17.143	0.227
APO	Apollo	101	38	17.749	0.253
TNO	TransNeptunian Object	20	3	17.588	0.243
ATE	Aten	19	8	17.668	0.292
CEN	Centaur	5	3	18.481	0.237
AST	Asteroid	3	1	17.074	0.199




Fig. 3: Left: detection-level sky distribution of recovered known asteroids in equatorial coordinates, binned with HEALPix $n_{\text{side}} 64$. The color scale is clipped at 100 detections per pixel for readability. Right: orbital-element distribution of the recovered known-object sample.

main-belt opposition geometries. The smaller NEO and PHA subsets should therefore be viewed as validation samples for the recovery path rather than as indicators of discovery performance.

Figure 3 shows the detection-level sky distribution and the orbital-element distribution of the recovered objects. The sky distribution is shaped jointly by the prototype survey footprint, seasonal visibility, and the instantaneous distribution of known minor planets during the observing campaign. The orbital-element panel confirms that the recovered sample is dominated by main-belt objects, with smaller contributions from inner and outer main-belt objects, Jupiter Trojans, Mars-crossers, and NEOs.

4.2 Photometric Properties of the Recovered Asteroids

For the photometric analysis we use the aperture magnitude corresponding to the optimal aperture defined by the basic GOTTA photometric pipeline. In that pipeline, aperture photometry is measured in a series of circular apertures, and the optimal aperture is determined from the signal-to-noise ratio of stellar sources as a function of aperture radius. A larger reference aperture is used for aperture correction, so that the cataloged aperture magnitudes are placed on a consistent photometric scale. We therefore do not re-derive an asteroid-specific optimal aperture in this

work. Instead, we use the pipeline-defined optimal aperture as a homogeneous measurement for the recovered asteroid sample.

This choice is preferable to Kron-like photometry for the statistical analysis presented here. Kron apertures are adaptive and can respond differently to background noise, faint-source segmentation, and possible source elongation, whereas the adopted aperture magnitude provides a uniform measurement across the sample. Kron photometry is retained in the catalog for comparison, but it is not used as the primary quantity in the magnitude, uncertainty, and astrometric-residual diagnostics below.

For the current catalog, the optimal aperture corresponds to g_{aper} below. Figure 4 shows the photometric diagnostics of the recovered sample. Most recovered asteroids lie close to the faint end of routine single-exposure catalog measurements rather than in the high signal-to-noise regime used for stellar calibration. The uncertainty–magnitude relation increases toward the faint end, as expected from decreasing source signal-to-noise ratio. The difference between predicted ephemeris magnitude and measured GOTTA g -band aperture magnitude is used only as a consistency diagnostic, because the two quantities are not necessarily defined in the same photometric system and may also differ because of asteroid color, phase behavior, rotation, shape, and ephemeris-magnitude uncertainties.

4.3 Astrometric Residuals and Apparent Motion

The observed-minus-predicted positional residual has a median separation of 0.220 arcsec and a two-dimensional RMS of 0.355 arcsec. The RMS values of $\Delta\alpha \cos \delta$ and $\Delta\delta$ are 0.250 arcsec and 0.252 arcsec, respectively. The coordinate residuals are centered close to zero, indicating that the accepted associations do not show a large global offset between the predicted ephemerides and the calibrated source positions.

Figure 5 shows the residual distribution and its dependence on the adopted aperture magnitude. The residual generally increases toward fainter magnitudes, consistent with centroiding uncertainty and decreasing signal-to-noise ratio. Table 3 summarizes the same behavior in magnitude bins: the median separation increases from about 0.09–0.11 arcsec for relatively bright objects to about 0.27 arcsec near $g_{\text{aper}} \simeq 18$ –19 mag.

The dependence on sky-plane angular rate is shown in Figure 6 and summarized in Table 4. Most recovered objects have moderate angular rates, with a median of 29.276 arcsec hr^{−1}. Over this dominant part of the sample, the residual distribution is similar to that of the full recovered catalog. The sparsely populated high-rate tail shows larger scatter and should be interpreted with caution because it contains few detections and may be more affected by trailing, centroiding biases, or matching ambiguity.

4.4 Temporal Sampling and Light-Curve Potential

The median heliocentric distance is 2.581 AU, and the median topocentric distance is 1.840 AU. The median phase angle is 13.245 deg. These values are consistent with a sample dominated by




Fig. 4: Aperture-photometry diagnostics for recovered known asteroids. The adopted aperture follows the optimal aperture defined by the GOTTA basic photometric pipeline. The panels show the adopted aperture magnitude distribution, the corresponding magnitude-uncertainty distribution, the aperture S/N proxy, and the dependence of magnitude uncertainty on adopted aperture magnitude. The last panel is shown as a density map to emphasize the dominant distribution of faint asteroid detections.

main-belt asteroids observed in typical survey geometry. Figure 7 shows the angular-rate, phase-angle, heliocentric-distance, and topocentric-distance distributions for rows with successful geometry augmentation.

Table 3: Astrometric residuals as a function of adopted aperture magnitude g_{aper} . Separations are in arcsec.

Magnitude bin	Detections	Median separation	RMS separation	Median mag. error
[12, 13)	340	0.113	0.210	0.005
[13, 14)	698	0.086	0.174	0.007
[14, 15)	1,072	0.089	0.194	0.010
[15, 16)	2,129	0.104	0.225	0.018
[16, 17)	5,447	0.135	0.259	0.039
[17, 18)	15,625	0.204	0.343	0.085
[18, 19)	24,990	0.270	0.391	0.155
[19, 20)	5,687	0.268	0.380	0.234




Fig. 5: Astrometric residual diagnostics. The panels show the observed-minus-predicted separation, $\Delta\alpha \cos \delta$, $\Delta\delta$, and separation as a function of the adopted aperture magnitude g_{aper} .




Fig. 6: Observed-minus-predicted separation as a function of sky-plane angular rate. The gray density map shows the dominant range of the recovered sample, while the red curve and shaded region show the running median and the 16th–84th percentile range in logarithmic rate bins. Most detections lie at moderate angular rates, where the residual distribution remains close to the full-sample behavior. A small number of higher-rate detections are excluded from the displayed range for clarity but are included in the binned statistics.

Table 4: Astrometric residuals as a function of apparent angular rate. Rate bins are in arcsec hr^{-1} and separations are in arcsec.

Rate bin	Detections	Median separation	RMS separation	Median g_{aper}
[0, 10)	4,917	0.213	0.356	18.053
[10, 20)	10,841	0.218	0.351	18.083
[20, 40)	31,792	0.222	0.357	18.136
[40, 80)	8,262	0.219	0.348	18.099
[80, 160)	224	0.222	0.337	17.790
[160, 320)	64	0.253	0.358	17.699
[320, 1000)	20	0.467	0.543	17.389




Fig. 7: Observing-geometry diagnostics for recovered known asteroids with successful geometry augmentation. The panels show apparent angular rate, phase angle, heliocentric distance, and topocentric distance. Vertical dashed lines mark the corresponding medians.

Repeated observations are essential for future light-curve analysis. In the present recovered sample, the median number of detections per object is two, while the mean is 3.50 and the 84th percentile is six. The most frequently recovered object has 317 detections over nine distinct nights, while only a small fraction of objects have more than several tens of detections. These highly repeated objects are useful for checking time-domain consistency, but they are not representative of the full recovered sample. The temporal baseline distribution indicates that many objects




Fig. 8: Temporal sampling diagnostics. The panels show nightly accepted known-asteroid associations, the detection-count distribution per object, the cumulative distribution of detections per object, and the cumulative distribution of object time baseline.

Table 5: Five UTC nights with the largest numbers of accepted known-asteroid associations.

UTC date	Detections	Unique objects	Exposure catalogs	Median g_{aper}	Median separation
2025-02-05	3,385	3,362	353	18.739	0.187
2025-01-19	3,119	3,065	576	18.354	0.189
2025-03-05	2,657	2,639	372	18.439	0.220
2025-01-28	2,656	2,638	395	18.499	0.189
2025-01-29	2,645	2,621	307	18.498	0.201

are sampled over multiple nights, but the prototype-only sample is still sparse for robust period determination for most objects.

Figure 8 shows nightly matched detections, the detection-count distribution per object, the cumulative distribution of detections per object, and the cumulative distribution of object time baseline. These diagnostics quantify not only how many objects were recovered, but also how often they were revisited. Table 5 lists the five UTC nights with the largest numbers of accepted associations. Thus, the prototype sample is useful for testing the construction of sparse asteroid time series, while dense rotational light-curve recovery requires either repeated visits to selected fields or auxiliary observations in future work.




Fig. 9: Example cross-matching diagnostic for one exposure. The panels show the detector distribution of accepted known-asteroid associations, residual vectors between predicted and measured positions, the separation distribution, and the relation between predicted ephemeris magnitude and measured aperture magnitude. Only accepted associations are shown; candidate-level rejection diagnostics require additional per-exposure products and are left for future development.

Figure 9 shows an example cross-matching diagnostic for one exposure. The panels show the detector distribution of accepted known-asteroid associations, residual vectors between predicted and measured positions, the separation distribution, and the relation between predicted ephemeris magnitude and measured aperture magnitude. Only accepted associations are shown; candidate-level rejection diagnostics require additional per-exposure products and are left for future development.

5 DISCUSSION AND CONCLUSIONS

The recovered known-asteroid sample demonstrates that calibrated GOTTA Prototype catalogs can already support systematic association with cataloged Solar System objects. The main validation provided by this work is the closure of the known-object association workflow: calibrated source catalogs are combined with exposure geometry and mid-exposure timing, known-object ephemerides are filtered by the real CCD footprint, accepted detections are matched in the source catalog, and the resulting associations are augmented with small-body database information. This workflow is a prerequisite for future moving-object products, including candidate-level bookkeeping, tracklet construction, and moving-object-aware transient filtering.

Known-object recovery is a useful validation mode because the predicted positions of cataloged asteroids are obtained independently of the GOTTA source extraction. The observed-minus-predicted residuals therefore test the combined effects of WCS quality, timing accuracy, centroiding, ephemeris uncertainty, and matching behavior. The median separation of 0.220 arcsec and the two-dimensional RMS of 0.355 arcsec show that the accepted associations are well localized on the scale required for known-object recovery. The increase of residual with fainter aperture magnitude indicates that centroiding precision and signal-to-noise ratio dominate the residual distribution for most of the sample. The weak dependence on angular rate over the main rate range suggests that trailing is not the primary limitation for the bulk of recovered objects, although the sparsely populated high-rate tail still requires dedicated analysis.

The photometric results should be interpreted as survey-catalog diagnostics rather than asteroid physical measurements. The adopted aperture magnitude provides a homogeneous brightness estimate, but the ephemeris magnitude is not a calibrated reference in the GOTTA g band. Asteroid color, rotational modulation, phase-angle dependence, shape, and uncertainties in the magnitude model can all contribute to the difference between predicted and observed brightness. Physical interpretation of the photometry will require phase-function modeling and, for selected targets, light-curve analysis over sufficient temporal and viewing-geometry coverage.

Several limitations should be kept in mind when interpreting the present recovered sample. Because the catalog contains only accepted associations, it does not provide the denominator required for a formal recovery-efficiency estimate. It also cannot quantify the false-association rate without the rejected candidates or the full per-exposure source catalogs. Future processing should therefore preserve the complete candidate list and should include random-shift, time-scrambling, or injection-recovery tests. A quantitative trailing analysis will also require exposure-level seeing and exposure-time information, so that the expected trail length,

$$L_{\text{trail}} = \mu t_{\text{exp}}, \quad (5)$$

can be compared with astrometric residuals and photometric uncertainties. Here μ is the apparent angular rate and t_{exp} is the exposure time.

For the future GOTTA array, the strength of the system will lie in the combination of large sky coverage and frequent temporal sampling. The prototype results show that the association and augmentation framework needed for known-object recovery can be implemented on GOTTA data. With candidate-level bookkeeping, repeated field visits, multi-band observations, and moving-object-aware scheduling, the same framework can be extended toward NEO monitoring, asteroid light-curve studies, and eventually discovery-oriented moving-object processing.

In summary, we have developed and tested a known-asteroid recovery workflow for GOTTA Prototype photometric catalogs. The recovered sample contains 56,230 detections of 16,053 unique known asteroids over 73 UTC nights, including small subsets of NEOs and PHAs. The sample is dominated by main-belt asteroids and provides statistically useful distributions in sky position, orbital class, aperture magnitude, angular rate, observing geometry, astrometric residual, and repeated temporal sampling. The sub-arcsecond observed-minus-predicted residuals demonstrate

that the current catalog products are suitable for known-object association. This work establishes the first moving-object validation layer for the GOTTA Prototype and provides the basis for future candidate-level detection, tracklet construction, and asteroid light-curve analysis.

Acknowledgements The authors acknowledge support from the Strategic Priority Research Program of the Chinese Academy of Sciences with Grant Nos. XDB0550103, the National Key R&D Program of China, the National Natural Science Foundation of China, and China Manned Space Project. The GOTTA Prototype Telescope is the pioneering unit of the global network dedicated to discovering and studying astronomical transients. The development and operation of the telescope are supported by the National Astronomical Observatories, Chinese Academy of Sciences and the University of Chinese Academy of Sciences.

References

- Astropy Collaboration, Robitaille, T. P., Tollerud, E. J., et al. 2013, *A&A*, 558, A33
- Astropy Collaboration, Price-Whelan, A. M., Sipocz, B. M., et al. 2018, *AJ*, 156, 123
- Astropy Collaboration, Price-Whelan, A. M., Lim, P. L., et al. 2022, *ApJ*, 935, 167
- Bellm, E. C., Kulkarni, S. R., Graham, M. J., et al. 2019, *PASP*, 131, 018002
- Bertin, E., & Arnouts, S. 1996, *A&AS*, 117, 393
- Bertin, E. 2006, in *ASP Conf. Ser.*, Vol. 351, *Astronomical Data Analysis Software and Systems XV*, ed. C. Gabriel, C. Arviset, D. Ponz, & S. Enrique, 112
- Denneau, L., Jedicke, R., Grav, T., et al. 2013, *PASP*, 125, 357
- Gaia Collaboration, Vallenari, A., Brown, A. G. A., et al. 2023, *A&A*, 674, A1
- Han, H., Huang, Y., Wang, B., et al. 2025, *Research in Astronomy and Astrophysics*, 25, 044001
- Ivezić, Z., Kahn, S. M., Tyson, J. A., et al. 2019, *ApJ*, 873, 111
- Jones, R. L., Slater, C. T., Moeyens, J., et al. 2018, *Icarus*, 303, 181
- Kaasalainen, M., & Torppa, J. 2001, *Icarus*, 153, 24
- Kaasalainen, M., Torppa, J., & Muinonen, K. 2001, *Icarus*, 153, 37
- Kubica, J., Denneau, L., Grav, T., et al. 2007, *Icarus*, 189, 151
- Lang, D., Hogg, D. W., Mierle, K., Blanton, M., & Roweis, S. 2010, *AJ*, 139, 1782
- Larsen, J. A. 2007, *AJ*, 133, 1247
- Larson, S., Beshore, E., Hill, R., et al. 2003, *BAAS*, 35, 982
- Liu, J., Soria, R., Wu, X.-F., Wu, H., & Shang, Z. 2021, *Anais da Academia Brasileira de Ciencias*, 93, 20200628
- Liu, Z., Gao, J., Gu, H., et al. 2025, *Research in Astronomy and Astrophysics*, 25, 044010
- Li, Z., et al. in preparation
- Li, N., et al. in preparation
- Mainzer, A. K., Masiero, J. R., Abell, P. A., et al. 2023, *PSJ*, 4, 224
- Morrison, D. 1992, *The Spaceguard Survey: Report of the NASA International Near-Earth-Object Detection Workshop*, NASA Technical Memorandum 107979
- Muinonen, K., Belskaya, I. N., Cellino, A., et al. 2010, *Icarus*, 209, 542

- Patterson, M. T., Bellm, E. C., Rusholme, B., et al. 2019, *PASP*, 131, 018001
- Schwamb, M. E., Jones, R. L., Yoachim, P., et al. 2023, *ApJS*, 266, 22
- Stokes, G. H., Evans, J. B., Viggh, H. E. M., Shelly, F. C., & Pearce, E. C. 2000, *Icarus*, 148, 21
- Tonry, J. L., Denneau, L., Heinze, A. N., et al. 2018, *PASP*, 130, 064505
- Vereš, P., & Chesley, S. R. 2017, *Icarus*, 296, 139
- Warner, B. D., Harris, A. W., & Pravec, P. 2009, *Icarus*, 202, 134
- Xiao, K., Li, Z., Huang, Y., et al. 2025, *Research in Astronomy and Astrophysics*, 25, 044006