

## 2A degree exam 2017–18, solutions

1. With  $z(x, y) = \ln(x^2 + y)$  we consider  $Z(s, t) = z(x(s, t), y(s, t))$  with  $x(s, t) = se^t$  and  $y = te^{-s}$ . Then the chain rule gives

$$\begin{aligned}\frac{\partial Z}{\partial s} &= \frac{\partial x}{\partial s} \frac{\partial z}{\partial x} + \frac{\partial y}{\partial s} \frac{\partial z}{\partial y} \\ \frac{\partial Z}{\partial t} &= \frac{\partial x}{\partial t} \frac{\partial z}{\partial x} + \frac{\partial y}{\partial t} \frac{\partial z}{\partial y}.\end{aligned}$$

The relevant partial derivatives are

$$\frac{\partial x}{\partial s} = e^t, \quad \frac{\partial x}{\partial t} = se^t, \quad \frac{\partial y}{\partial s} = -te^{-s}, \quad \frac{\partial y}{\partial t} = e^{-s}$$

and

$$\frac{\partial z}{\partial x} = \frac{2x}{x^2 + y}, \quad \frac{\partial z}{\partial y} = \frac{1}{x^2 + y}.$$

Putting these together

$$\begin{aligned}\frac{\partial Z}{\partial s} &= e^t \cdot \frac{2se^t}{(se^t)^2 + te^{-s}} - te^{-s} \cdot \frac{1}{(se^t)^2 + te^{-s}} = \frac{2se^{2t} - te^{-s}}{s^2e^{2t} + te^{-s}} \\ \frac{\partial Z}{\partial t} &= e^t \cdot \frac{2se^t}{(se^t)^2 + te^{-s}} + e^{-s} \cdot \frac{1}{(se^t)^2 + te^{-s}} = \frac{2se^{2t} + e^{-s}}{s^2e^{2t} + te^{-s}}.\end{aligned}$$

2. We seek to simplify the PDE by writing  $f(x, y) = F(u(x, y), v(x, y))$ . Then with  $u = xy^4$  and  $v = xy$  we have, using the chain rule,

$$\begin{aligned}\frac{\partial f}{\partial x} &= \frac{\partial u}{\partial x} \frac{\partial F}{\partial u} + \frac{\partial v}{\partial x} \frac{\partial F}{\partial v} = y^4 \frac{\partial F}{\partial u} + y \frac{\partial F}{\partial v} \\ \frac{\partial f}{\partial y} &= \frac{\partial u}{\partial y} \frac{\partial F}{\partial u} + \frac{\partial v}{\partial y} \frac{\partial F}{\partial v} = 4xy^3 \frac{\partial F}{\partial u} + x \frac{\partial F}{\partial v}.\end{aligned}$$

Substitute into the PDE  $xf_x - yf_y = 9xy$  to give

$$x(y^4 F_u + y F_v) - y(4xy^3 F_u + x F_v) = 9xy$$

which simplifies to  $-3xy^4 F_u = 9xy$  and using the change of variable to eliminate  $x$  and  $y$  we obtain the simplified PDE

$$\frac{\partial F}{\partial u} = -\frac{3v}{u}$$

a partial integration gives the general solution as

$$F(u, v) = -3v \log u + A(v)$$

where  $A$  is an arbitrary function of a single variable. The solution to the original PDE is therefore

$$f(x, y) = -3xy \log(xy^4) + A(xy).$$

3. (i) The Laplacian of  $f$  is

$$\nabla^2 f = \operatorname{div}(\operatorname{grad} f) = \frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} + \frac{\partial^2 f}{\partial z^2}.$$

- (ii) The gradient of  $1/r$  (using the chain rule and implicit differentiation to give  $r_x = x/r$  etc.) is

$$\nabla \left( \frac{1}{r} \right) = - \left( \frac{\partial r}{\partial x} \frac{1}{r^2}, \frac{\partial r}{\partial y} \frac{1}{r^2}, \frac{\partial r}{\partial z} \frac{1}{r^2} \right) = -\frac{1}{r^2} \left( \frac{\partial r}{\partial x}, \frac{\partial r}{\partial y}, \frac{\partial r}{\partial z} \right) = -\frac{1}{r^2} \left( \frac{x}{r}, \frac{y}{r}, \frac{z}{r} \right) = -\frac{\mathbf{x}}{r^3}$$

where  $\mathbf{x} = (x, y, z)$ . Now we calculate the divergence, using the nabla identity

$$\nabla \cdot (f\mathbf{F}) = \mathbf{F} \cdot \nabla f + f \nabla \cdot \mathbf{F}$$

and that  $\nabla \cdot \mathbf{x} = 3$ ,


$$\nabla^2 \left( \frac{1}{r} \right) = -\nabla \cdot \left( \frac{\mathbf{x}}{r^3} \right) = -\frac{1}{r^3} \nabla \cdot \mathbf{x} - \mathbf{x} \cdot \nabla \left( \frac{1}{r^3} \right) = -\frac{3}{r^3} - \mathbf{x} \cdot \left( -3 \frac{\mathbf{x}}{r^5} \right) = -\frac{3}{r^3} + \frac{3}{r^3} = 0$$

- (iii) The curl of the gradient is

$$\nabla \times \nabla \phi = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \partial_x & \partial_y & \partial_z \\ \phi_x & \phi_y & \phi_z \end{vmatrix} = (\phi_{zy} - \phi_{yz}, \phi_{xz} - \phi_{zx}, \phi_{yx} - \phi_{xy})$$

If  $\phi$  has continuous second derivatives then the order of differentiation doesn't matter and so by Clairaut's theorem  $\nabla \times \nabla \phi = \mathbf{0}$ .

4. The sketch is shown below.



The integral is written as a type-II integral ( $x$  integral first) and so we convert to type-I. The type-I description of the region is  $x^{1/4} \leq y \leq 3$  and  $1 \leq x \leq 81$ , therefore the integral can be written

$$I = \int_1^{81} \int_{x^{1/4}}^3 \frac{y^2}{x} dx dy = \int_1^{81} \int_{x^{1/4}}^3 \frac{y^2}{x} dy dx.$$

Performing the iterated integral gives

$$\int_1^{81} \left[ \frac{y^3}{3x} \right]_{x^{1/4}}^3 dx = \int_1^{81} \frac{9}{x} - \frac{1}{3x^{1/4}} dx = \left[ 9 \log x - \frac{4x^{3/4}}{9} \right]_1^{81} = 9 \log 81 - \frac{104}{9}$$

5. In polar coordinates, in the  $r$ - $\theta$  plane the region is rectangular with  $(r, \theta) \in [0, 2] \times [\pi/4, \pi/2]$ : the  $y$ -axis is the line  $\theta = \pi/2$  and the line  $y = x$ , in the first quadrant, is  $\theta = \pi/4$ ; the circle  $x^2 + y^2 = 4$  corresponds to  $r = 2$ . Therefore, substitution of  $x = r \cos \theta$  and  $y = r \sin \theta$  in the integrand and use of the Jacobian  $J = r$  (scale factor of areas) gives

$$\iint_D xy dx dy = \int_{\pi/4}^{\pi/2} \left( \int_0^2 (r \cos \theta) (r \sin \theta) r dr \right) d\theta$$

note that the integrand is separable and the region is rectangular so the integral can be written as the product of two one-dimensional integrals

$$\left[ \int_{\pi/4}^{\pi/2} \sin \theta \cos \theta d\theta \right] \left[ \int_0^2 r^3 dr \right] = \left[ -\frac{1}{4} \cos 2\theta \right]_{\pi/4}^{\pi/2} \left[ \frac{1}{4} r^4 \right]_0^2 = 1$$

6. In spherical polar coordinates the hemisphere is described by  $r \in [0, \sqrt{3}]$ ,  $\theta \in [0, 2\pi]$  and  $\phi \in [0, \pi/2]$ . Using spherical polar coordinates we note that  $x^2 + y^2 + z^2 = r^2$  and the Jacobian of the change of variables is  $J = r^2 \sin \phi$ , so the integral becomes

$$\iiint_{V'} (2 + 5r^2) r^2 \sin \phi dr d\theta d\phi$$

and since the integrand is separable and the region is cuboidal we can write the integral as the product of three one-dimensional integrals as follows

$$\left[ \int_0^{\sqrt{3}} (2 + 5r^2) r^2 dr \right] \left[ \int_0^{\pi/2} \sin \phi d\phi \right] \left[ \int_0^{2\pi} 1 d\theta \right] = \left[ \frac{2r^3}{3} + r^5 \right]_0^{\sqrt{3}} [-\cos \phi]_0^{\pi/2} [\theta]_0^{2\pi} = 22\pi\sqrt{3}.$$

7. The projection of the surface onto the  $xy$ -plane is the quadrilateral given ( $0 \leq x \leq 1$  and  $0 \leq y \leq x + 3$ ). The surface is the graph  $z = 25 - 3x - 2y$ , to convert the surface integral into a double integral we calculate

$$\sqrt{1 + z_x^2 + z_y^2} = \sqrt{1 + (-3)^2 + (-2)^2} = \sqrt{14}.$$

The quadrilateral is a type-I region so the surface integral becomes

$$\int_0^1 \left( \int_0^{x+3} y \sqrt{14} dy \right) dx = \int_0^1 \frac{1}{2} (x+3)^2 \sqrt{14} dx = \left[ \frac{1}{6} (x+3)^3 \sqrt{14} \right]_0^1 = \frac{37\sqrt{14}}{6}.$$

8. A parametric description of the given curve is  $\mathbf{r}(t) = (\cos t, \sin t)$  for  $t \in [-\pi/2, \pi/2]$ . We can check that all points on this curve lie on the circle  $x^2 + y^2 = 1$ , check that  $|\mathbf{r}'|^2 = 1$  (the calculation  $\cos^2 t + \sin^2 t = 1$  confirms this). The start, when  $t = -\pi/2$  gives the point  $(0, -1)$  and the end, when  $t = \pi/2$ , gives the point  $(0, 1)$ . The orientation is anticlockwise (the positive direction), we can confirm this by calculating  $\dot{\mathbf{r}}(-\pi/2) = (1, 0)$  (as  $\dot{\mathbf{r}} = (-\sin t, \cos t)$ ). Now we have

$$\int_C \mathbf{F} \cdot d\mathbf{x} = \int_{-\pi/2}^{\pi/2} (-\sin t, \cos t) \cdot (-\sin t, \cos t) dt = \int_{-\pi/2}^{\pi/2} \sin^2 t + \cos^2 t dt = \pi.$$

9. The divergence says that, given  $\mathbf{F}$ , a vector field in  $\mathbb{R}^3$  we have

$$\iint_S \mathbf{F} \cdot \mathbf{n} dS = \iiint_V \nabla \cdot \mathbf{F} dV$$

where  $S$  is a closed orientable surface enclosing the region  $V$ , with outward pointing unit normal  $\mathbf{n}$ .

In our case  $\nabla \cdot \mathbf{F} = -3x^2 - 2y + 6z$  and we can write the triple integral as the following iterated integral

$$\begin{aligned} \int_0^1 \left( \int_{-1}^1 \left( \int_{-1}^2 -3x^2 - 2y + 6z \, dz \right) dy \right) dx &= \int_0^1 \left( \int_{-1}^1 [-3x^2z - 2yz + 3z^2]_{-1}^2 dy \right) dx \\ &= \int_0^1 \left( \int_{-1}^1 -9x^2 - 6y + 9 \, dy \right) dx \\ &= \int_0^1 [-9x^2y - 3y^2 + 9y]_{-1}^1 dx \\ &= \int_0^1 -18x^2 + 18 \, dx \\ &= [-6x^3 + 18x]_0^1 = 12 \end{aligned}$$