



University
of Glasgow

EXAMINATION FOR THE DEGREES OF
M.A. AND B.Sc.

Mathematics 2B - Linear Algebra

An electronic calculator may be used provided that it does not have a facility for either textual storage or display, or for graphical display.

Candidates must attempt all questions.

Specimen exam paper - SOLUTIONS

1. We consider a subset of \mathbb{R}^3 in the form

$$\mathcal{C} = \{[1, 1, 0], [0, 1, 1], [0, 0, 1]\}$$

- (i) Show that \mathcal{C} forms a basis for \mathbb{R}^3 .
- (ii) Construct the change-of-basis matrix $P_{\mathcal{C} \leftarrow \mathcal{E}}$, where \mathcal{E} is the standard basis for \mathbb{R}^3

- (iii) Consider the vector $\mathbf{w} = \begin{bmatrix} 1 \\ 2 \\ -1 \end{bmatrix} \in \mathbb{R}^3$. Using your solution to part(ii), or otherwise, compute the coordinates of \mathbf{w} with respect to the basis \mathcal{C} .

•

- (i) We first show that the vectors in \mathcal{C} are linearly independent by considering solutions to the equation

$$c_1 \mathbf{v}_1 + c_2 \mathbf{v}_2 + c_3 \mathbf{v}_3 = \mathbf{0},$$

where c_1, c_2 and c_3 are scalars. We form the augmented matrix

$$\left[\begin{array}{ccc|c} 1 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 0 \end{array} \right] \rightarrow \left[\begin{array}{ccc|c} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{array} \right]$$

Hence, the only solution to the homogeneous system is $c_1 = c_2 = c_3 = 0$, so the vectors are linearly independent. Since $\dim(\mathbb{R}^3) = 3$ and we have three

linearly independent vectors, these vectors must form a basis for \mathbb{R}^3 using a theorem from lectures.

(ii) We let C be the matrix with columns formed from the coordinate vectors of the vectors in \mathcal{C} with respect to the standard basis \mathcal{E} . Further, we let E be the matrix with columns as the vectors in the standard basis \mathcal{E} . We compute the change-of-basis matrix using the Gauss-Jordan method

$$\begin{aligned} [C|E] &= \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 1 & 0 & 0 \\ 1 & 1 & 0 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 & 0 & 1 \end{array} \right] \rightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & -1 & 1 & 0 \\ 0 & 1 & 1 & 0 & 0 & 1 \end{array} \right] \\ &\rightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & -1 & 1 & 0 \\ 0 & 0 & 1 & 1 & -1 & 1 \end{array} \right] = [\mathbb{I}|P_{\mathcal{C} \leftarrow \mathcal{E}}] \end{aligned}$$

so we have

$$P_{\mathcal{C} \leftarrow \mathcal{E}} = \begin{bmatrix} 1 & 0 & 0 \\ -1 & 1 & 0 \\ 1 & -1 & 1 \end{bmatrix}$$

(iii) We have $[\mathbf{w}]_{\mathcal{E}} = \begin{bmatrix} 1 \\ 2 \\ -1 \end{bmatrix}$. Hence, using the change-of-basis matrix

$$[\mathbf{w}]_{\mathcal{C}} = P_{\mathcal{C} \leftarrow \mathcal{E}} [\mathbf{w}]_{\mathcal{E}} = \begin{bmatrix} 1 & 0 & 0 \\ -1 & 1 & 0 \\ 1 & -1 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \\ -1 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \\ -2 \end{bmatrix}.$$

2. Consider two $n \times n$ matrices C and D , where D is invertible. Prove that the matrix C is non-invertible if and only if the matrix DC is non-invertible.

. Suppose C is non-invertible. Then C must have a zero eigenvalue by Thm 4.16. We let \mathbf{x} be the eigenvector corresponding to the zero eigenvalue and so $C\mathbf{x} = 0\mathbf{x} = \mathbf{0}$. Multiplying both sides by the matrix D , we obtain $DC\mathbf{x} = D\mathbf{0} = \mathbf{0} = 0\mathbf{x}$ and so DC has a zero eigenvalue. Hence, DC must be non-invertible by Thm 4.16.

Conversely suppose DC is non-invertible. Then by Thm 4.16 DC has a zero eigenvalue with corresponding eigenvector \mathbf{y} and so $DC\mathbf{y} = \mathbf{0} = 0\mathbf{y}$. Since D is invertible, then $C\mathbf{y} = D^{-1}\mathbf{0} = 0\mathbf{y}$ and so C is non-invertible (by Thm 4.16) as it has a zero eigenvalue.

Hence, C is non-invertible if and only if the matrix DC is non-invertible.

3. Consider an $n \times n$ matrix A . Suppose that A has an eigenvalue λ . Show that the eigenspace E_λ is a subspace of \mathbb{R}^n .

. By definition the eigenspace E_λ contains the eigenvectors of A with eigenvalue λ as well as the zero vector. Consider two vectors \mathbf{x} and \mathbf{y} which belong to E_λ . Hence $A\mathbf{x} = \lambda\mathbf{x}$ and $A\mathbf{y} = \lambda\mathbf{y}$. We consider

$$A(\mathbf{x} + \mathbf{y}) = A\mathbf{x} + A\mathbf{y} = \lambda\mathbf{x} + \lambda\mathbf{y} = \lambda(\mathbf{x} + \mathbf{y}),$$

so $\mathbf{x} + \mathbf{y} \in E_\lambda$ and the subset is closed under addition. Consider a scalar c , then

$$A(c\mathbf{x}) = cA\mathbf{x} = c\lambda\mathbf{x} = \lambda(c\mathbf{x}),$$

so $c\mathbf{x} \in E_\lambda$ and the subset is closed under scalar multiplication. Hence, E_λ forms a subspace of \mathbb{R}^n .

4. Let U be the subspace of \mathbb{R}^4 spanned by

$$(1, 1, 0, 0), \quad (2, 0, 1, 0), \quad (1, 0, 2, 1)$$

Use the Gram-schmidt Process to find an orthogonal basis for U .

. We first show that the vectors in \mathcal{C} are linearly independent by considering solutions to the equation

$$c_1\mathbf{u}_1 + c_2\mathbf{u}_2 + c_3\mathbf{u}_3 = \mathbf{0},$$

where c_1, c_2 and c_3 are scalars. We form the augmented matrix

$$\left[\begin{array}{ccc|c} 1 & 2 & 1 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & 1 & 0 \end{array} \right]$$

Gaussian-Jordan elimination gives the final augmented form as

$$\left[\begin{array}{ccc|c} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right],$$

so if $c_1\mathbf{u}_1 + c_2\mathbf{u}_2 + c_3\mathbf{u}_3 = \mathbf{0}$, then $c_1 = c_2 = c_3 = 0$. So $\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3$ is a basis for U .

We now apply the Gram-Schmidt process to construct an orthogonal basis:

$$\begin{aligned}
\mathbf{v}_1 &= \mathbf{u}_1 = \begin{bmatrix} 1 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \\
\mathbf{v}_2 &= \mathbf{u}_2 - \frac{\mathbf{u}_2 \cdot \mathbf{v}_1}{\mathbf{v}_1 \cdot \mathbf{v}_1} \mathbf{v}_1 \\
&= \begin{bmatrix} 2 \\ 0 \\ 1 \\ 0 \end{bmatrix} - \frac{2}{2} \begin{bmatrix} 1 \\ 1 \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 1 \\ -1 \\ 1 \\ 0 \end{bmatrix} \\
\mathbf{v}_3 &= \mathbf{u}_3 - \frac{\mathbf{u}_3 \cdot \mathbf{v}_1}{\mathbf{v}_1 \cdot \mathbf{v}_1} \mathbf{v}_1 - \frac{\mathbf{u}_3 \cdot \mathbf{v}_2}{\mathbf{v}_2 \cdot \mathbf{v}_2} \mathbf{v}_2 \\
&= \begin{bmatrix} 1 \\ 0 \\ 2 \\ 1 \end{bmatrix} - \frac{1}{2} \begin{bmatrix} 1 \\ 1 \\ 0 \\ 0 \end{bmatrix} - \frac{3}{3} \begin{bmatrix} 1 \\ -1 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} -1/2 \\ 1/2 \\ 1 \\ 1 \end{bmatrix}.
\end{aligned}$$

So $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3$ is an orthonormal basis for U .

5. Consider the transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by

$$(x_1, x_2) \rightarrow (x_1 + 2x_2, -x_1).$$

Show that T is a linear transformation.

. Let $\mathbf{x}, \mathbf{y} \in \mathbb{R}^2$ and $c \in \mathbb{R}$, where $\mathbf{x} = (x_1, x_2)$, and $\mathbf{y} = (y_1, y_2)$. We first show T preserves addition, so we consider

$$\begin{aligned}
T(\mathbf{x} + \mathbf{y}) &= T(x_1 + y_1, x_2 + y_2) = ((x_1 + y_1) + 2(x_2 + y_2), -(x_1 + y_1)) \\
&= (x_1 + 2x_2 + y_1 + 2y_2, -x_1 - y_1) = (x_1 + 2x_2, -x_1) + (y_1 + 2y_2, -y_1) = T(\mathbf{x}) + T(\mathbf{y})
\end{aligned}$$

and so T preserves addition. We then show T preserves scalar multiplication, so we consider

$$T(c\mathbf{x}) = T(cx_1, cx_2) = (cx_1 + 2cx_2, -cx_1) = c(x_1 + 2x_2, -x_1) = cT(\mathbf{x})$$

so T also preserves scalar multiplication and hence T is a linear transformation.

6. Let q be the quadratic form defined by

$$q(x_1, x_2) = -3x_1^2 + 5x_2^2 - 6x_1x_2.$$

(i) Find a non-singular change of variables $\mathbf{x} = Q\mathbf{y}$, where

$$\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} \text{ and } \mathbf{y} = \begin{bmatrix} y_1 \\ y_2 \end{bmatrix}$$

such that $q(y_1, y_2) = \lambda_1 y_1^2 + \lambda_2 y_2^2$, for some $\lambda_1, \lambda_2 \in \mathbb{R}$.

(ii) Find a diagonal matrix D such $Q^T A Q = D$, where A is the matrix associated to q and determine the rank and the signature of q .

.

(i) The quadratic can be written as $\mathbf{x}^T A \mathbf{x}$ where $\mathbf{x} \in \mathbb{R}^2$ and

$$A = \begin{bmatrix} -3 & -3 \\ -3 & 5 \end{bmatrix}.$$

The eigenvalues of A satisfy $\lambda^2 - 2\lambda - 24 = (\lambda - 6)(\lambda + 4)$, hence $\lambda_1 = 6, \lambda_2 = -4$.

The corresponding eigenspaces E_6 and E_{-4} are spanned by $\mathbf{v}_1 = \frac{1}{\sqrt{10}} \begin{bmatrix} 1 \\ -3 \end{bmatrix}$ and

$\mathbf{v}_2 = \frac{1}{\sqrt{10}} \begin{bmatrix} 3 \\ 1 \end{bmatrix}$ respectively. These vectors are orthonormal and we construct an orthogonal matrix Q whose columns are \mathbf{v}_1 and \mathbf{v}_2 :

$$Q = \frac{1}{\sqrt{10}} \begin{bmatrix} 1 & 3 \\ -3 & 1 \end{bmatrix}$$

Applying orthogonal diagonalisation, $\mathbf{x}^T A \mathbf{x} = \mathbf{x}^T Q Q^T A Q Q^T \mathbf{x} = (Q^T \mathbf{x})^T D Q^T \mathbf{x} = \mathbf{y}^T D \mathbf{y}$, where $D = \begin{bmatrix} 6 & 0 \\ 0 & -4 \end{bmatrix}$ and so the quadratic in the new variables is

$$6(y_1)^2 - 4(y_2)^2,$$

where $Q\mathbf{y} = \mathbf{x}$, since $Q^T = Q^{-1}$.

(ii) From part (i) $D = \begin{bmatrix} 6 & 0 \\ 0 & -4 \end{bmatrix}$ and the rank of q is 2 and the signature is 0.

END]