01 ORTHOPEDIC SIGNAL 85

For chronic stroke hand rehab, sensory ES and Alt-EMG are equally effective.

**Why this matters*: This RCT compared two types of electrical stimulation, both combined with conventional therapy, for improving upper-limb motor function.*

NeuroRehabilitation

ALSO TODAY

01 ORTHO Journal of physiotherapy

01 ORTHO Journal of Rehabilitation Medicine

02 NEURO Sports medicine (Auckland, N.Z.)

05 GERIATRIC BMC Geriatrics

Adding PBMT to exercise offers no extra benefit for chronic non-specific low back pain.

**Why this matters*: This RCT tested if adding light therapy to exercise improved pain or disability in people with long-term lower back issues.*

SIGNAL 85

Journal of physiotherapy

Supervised telerehab with Pilates and self-massage aids fibromyalgia recovery.

**Why this matters*: This RCT evaluated a 6-month remote program combining exercise and self-care for fibromyalgia impact and symptoms.*

SIGNAL 80
Journal of Rehabilitation Medicine

Exercise improves walking economy in healthy adults; effects are smaller in stroke patients.

**Why this matters*: This systematic review synthesised evidence on how exercise training affects the energy used for walking across different health conditions.*

SIGNAL 80
Sports medicine (Auckland, N.Z.)

In frail older adults, swallowing problems link to poor physical function and falls.

**Why this matters*: This study explored the relationship between difficulty swallowing and physical performance and fall risk in older individuals.*

SIGNAL 80

BMC Geriatrics