01 ORTHOPEDIC SIGNAL 75

Prehabilitation exercise may improve ankle surgery outcomes and speed return to sport.

**Why this matters*: A 4-week exercise program before arthroscopic Broström procedure for chronic ankle instability showed better functional recovery and faster return to sports at 2 years.*

BMC Sports Sci Med Rehabil

ALSO TODAY

02 NEURO — Physiotherapy theory and practice

05 GERIATRIC — BMC musculoskeletal disorders

04 PEDIATRIC — medRxiv

Therapists' confidence and resources impact stroke walking program delivery.

**Why this matters*: Therapists believe high-dose walking training is important but need confidence and resources to ensure patients complete it.*

SIGNAL 75

Physiotherapy theory and practice

Early exercise adherence after hip fracture surgery links to better function.

**Why this matters*: Moderate early functional exercise adherence in older adults after hip fracture surgery was associated with better 3-month hip functional recovery.*

SIGNAL 75
BMC musculoskeletal disorders

Robot-assisted walking in schools may increase participation for children with physical disabilities.

This pilot randomized crossover trial found that robot-assisted walking in schools was feasible and showed potential for improving participation for children with physical disabilities.

SIGNAL 70

medRxiv

Five papers a day, read so you don't have to.

Cadence Evidence

incadencept.com