02 NEUROLOGICAL

SIGNAL 85

Consider NMES to improve lower limb function in post-stroke patients.

A systematic review and meta-analysis found neuromuscular electrical stimulation improved several measures of lower limb mobility and daily activities in post-stroke patients.

AJPM&R

ALSO TODAY

02 NEURO — BMC Sports Sci Med Rehabil

02 NEURO — Disability and rehabilitation

02 NEURO — Clin Biomech

03 SPORTS — BMC Sports Sci Med Rehabil

For stroke balance, is whole-body vibration most beneficial?

A network meta-analysis of stroke rehabilitation interventions found whole-body vibration most beneficial for balance, while other exercise types improved walking endurance.

SIGNAL 80

BMC Sports Sci Med Rehabil

No single observational gait tool is fully recommended for neurological populations.

A COSMIN systematic review and meta-analysis found no observational gait analysis tool was fully recommended for use in neurological populations, though some showed potential for stroke.

Balance confidence and objective balance are distinct in chronic stroke.

A secondary analysis of adults with chronic stroke found small correlations between balance confidence and objective balance and gait measures.

SIGNAL 75
Clin Biomech

Can machine learning predict peak ACL force from biomechanical parameters?

An interpretable machine learning model demonstrated high accuracy in predicting peak ACL force during side-cutting maneuvers in male football and basketball athletes.

SIGNAL 75
BMC Sports Sci Med Rehabil

Five papers a day, read so you don't have to.

Cadence Evidence incadencept.com