03 SPORTS & ATHLETIC SIGNAL 75

For refractory plantar fasciopathy, exercise is not inferior to microtenotomy.

*Why this matters: *This randomized controlled trial compared radiofrequency microtenotomy with heavy-slow resistance training for pain in refractory plantar fasciopathy.**

Sports Medicine - Open

ALSO TODAY

04 PEDIATRIC medRxiv

07 MANUAL BMC Musculoskeletal Disorders

01 ORTHO BMC Musculoskeletal Disorders

05 GERIATRIC Aging Clin Exp Res

Children with limb loss after trauma may need emotional support.

*Why this matters: *This retrospective analysis examined the physical and psychosocial outcomes in children with earthquake-related limb loss two years post-amputation.**

SIGNAL 75

medRxiv

Can assessing a patient's self-correction ability predict scoliosis outcomes?

*Why this matters: *This prospective study quantified active self-correction during physiotherapeutic scoliosis-specific exercises and its association with Cobb angle change.**

SIGNAL 75
BMC Musculoskeletal Disorders

Adding hip-ankle training after TKA offers only modest, short-term benefits.

*Why this matters: *This randomized controlled trial investigated the effect of multi-joint lower-limb training on function after total knee arthroplasty.**

SIGNAL 70

BMC Musculoskeletal Disorders

Malnutrition is common and linked to frailty in older diabetic outpatients.

*Why this matters: *This retrospective study examined the prevalence of GLIM-defined malnutrition and its association with frailty in older outpatients with diabetes.**

SIGNAL 70

Aging Clin Exp Res