01 ORTHOPEDIC

SIGNAL 85

Exercise helps reduce kinesiophobia and improve function after TKA.

This systematic review examined if exercise-based rehabilitation could reduce kinesiophobia and improve knee function after total knee arthroplasty.

BMC musculoskeletal disorders

01

ALSO TODAY

06	CARDIOPULM	The Lancet
02	NEURO	NeuroRehabilitation
02	NEURO	Disability and rehabilitation
02	NEURO	Topics in stroke rehabilitation

Prone positioning and HFOV are not first-line treatments for early PARDS.

This randomized controlled trial investigated whether prone positioning or high-frequency oscillatory ventilation improved ventilator-free days in paediatric acute respiratory distress syndrome.

SIGNAL 80
The Lancet

Hybrid technology is a feasible option for upper limb neurorehabilitation in stroke.

This scoping review explored the use of hybrid technology in upper limb neurorehabilitation for adult stroke patients in hospital settings.

SIGNAL 75

NeuroRehabilitation

Psychosocial interventions for ABI need three key components.

This realist evaluation aimed to identify essential program theories for psychosocial interventions in young adults with acquired brain injury.

SIGNAL 75

Disability and rehabilitation

Could timing stroke rehab closer to sleep improve motor recovery?

This review explored the hypothesis that scheduling motor therapy close to sleep could enhance motor memory consolidation and recovery after stroke.

Five papers a day, read so you don't have to.

Cadence Evidence incadencept.com