01 ORTHOPEDIC

SIGNAL 85

# Prioritize short-foot exercise in early ACLR rehab for better hip control.

*Why this matters: This exploratory RCT compared short-foot and toe-curl exercises in ACLR patients, observing hip adduction during single-leg landing.*

Phys Ther Sport

01

ALSO TODAY

02 NEURO — Physiotherapy theory and practice

02 NEURO — medRxiv

05 GERIATRIC — BMC Geriatrics

06 CARDIOPULM — Braz J Phys Ther

# Use the Stair Test or Step Test to monitor mobility in MS patients.

*Why this matters: This study evaluated the reliability and validity of the Stair Test and Step Test for monitoring mobility in individuals with multiple sclerosis.*

# Personalize BCI parameters for stroke patients; MI timing may not matter.

*Why this matters: This single-case design study explored individual learning trajectories and the effect of motor imagery timing in network-based BCI rehabilitation for stroke patients.*

SIGNAL 75
medRxiv

# Protein-enriched diets may not improve early recovery from fragility fractures.

*Why this matters: *This RCT investigated if a protein-enriched diet during acute hospital stays improved recovery in older adults with fragility fractures.**

SIGNAL 75

BMC Geriatrics

# Use the 5STS test to assess muscle strength and readmission risk in severe COPD exacerbation.

*Why this matters: This study validated the five-repetition sit-to-stand (5STS) test for patients with severe COPD exacerbation, exploring its prognostic value for readmission risk.*

SIGNAL 75

Braz J Phys Ther