

# Take Home Exam II

Xingche Guo

November 28, 2017

## 1 Introduction

The main purpose of this study is to learn the different parametric methods, (e.g. **Wald's Theory**, **Inversion of Likelihood Ratio**, and **Parametric Bootstrap**), for constructing confident intervals of the parameters in Inverse Gaussian distribution under different conditions (e.g. differences in sample size  $n$  and in dispersion parameter  $\lambda$ ). On the other hand, this research gives me a chance to practice conducting a Monte Carlo study and to learn how to compare methods by different criteria.

I'm interested in doing this because this study is very meaningful. The results tells me the pros and cons of these methods under different conditions, which might give me a better idea of selecting methods to conduct confidence intervals. This study also shows how the sample size  $n$  and dispersion parameter  $\lambda$  affect the confidence interval features. Besides, it sheds light on determining how big the Monte Carlo sample size  $M$  and bootstrap sample size  $N$  should be, which are all very useful in my later research.

## 2 Methods

### 2.1 Method 1

The density function of an Inverse Gaussian distribution can be written as:

$$f(y|\mu, \lambda) = \sqrt{\frac{\lambda}{2\pi y^3}} \exp\left(-\frac{\lambda(y - \mu)^2}{2\mu^2 y}\right); \quad y > 0; \mu > 0; \lambda > 0$$

Thus, the joint density of iid r.v.  $Y_1, \dots, Y_n$  is:

$$f(\mathbf{y}|\mu, \lambda) = \prod_{i=1}^n f(y_i|\mu, \lambda)$$

The log likelihood function of the joint density is:

$$\begin{aligned}
l(\mu, \lambda; \mathbf{y}) &= \sum_{i=1}^n \log f(y_i | \mu, \lambda) \\
&= \sum_{i=1}^n \left( \frac{1}{2} \log \frac{\lambda}{2\pi y^3} - \frac{\lambda(y_i^2 - 2y_i\lambda + \mu^2)}{2\mu^2 y_i} \right) \\
&= \frac{n}{2} \log \lambda - \frac{\lambda}{2\mu^2} \sum_{i=1}^n y_i + \frac{n\lambda}{\mu} - \frac{\lambda}{2} \sum_{i=1}^n \frac{1}{y_i} - \frac{1}{2} \sum_{i=1}^n \log 2\pi y_i^3
\end{aligned}$$

Compute the MLE by solving:

$$\begin{cases} \frac{\partial l}{\partial \mu} = 0 \\ \frac{\partial l}{\partial \lambda} = 0 \end{cases}$$

The MLE of  $\mu$  and  $\lambda$  is:

$$\begin{cases} \hat{\mu} = \frac{\sum_{i=1}^n y_i}{n} \\ \hat{\lambda} = \left( \frac{\sum_{i=1}^n \frac{1}{y_i}}{n} - \frac{n}{\sum_{i=1}^n y_i} \right)^{-1} \end{cases}$$

The Fisher's Information Matrix is:

$$I(\mu, \lambda) = \begin{pmatrix} \frac{n\lambda}{\mu^3} & 0 \\ 0 & \frac{n}{2\lambda^2} \end{pmatrix}$$

By Wald's theory, the 95% Confidence Interval of  $\mu$  and  $\lambda$  is:

$$\begin{cases} \mu \in \hat{\mu} \pm 1.96 \times \sqrt{\frac{\hat{\mu}^3}{n\hat{\lambda}}} \\ \lambda \in \hat{\lambda} \pm 1.96 \times \sqrt{\frac{2\hat{\lambda}^2}{n}} \end{cases}$$

## 2.2 Method 2

For  $\lambda$ , using the profile likelihood thought, define:  $l_1(\lambda) = l(\hat{\mu}, \lambda)$ . By Wilk's theorem, we have:

$$-2(l_1(\lambda_0) - l_1(\hat{\lambda})) \sim \chi_1^2, \text{ when } n \rightarrow \infty.$$

Thus, by the inversion of likelihood ratio statistics, the 95% Confidence Interval of  $\lambda$  is:

$$\{\lambda \in R^+ : -2(l_1(\lambda) - l_1(\hat{\lambda})) \leq \chi_1^2(0.95)\}$$

Define:

$$T_\lambda = -2(l_1(\lambda) - l_1(\hat{\lambda}))$$

Then:

$$\begin{aligned} T_\lambda &= -2[(\frac{n}{2}\log\lambda + \frac{n\lambda}{2\hat{\mu}} - \frac{\lambda}{2}\sum_{i=1}^n \frac{1}{y_i}) - (\frac{n}{2}\log\hat{\lambda} + \frac{n\hat{\lambda}}{2\hat{\mu}} - \frac{\hat{\lambda}}{2}\sum_{i=1}^n \frac{1}{y_i})] \\ &= (\sum_{i=1}^n \frac{1}{y_i} - \frac{n}{\hat{\mu}})(\lambda - \hat{\lambda}) - n\log \frac{\lambda}{\hat{\lambda}} \end{aligned}$$

For  $\mu$ , notice that:

$$\frac{\partial l}{\partial \lambda} = 0 \rightarrow \lambda = (\frac{\hat{\mu}}{\mu^2} - \frac{2}{\mu} + \frac{\sum_{i=1}^n \frac{1}{y_i}}{n})^{-1}$$

So, we define:

$$\tilde{\lambda} \equiv \tilde{\lambda}(\mu) = (\frac{\hat{\mu}}{\mu^2} - \frac{2}{\mu} + \frac{\sum_{i=1}^n \frac{1}{y_i}}{n})^{-1}$$

Then define:  $l_2(\mu) = l(\mu, \tilde{\lambda})$ , similarly, the 95% Confidence Interval of  $\mu$  is:

$$\{\mu \in R^+ : -2(l_2(\mu) - l_2(\hat{\mu})) \leq \chi_1^2(0.95)\}$$

Define:

$$T_\mu = -2(l_2(\mu) - l_2(\hat{\mu}))$$

Then:

$$T_\mu = -2[(\frac{n}{2}\log\tilde{\lambda} - \frac{\tilde{\lambda}}{2\mu^2}\sum_{i=1}^n y_i + \frac{n\tilde{\lambda}}{\mu} - \frac{\tilde{\lambda}}{2}\sum_{i=1}^n \frac{1}{y_i}) - (\frac{n}{2}\log\hat{\lambda} + \frac{n\hat{\lambda}}{2\hat{\mu}} - \frac{\hat{\lambda}}{2}\sum_{i=1}^n \frac{1}{y_i})]$$

To draw a conclusion, the 95% Confidence Interval of  $\mu$  and  $\lambda$  is:

$$\begin{cases} \mu \in \{\mu \in R^+ : T_\mu \leq \chi_1^2(0.95)\} \\ \lambda \in \{\lambda \in R^+ : T_\lambda \leq \chi_1^2(0.95)\} \end{cases}$$

where  $T_\mu$  and  $T_\lambda$  are defined above.

## 2.3 Method 3

By Maximum likelihood and parametric bootstrap using the comparison function:  $h(\theta, \hat{\theta}) = n^{1/2}(\hat{\theta} - \theta)$ . We can write down the algorithm as follow:

- Step1: Calculate  $\hat{\mu}$  and  $\hat{\lambda}$  (the MLE of  $\mu$  and  $\lambda$  using the original data).
- Step2: Generate observations  $\mathbf{y}_m^* = (y_{1m}^*, \dots, y_{nm}^*)$ ,  $m = 1, \dots, N$ , from the fitted model  $Inv.Gauss(\hat{\mu}, \hat{\lambda})$ . ( $N$  will be decided later).

- Step3: Calculate  $\mu_m^*$  and  $\lambda_m^*$  (the MLE of  $\mu$  and  $\lambda$  using the  $m$ th generated data).
- Step4: Calculate  $\mu_{[(N+1)(0.025)]}^*$ ,  $\mu_{[(N+1)(0.975)]}^*$ ,  $\lambda_{[(N+1)(0.025)]}^*$ ,  $\lambda_{[(N+1)(0.975)]}^*$  (e.g.  $\mu_{[(N+1)(0.025)]}^*$  is the  $[(N+1)(0.025)]$ th order statistics of  $\{\mu_m^*\}_{m=1}^N$ ).

Then the 95% Confidence Interval of  $\mu$  and  $\lambda$  is:

$$\begin{cases} \mu \in (2\hat{\mu} - \mu_{[(N+1)(0.975)]}^*, 2\hat{\mu} - \mu_{[(N+1)(0.025)]}^*) \\ \lambda \in (2\hat{\lambda} - \lambda_{[(N+1)(0.975)]}^*, 2\hat{\lambda} - \lambda_{[(N+1)(0.025)]}^*). \end{cases}$$

## 2.4 Method 4

By Maximum likelihood and parametric bootstrap using the comparison function:  $h(\theta, \hat{\theta}) = n^{1/2}(\frac{\theta}{\hat{\theta}} - 1)$ . The algorithm is the same as in Method 3.

The 95% Confidence Interval of  $\mu$  and  $\lambda$  is:

$$\begin{cases} \mu \in (\frac{\hat{\mu}^2}{\mu_{[(N+1)(0.975)]}^*}, \frac{\hat{\mu}^2}{\mu_{[(N+1)(0.025)]}^*}) \\ \lambda \in (\frac{\hat{\lambda}^2}{\lambda_{[(N+1)(0.975)]}^*}, \frac{\hat{\lambda}^2}{\lambda_{[(N+1)(0.025)]}^*}). \end{cases}$$

## 2.5 Method 5

By Maximum likelihood and parametric bootstrap using the comparison function:  $h(\theta, \hat{\theta}) = \frac{\hat{\theta} - \theta}{V(\hat{\theta})^{1/2}}$ . Only step 4 of the algorithm will change. We now need to calculate  $(\frac{\theta^* - \hat{\theta}}{V(\theta^*)^{1/2}})_{[(N+1)(0.025)]}$  and  $(\frac{\theta^* - \hat{\theta}}{V(\theta^*)^{1/2}})_{[(N+1)(0.975)]}$  instead of  $\theta_{[(N+1)(0.025)]}^*$  and  $\theta_{[(N+1)(0.975)]}^*$ , where  $\theta$  denote  $\mu$  or  $\lambda$ .

Thus, the 95% Confidence Interval of  $\mu$  and  $\lambda$  is:

$$\begin{cases} \mu \in (\hat{\mu} - [V(\hat{\mu})]^{1/2}(\frac{\mu^* - \hat{\mu}}{V(\mu^*)^{1/2}})_{[(N+1)(0.975)]}, \hat{\mu} - [V(\hat{\mu})]^{1/2}(\frac{\mu^* - \hat{\mu}}{V(\mu^*)^{1/2}})_{[(N+1)(0.025)]}) \\ \lambda \in (\hat{\lambda} - [V(\hat{\lambda})]^{1/2}(\frac{\lambda^* - \hat{\lambda}}{V(\lambda^*)^{1/2}})_{[(N+1)(0.975)]}, \hat{\lambda} - [V(\hat{\lambda})]^{1/2}(\frac{\lambda^* - \hat{\lambda}}{V(\lambda^*)^{1/2}})_{[(N+1)(0.025)]}). \end{cases}$$

Where:

$$V(\tilde{\mu}) = \frac{\tilde{\mu}^3}{n\tilde{\lambda}} ; \quad V(\tilde{\lambda}) = \frac{2\tilde{\lambda}^2}{n}.$$

( $\tilde{\mu}$  denotes  $\hat{\mu}$  or  $\mu^*$ ,  $\tilde{\lambda}$  denotes  $\hat{\lambda}$  or  $\lambda^*$ ).

Note that:

$$\begin{aligned}
& \hat{\lambda} - [V(\hat{\lambda})]^{1/2} \left( \frac{\lambda^* - \hat{\lambda}}{V(\lambda^*)^{1/2}} \right)_{[(N+1)(0.975)]} \\
&= \hat{\lambda} - \sqrt{\frac{2}{n}} \hat{\lambda} \left( \frac{\lambda^* - \hat{\lambda}}{\sqrt{\frac{2}{n}} \lambda^*} \right)_{[(N+1)(0.975)]} \\
&= \hat{\lambda} - \hat{\lambda} \left( 1 - \frac{\hat{\lambda}}{\lambda^*} \right)_{[(N+1)(0.975)]} \\
&= \hat{\lambda} - \hat{\lambda} \left( 1 - \frac{\hat{\lambda}}{\lambda_{[(N+1)(0.975)]}^*} \right) \\
&= \frac{\hat{\lambda}^2}{\lambda_{[(N+1)(0.975)]}^*}.
\end{aligned}$$

Similarly:

$$\begin{aligned}
& \hat{\lambda} - [V(\hat{\lambda})]^{1/2} \left( \frac{\lambda^* - \hat{\lambda}}{V(\lambda^*)^{1/2}} \right)_{[(N+1)(0.025)]} \\
&= \frac{\hat{\lambda}^2}{\lambda_{[(N+1)(0.025)]}^*}.
\end{aligned}$$

Hence, method 4 and method 5 will have the same Confidence Interval for  $\lambda$  given the same seed of randomness.

## 2.6 Method 6

By Method of Moment estimation and parametric bootstrap using the comparison function:  $h(\theta, \hat{\theta}) = n^{1/2}(\hat{\theta} - \theta)$ . The algorithm will be similar, the only difference is to replace all the  $\hat{\mu}$  and  $\hat{\lambda}$  by  $\mu_{MME}$  and  $\lambda_{MME}$ .

$$\begin{cases} \mu_{MME} = \frac{1}{n} \sum_{i=1}^n Y_i \\ \lambda_{MME} = \frac{\mu_{MME}^3}{\frac{1}{n} \sum_{i=1}^n Y_i^2 - \mu_{MME}^2} \end{cases}$$

Thus, the 95% Confidence Interval of  $\mu$  and  $\lambda$  is:

$$\begin{cases} \mu \in (2\mu_{MME} - \mu_{MME,[(N+1)(0.975)]}^*, 2\mu_{MME} - \mu_{MME,[(N+1)(0.025)]}^*) \\ \lambda \in (2\lambda_{MME} - \lambda_{MME,[(N+1)(0.975)]}^*, 2\lambda_{MME} - \lambda_{MME,[(N+1)(0.025)]}^*) \end{cases}$$

## 2.7 R Functions and Packages Used

- Generate Inverse Gaussian distribution: `statmod::rinvgauss(x, mean, shape)`.
- All the other R functions are written by me (In Appendix).

## 2.8 Choose Bootstrap — N

There are too many combinations of experiments:  $\lambda \in \{2, 4, 8, 12\}$ ,  $n \in \{10, 25, 50, 100, 500\}$ ,  $method \in \{3, 4, 5, 6\}$ . It's not efficient to find the best N for all these combinations. So, I use the thought of "Latin Square Design" to compute the confidence interval width of the following combinations when  $N \in \{50, 100, 200, 300, \dots, 10000\}$ . The results are in figure[1] and figure[2].

	method3	method4	method5	method6
$\lambda = 2$	n=10	n=50	n=100	n=500
$\lambda = 4$	n=50	n=100	n=500	n=10
$\lambda = 8$	n=100	n=500	n=10	n=50
$\lambda = 12$	n=500	n=10	n=50	n=100

The figure[1] and figure[2] show that the CI widths converge for most of the combinations when  $N=2000$ . Thus, we pick the bootstrap N to be 2000.

## 2.9 Choose Monte Carlo — M

In order to make the comparison between different combinations of experiments, we need to find one unified M. In this case, I compute:

- the coverage rate of  $\mu$ : given  $n = 100$  and  $method = 1$  for the four different  $\lambda$  (figure[3]);
- the coverage rate of  $\mu$ : given  $\lambda = 4$  and  $method = 1$  for the five different  $n$  (figure[4]);
- the coverage rate of  $\mu$ : given  $\lambda = 4$  and  $n = 10$  for the six different methods (figure[5]);

I also compute the coverage rate of  $\lambda$  in the same conditions, the results are similar. By checking the figures, I find the coverage rates become stable when  $\sqrt{M} = 100$ , which means  $M = 10000$ . So, I choose M to be 10000 for all the cases.

### 3 Results

The **coverage rate**, **median CI width** and the **out of bound probability** of all the combinations of experiments ( $\lambda \in \{2, 4, 8, 12\}$ ,  $n \in \{10, 25, 50, 100, 500\}$ , and  $method \in \{1, 2, 3, 4, 5, 6\}$ ) are shown in table[1] –table[6] in the Appendix.

## 4 Discussion and Conclusions

### 4.1 Performance of Methods When $n$ Different

**For confidence intervals of  $\mu$ :**

When the sample size  $n$  small, the actual coverage rate of method 1, 3, and 6 are small, (i.e. when  $n = 10$ ,  $\lambda = 2$ :  $cv1 = 0.7941$ ,  $cv3 = 0.7339$ ,  $cv6 = 0.7042$ ) while the coverage rate of method 2 are very close to 0.95 (i.e. when  $n = 10$ ,  $\lambda = 2$ :  $cv2 = 0.9324$ ). However, the median width of method 2 is too large compared to the other methods (i.e. when  $n = 10$ ,  $\lambda = 2$ :  $med.wid2 = 45.7270$ ,  $med.wid1 = 7.4000$ ). I guess the coverage rate of method 2 is close to 0.95 because the CI computed by method 2 is too wide so that the true  $\mu$  falls into most of the CI, while the coverage rate of method 1, 3, and 6 is small means the CI computed by these methods are not wide enough to capture enough information. So, by the coverage rate and median width trade-off, I prefer to pick method 4 and method 5 (i.e. when  $n = 10$ ,  $\lambda = 2$ :  $cv4 = 0.8583$ ,  $cv5 = 0.8860$ ,  $med.wid4 = 8.4691$ ,  $med.wid5 = 13.4934$ ). Besides, the out of bound probability of method 4 and method 5 is always 0, which also means method 4 and method 5 are better.

When the sample size  $n$  large, the differences between the different methods become less. when  $n = 500$ , the differences of the first five methods are nearly negligible (The coverage rate of method 6 when  $n = 500$  is a little small than the first five methods) and all of the coverage rate of the six methods are approaching to 0.95. So, the first five methods (even method 6) are applicable. In this case, I prefer method 1 because it's the fastest method.

**For confidence intervals of  $\lambda$ :**

Firstly, we should notice a fact that the confidence interval computed by method 4 and method 5 are the same give the same seed. (It has been discussed in section 2.5.) Also, the results of method 1 and method 2 perform very similar (nearly the same) by checking table [2], [4], and [6].

When the sample size  $n$  small, method 1, 2, 4, and 5 have the similar median CI width, method 3

and method 6 have a much larger median CI width than the front four methods. (i.e. when  $n = 10$ ,  $\lambda = 2$ :  $med.wid1 = 4.1874$ ,  $med.wid2 = 4.276$ ,  $med.wid4\&5 = 3.9031$ ,  $med.wid3 = 7.5531$ ,  $med.wid6 = 17.6629$ ). On the other hand, the out of bound probability of method 3 and method 6 are very large while the other four methods always have 0 probability (i.e. when  $n = 10$ ,  $\lambda = 2$ :  $P(out\ of\ bound\ of\ method\ 3) = 1$ ,  $P(out\ of\ bound\ of\ method\ 6) = 1$ ). In the table [2], I notice that the coverage rate of method 4 and 5 are closer to 0.95 than method 1 and 2, while the confidence intervals are narrower. In this case, I recommend to choose method 4 or method 5. (Method 4 is a little faster).

When the sample size  $n$  large, the first five methods outperform the method 6 (i.e. when  $n = 500$ ,  $\lambda = 2$ : the median width of the first fives methods are about 0.497, while  $med.wid6 = 1.3649$ ). Besides, the out of bound probability of method 6 when  $n = 100$ ,  $\lambda = 2$  is still big (i.e.  $P=0.1697$ ). As a result, first five methods are applicable. In this case, I prefer method 1 because it's the fastest.

## 4.2 Performance of Methods When $\lambda$ Different

The change of  $\lambda$  will not affect the performance of the confidence interval for  $\lambda$ , however, it does have influence on the confidence interval for  $\mu$ . It can be found in table [1], [3] and [5] that the performance of the confidence interval for  $\mu$  becomes better when  $\lambda$  increase (i.e. the coverage rate is closer to 0.95, the out of bound probability is smaller).

By checking table [1], I found the coverage rate of method 2 and method 5 for  $\mu$  is closer to 0.95 than the other four methods when  $\lambda$  is large and  $n$  is small (e.g. when  $n = 10$ ,  $\lambda = 12$ :  $cv2 = 0.9282$ ,  $cv5 = 0.9376$ , the other methods are all less than 0.9). At the same time, the median CI width of these two methods are not too large compared to the other four methods, the out of bound probability is 0 for these two methods. So, I would like to pick method 2 or method 5 for conducting CI of  $\mu$  when  $\lambda$  large and  $n$  small.

The method 4 is also applicable since the out of bound probability is 0 and the coverage rate is higher than method 1, 3 and 6, though it's not as good as method 2 and method 5.



### 4.3 Conclusions

For the confidence interval of  $\mu$ , the following methods have good performance (\* The method in bracket means: applicable but not as good as the other ones.):

	$n$ small	$n$ large
$\lambda$ small	method 4 5	method 1 2 3 4 5 (6)
$\lambda$ large	method 2 (4) 5	method 1 2 3 4 5 (6)

For the confidence interval of  $\lambda$ , the following methods have good performance:

$n$ small	$n$ large
method 4 5	method 1 2 3 4 5

Over all, my favorite method is method 5 since it has a good performance in any condition. However, I recommend to use method 1 when the sample size is large since it's much faster than the other ones.

## 5 Appendix

### 5.1 R Functions of Method 1-6

---

```
##### Method1 MLE & Wald Theory #####
mle_mu <- function(x) return( mean(x) )
mle_la <- function(x) return( 1 / ( mean(1/x) - 1/mean(x) ) )

ci_mu_1 <- function(x){
  sigma <- sqrt( (mle_mu(x))^3/(length(x)*mle_la(x)) )
  lb <- mle_mu(x) - 1.96*sigma
  ub <- mle_mu(x) + 1.96*sigma
  return(c(lb,ub))
}

ci_la_1 <- function(x){
  sigma <- sqrt( (2*(mle_la(x))^2)/length(x) )
  lb <- mle_la(x) - 1.96*sigma
  ub <- mle_la(x) + 1.96*sigma
  return(c(lb,ub))
}

##### Method2 MLE & Inverse LRT #####

## lambda
LR_la <- function(la,x){
  n <- length(x)
  return((sum(1/x) - (n/mle_mu(x)) )*(la - mle_la(x)) - n*log((la)/mle_la(x)))
}

ci_la_2 <- function(x){
  me <- mle_la(x)
  ci <- ci_la_1(x)
```

```

n <- length(x)
if (n <= 20) b <- 5*(me-ci[1])
if (n>20 & n <=50) b <- 3*(me-ci[1])
if (n>50) b <- 2*(me-ci[1])
st <- max(me-b,0.001)
ed <- me+b
XX <- seq(st,ed,0.001)
LR <- LR_la( XX, x )
C <- qchisq(0.95,1)
index1 <- which(LR < C)[1]
index2 <- which(LR < C)[length(which(LR < C))]
return(c(XX[index1],XX[index2]))
}

```

```

##mu
til_la <- function(mu,x){
  Y <- (mle_mu(x))/(mu^2) - (2/mu) + mean(1/x)
  return(1/Y)
}

```

```

LR_mu <- function(mu,x){
  n <- length(x)
  L1 <- (n/2)*log(til_la(mu,x)) - ((til_la(mu,x))/(2*mu^2))*sum(x) +
    (n*til_la(mu,x))/mu - (til_la(mu,x)/2)*sum(1/x)
  L2 <- (n/2)*log(mle_la(x)) + (n*mle_la(x))/(2*mle_mu(x)) -
    (mle_la(x)/2)*sum(1/x)
  return((-2)*(L1-L2))
}

```

```

ci_mu_2 <- function(x){
  me <- mle_mu(x)
  ci <- ci_mu_1(x)
  n <- length(x)
  if (n <= 20) b <- 15*(me-ci[1])

```

```

if (n>20 & n <=50) b <- 10*(me-ci[1])
if (n>50 & n<=100) b <- 5*(me-ci[1])
if (n>100 & n<=200) b <- 3*(me-ci[1])
if(n>200) b <- 2*(me-ci[1])
st <- max(me-b,0.00001)
ed <- me+b
XX <- seq(st,ed,0.001)
LR <- LR_mu( XX, x )
C <- qchisq(0.95,1)
index1 <- which(LR < C)[1]
index2 <- which(LR < C)[length(which(LR < C))]
return(c(XX[index1],XX[index2]))
}

```

##### Method3 MLE & Parametric bootstrap 1 #####

```

ci_boot_mu_1 <- function(x,N){
  library(statmod)
  mu <- mle_mu(x)
  la <- mle_la(x)
  n <- length(x)
  X <- matrix(rinvgauss(n*N,mu,la),ncol = N)
  mu1 <- apply(X,2,mle_mu)
  mu1 <- sort(mu1)
  qmu1 <- mu1[ floor( (N+1)*(1-(0.05/2))) ]
  qmu2 <- mu1[ floor( (N+1)*(0.05/2) ) ]
  lb <- 2*mu-qmu1
  ub <- 2*mu-qmu2
  return(c(lb,ub))
}

```

```

ci_boot_la_1 <- function(x,N){
  library(statmod)

```

```

mu <- mle_mu(x)
la <- mle_la(x)
n <- length(x)
X <- matrix(rinvgauss(n*N,mu,la),ncol = N)
la1 <- apply(X,2,mle_la)
la1 <- sort(la1)
qla1 <- la1[ floor( (N+1)*(1-(0.05/2))) ]
qla2 <- la1[ floor( (N+1)*(0.05/2) ) ]
lb <- 2*la-qla1
ub <- 2*la-qla2
return(c(lb,ub))
}

```

##### Method4 MLE & Parametric bootstrap 2 #####

```

ci_boot_mu_2 <- function(x,N){
  library(statmod)
  mu <- mle_mu(x)
  la <- mle_la(x)
  n <- length(x)
  X <- matrix(rinvgauss(n*N,mu,la),ncol = N)
  mu1 <- apply(X,2,mle_mu)
  mu1 <- sort(mu1)
  qmu1 <- mu1[ floor( (N+1)*(1-(0.05/2))) ]
  qmu2 <- mu1[ floor( (N+1)*(0.05/2) ) ]
  lb <- mu^2/qmu1
  ub <- mu^2/qmu2
  return(c(lb,ub))
}

```

```

ci_boot_la_2 <- function(x,N){
  library(statmod)
  mu <- mle_mu(x)
  la <- mle_la(x)

```

```

n <- length(x)
X <- matrix(rinvgauss(n*N,mu,la),ncol = N)
la1 <- apply(X,2,mle_la)
la1 <- sort(la1)
qla1 <- la1[ floor( (N+1)*(1-(0.05/2))) ]
qla2 <- la1[ floor( (N+1)*(0.05/2) ) ]
lb <- la^2/qla1
ub <- la^2/qla2
return(c(lb,ub))
}

```

##### Method5 MLE & Parametric bootstrap 3 #####

```

Vmu <- function(x){
  n <- length(x)
  la <- mle_la(x)
  mu <- mle_mu(x)
  return((mu^3)/(n*la))
}

```

```

Vla <- function(x){
  n <- length(x)
  la <- mle_la(x)
  return((2*la^2)/n)
}

```

```

ci_boot_mu_3 <- function(x,N){
  library(statmod)
  mu <- mle_mu(x)
  la <- mle_la(x)
  n <- length(x)
  X <- matrix(rinvgauss(n*N,mu,la),ncol = N)
  mu1 <- apply(X,2,mle_mu) - mu
  V1 <- apply(X,2,Vmu)
}

```

```

mu11 <- sort(mu1/sqrt(V1))
qmu1 <- mu11[ floor( (N+1)*(1-(0.05/2))) ]
qmu2 <- mu11[ floor( (N+1)*(0.05/2) ) ]
lb <- mu-sqrt(Vmu(x))*qmu1
ub <- mu-sqrt(Vmu(x))*qmu2
return(c(lb,ub))
}

```

```

ci_boot_la_3 <- function(x,N){
  library(statmod)
  mu <- mle_mu(x)
  la <- mle_la(x)
  n <- length(x)
  X <- matrix(rinvgauss(n*N,mu,la),ncol = N)
  la1 <- apply(X,2,mle_la) - la
  V1 <- apply(X,2,Vla)
  la11 <- sort(la1/sqrt(V1))
  qla1 <- la11[ floor( (N+1)*(1-(0.05/2))) ]
  qla2 <- la11[ floor( (N+1)*(0.05/2) ) ]
  lb <- la-sqrt(Vla(x))*qla1
  ub <- la-sqrt(Vla(x))*qla2
  return(c(lb,ub))
}

```

##### Method6 MME & Parametric bootstrap 4 #####

```

mme_mu <- function(x) return(mean(x))
mme_la <- function(x) return( ((mean(x))^3)/(mean(x^2)-(mean(x))^2) )

```

```

ci_boot_mu_4 <- function(x,N){
  library(statmod)
  mu <- mme_mu(x)
  la <- mme_la(x)

```

```

n <- length(x)
X <- matrix(rinvgauss(n*N,mu,la),ncol = N)
mu1 <- apply(X,2,mme_mu)
mu1 <- sort(mu1)
qmu1 <- mu1[ floor( (N+1)*(1-(0.05/2))) ]
qmu2 <- mu1[ floor( (N+1)*(0.05/2) ) ]
lb <- 2*mu-qmu1
ub <- 2*mu-qmu2
return(c(lb,ub))
}

```

```

ci_boot_la_4 <- function(x,N){
  library(statmod)
  mu <- mme_mu(x)
  la <- mme_la(x)
  n <- length(x)
  X <- matrix(rinvgauss(n*N,mu,la),ncol = N)
  la1 <- apply(X,2,mme_la)
  la1 <- sort(la1)
  qla1 <- la1[ floor( (N+1)*(1-(0.05/2))) ]
  qla2 <- la1[ floor( (N+1)*(0.05/2) ) ]
  lb <- 2*la-qla1
  ub <- 2*la-qla2
  return(c(lb,ub))
}

```

---

## 5.2 R Functions of 3 criteria

---

```

criteria <- function(M,n = 10, mu = 5,lambda = 2,

```



```

para = c("mu","lambda"), method = c(1,2,3,4,5,6)){
  if (method == 1){
    ci_mu <- function(x) ci_mu_1(x)
    ci_la <- function(x) ci_la_1(x)
  }
  if (method == 2){
    ci_mu <- function(x) ci_mu_2(x)
    ci_la <- function(x) ci_la_2(x)
  }
  if (method == 3){
    ci_mu <- function(x) ci_boot_mu_1(x,2000)
    ci_la <- function(x) ci_boot_la_1(x,2000)
  }
  if (method == 4){
    ci_mu <- function(x) ci_boot_mu_2(x,2000)
    ci_la <- function(x) ci_boot_la_2(x,2000)
  }
  if (method == 5){
    ci_mu <- function(x) ci_boot_mu_3(x,2000)
    ci_la <- function(x) ci_boot_la_3(x,2000)
  }
  if (method == 6){
    ci_mu <- function(x) ci_boot_mu_4(x,2000)
    ci_la <- function(x) ci_boot_la_4(x,2000)
  }
  set.seed(520520)
  library(statmod)
  if (para == "mu"){
    IG <- matrix(rinvgauss(M*n,mu,lambda), ncol=M)
    CImu <- apply(IG,2,ci_mu)
    ID <- (CImu[1,]<mu)&(CImu[2,]>mu)
    CI <- t(CImu)
  }
}

```

```


if (para == "lambda"){
  IG <- matrix(rinvgauss(M*n,mu,lambda), ncol=M)
  CI1a <- apply(IG,2,ci_la)
  ID <- (CI1a[1,<lambda]&(CI1a[2,>lambda)
  CI <- t(CI1a)
}

rate <- mean(ID)
width <- CI[,2]-CI[,1]
med_width <- median(width)
prob <- mean(CI[,1]<0)
result <- list(rate = rate,medianWidth = med_width, Prob = prob)
return(result)
}

```

---

### 5.3 Some Tables and Figures



**Figure 1:** Choose bootstrap  $N$  using the CI width of  $\mu$




Figure 2: Choose bootstrap N using the CI width of  $\lambda$




Figure 3




Figure 4




Figure 5

"lambda"	"n"	"method1"	"method2"	"method3"	"method4"	"method5"	"method6"
2	10	0.7941	0.9324	0.7339	0.8583	0.886	0.7042
2	25	0.8688	0.9409	0.829	0.9074	0.9058	0.8061
2	50	0.9061	0.9479	0.878	0.9336	0.9272	0.8601
2	100	0.9213	0.946	0.905	0.9385	0.9359	0.8924
2	500	0.9462	0.951	0.9421	0.9491	0.9489	0.9359
4	10	0.8468	0.9309	0.7984	0.8812	0.9185	0.7741
4	25	0.8928	0.9431	0.8678	0.9191	0.929	0.8498
4	50	0.9243	0.9497	0.9052	0.939	0.9419	0.8928
4	100	0.9342	0.9428	0.9231	0.9372	0.9399	0.914
4	500	0.9483	0.9494	0.946	0.9487	0.9476	0.9442
8	10	0.872	0.9283	0.8461	0.8925	0.9328	0.8293
8	25	0.9137	0.9416	0.8984	0.9281	0.9382	0.8862
8	50	0.9363	0.9516	0.9247	0.9444	0.9491	0.9147
8	100	0.9401	0.9494	0.9348	0.9446	0.9472	0.9285
8	500	0.9475	0.9483	0.9458	0.9478	0.9476	0.9445
12	10	0.881	0.9282	0.863	0.8946	0.9376	0.8464
12	25	0.9191	0.9422	0.9056	0.9267	0.941	0.8957
12	50	0.9386	0.9478	0.9311	0.9426	0.9486	0.9249
12	100	0.9416	0.9491	0.9346	0.945	0.9499	0.9322
12	500	0.9472	0.9471	0.9466	0.9473	0.9483	0.9464

**Table 1:** Coverage Rate for  $\mu$

"lambda"	"n"	"method1"	"method2"	"method3"	"method4"	"method5"	"method6"
2	10	0.9706	0.9292	0.9026	0.9517	0.9517	0.9607
2	25	0.9598	0.9444	0.9193	0.9497	0.9497	0.9354
2	50	0.9554	0.9492	0.9288	0.9501	0.9501	0.9339
2	100	0.9507	0.9459	0.9404	0.948	0.948	0.9352
2	500	0.9529	0.9521	0.9505	0.952	0.952	0.9455
4	10	0.9698	0.9311	0.9023	0.951	0.951	0.9418
4	25	0.9588	0.9432	0.9191	0.9485	0.9485	0.9306
4	50	0.9567	0.9485	0.9302	0.9518	0.9518	0.9364
4	100	0.951	0.9456	0.9408	0.9481	0.9481	0.9389
4	500	0.9543	0.9517	0.9502	0.9523	0.9523	0.947
8	10	0.9699	0.9289	0.9019	0.9503	0.9503	0.9262
8	25	0.957	0.945	0.9192	0.9487	0.9487	0.9268
8	50	0.9547	0.9494	0.9305	0.9512	0.9512	0.9369
8	100	0.9517	0.9452	0.9406	0.9477	0.9477	0.9423
8	500	0.9538	0.9515	0.9498	0.9525	0.9525	0.9482
12	10	0.969	0.9272	0.9025	0.9511	0.9511	0.9218
12	25	0.9597	0.9449	0.9193	0.9494	0.9494	0.9268
12	50	0.9555	0.9495	0.9291	0.9526	0.9526	0.9372
12	100	0.9513	0.9458	0.9415	0.9477	0.9477	0.9431
12	500	0.9532	0.9521	0.9497	0.9525	0.9525	0.9492

**Table 2:** Coverage Rate for  $\lambda$

"lambda"	"n"	"method1"	"method2"	"method3"	"method4"	"method5"	"method6"
2	10	7.4001	45.727	7.199	8.4691	13.4934	5.7284
2	25	5.5346	9.1635	5.4744	5.9553	7.4617	4.7185
2	50	4.1392	5.1925	4.1	4.2945	4.856	3.7141
2	100	3.0102	3.3565	2.9955	3.0705	3.2708	2.8149
2	500	1.381	1.41	1.3768	1.3845	1.403	1.3548
4	10	5.7566	11.9495	5.6848	6.2592	8.9142	4.8243
4	25	4.063	5.222	4.033	4.2235	4.9223	3.6593
4	50	2.9679	3.344	2.9537	3.0301	3.2788	2.787
4	100	2.1491	2.276	2.1431	2.1721	2.2606	2.0653
4	500	0.976	0.986	0.975	0.9778	0.9861	0.966
8	10	4.2347	6.1545	4.198	4.4093	5.8501	3.8137
8	25	2.9183	3.354	2.898	2.9744	3.3387	2.7506
8	50	2.1296	2.28	2.1281	2.1552	2.2813	2.045
8	100	1.5265	1.578	1.5237	1.5339	1.5799	1.496
8	500	0.691	0.695	0.6901	0.6912	0.6953	0.6866
12	10	3.5064	4.6245	3.488	3.6108	4.6698	3.2474
12	25	2.3988	2.663	2.3907	2.4301	2.6898	2.2949
12	50	1.7477	1.84	1.7438	1.7582	1.8499	1.6899
12	100	1.2484	1.28	1.2462	1.2512	1.2845	1.2299
12	500	0.5641	0.566	0.5635	0.564	0.5672	0.5614

**Table 3:** Median Width for  $\mu$



"lambda"	"n"	"method1"	"method2"	"method3"	"method4"	"method5"	"method6"
2	10	4.1874	4.276	7.5531	3.9031	3.9031	17.6629
2	25	2.3757	2.395	2.9569	2.3133	2.3133	7.1005
2	50	1.6176	1.6235	1.7996	1.5981	1.5981	4.3961
2	100	1.1288	1.13	1.1883	1.1208	1.1208	2.9763
2	500	0.4973	0.497	0.5024	0.4966	0.4966	1.3649
4	10	8.3938	8.573	15.1829	7.8323	7.8323	26.827
4	25	4.7515	4.791	5.9095	4.6316	4.6316	10.8836
4	50	3.2364	3.249	3.6001	3.197	3.197	6.832
4	100	2.2568	2.261	2.3741	2.2398	2.2398	4.6577
4	500	0.9948	0.994	1.005	0.9936	0.9936	2.0995
8	10	16.7491	17.1075	30.2689	15.5985	15.5985	43.4247
8	25	9.4969	9.577	11.8034	9.2449	9.2449	17.6828
8	50	6.4756	6.502	7.1915	6.394	6.394	11.1394
8	100	4.514	4.523	4.7476	4.4786	4.4786	7.5286
8	500	1.9894	1.989	2.0105	1.9872	1.9872	3.3332
12	10	25.0642	25.601	45.2767	23.4108	23.4108	59.2827
12	25	14.2387	14.359	17.7368	13.8751	13.8751	24.1096
12	50	9.7206	9.761	10.7984	9.6105	9.6105	15.1377
12	100	6.7733	6.7865	7.1215	6.7167	6.7167	10.2105
12	500	2.984	2.984	3.0144	2.9794	2.9794	4.4586

**Table 4:** Median Width for  $\lambda$

"lambda"	"n"	"method1"	"method2"	"method3"	"method4"	"method5"	"method6"
2	10	0.296	0	0.55	0	0	0.2654
2	25	0.0041	0	0.0649	0	0	0.0422
2	50	0	0	0	0	0	0.001
2	100	0	0	0	0	0	0
2	500	0	0	0	0	0	0
4	10	0.038	0	0.176	0	0	0.0776
4	25	0	0	0	0	0	0.0014
4	50	0	0	0	0	0	0
4	100	0	0	0	0	0	0
4	500	0	0	0	0	0	0
8	10	2e-04	0	0.0074	0	0	0.0076
8	25	0	0	0	0	0	0
8	50	0	0	0	0	0	0
8	100	0	0	0	0	0	0
8	500	0	0	0	0	0	0
12	10	0	0	1e-04	0	0	5e-04
12	25	0	0	0	0	0	0
12	50	0	0	0	0	0	0
12	100	0	0	0	0	0	0
12	500	0	0	0	0	0	0

**Table 5:** Out of Bound Probability for  $\mu$

"lambda"	"n"	"method1"	"method2"	"method3"	"method4"	"method5"	"method6"
2	10	0	0	1	0	0	1
2	25	0	0	0.6154	0	0	1
2	50	0	0	0	0	0	0.9607
2	100	0	0	0	0	0	0.1697
2	500	0	0	0	0	0	0
4	10	0	0	1	0	0	1
4	25	0	0	0.6108	0	0	1
4	50	0	0	0	0	0	0.5671
4	100	0	0	0	0	0	0.0027
4	500	0	0	0	0	0	0
8	10	0	0	1	0	0	1
8	25	0	0	0.6097	0	0	1
8	50	0	0	0	0	0	0.0482
8	100	0	0	0	0	0	0
8	500	0	0	0	0	0	0
12	10	0	0	1	0	0	1
12	25	0	0	0.6088	0	0	1
12	50	0	0	0	0	0	0.0027
12	100	0	0	0	0	0	0
12	500	0	0	0	0	0	0

**Table 6:** Out of Bound Probability for  $\lambda$