

# A Regularity Condition of 3d Axisymmetric Navier-Stokes Equations

Xinghong Pan<sup>1</sup> 

Received: 20 March 2016 / Accepted: 9 March 2017 / Published online: 6 April 2017  
© Springer Science+Business Media Dordrecht 2017

**Abstract** In this paper, we study the regularity of 3d axisymmetric Navier-Stokes equations under a prior point assumption on  $v^r$  or  $v^z$ . That is, the weak solution of the 3d axisymmetric Navier-Stokes equations  $v$  is smooth if

$$rv^r \geq -1; \quad \text{or} \quad r|v^r(t, x)| \leq Cr^\alpha, \quad \alpha \in (0, 1]; \quad \text{or} \quad r|v^z(t, x)| \leq Cr^\beta, \quad \beta \in [0, 1];$$

where  $r$  is the distance from the point  $x$  to the symmetric axis.

**Keywords** Axisymmetric · Navier-Stokes equations · Regularity criteria

**Mathematics Subject Classification** 35Q30 · 76N10

## 1 Introduction

In the cylindrical coordinates  $(r, \theta, z)$ , we have  $x = (x_1, x_2, x_3) = (r \cos \theta, r \sin \theta, z)$  and the axisymmetric solution of the incompressible Navier-Stokes equations is given as

$$v = v^r(r, z, t)e_r + v^\theta(r, z, t)e_\theta + v^z(r, z, t)e_z,$$

where the basis vectors  $e_r, e_\theta, e_z$  are

$$e_r = \left( \frac{x_1}{r}, \frac{x_2}{r}, 0 \right), \quad e_\theta = \left( -\frac{x_2}{r}, \frac{x_1}{r}, 0 \right), \quad e_z = (0, 0, 1).$$

---

✉ X. Pan  
[math.scrat@gmail.com](mailto:math.scrat@gmail.com)

<sup>1</sup> Department of Mathematics, Nanjing University of Aeronautics and Astronautics, Nanjing 211106, China

The components  $v^r$ ,  $v^\theta$ ,  $v^z$ , independent of  $\theta$ , satisfy

$$\begin{cases} \partial_t v^r + (b \cdot \nabla) v^r - \frac{(v^\theta)^2}{r} + \partial_r p = \left( \Delta - \frac{1}{r^2} \right) v^r, \\ \partial_t v^\theta + (b \cdot \nabla) v^\theta + \frac{v^\theta v^r}{r} = \left( \Delta - \frac{1}{r^2} \right) v^\theta, \\ \partial_t v^z + (v \cdot \nabla) v^z + \partial_z p = \Delta v^z, \\ b = v^r e_r + v^z e_z, \quad \nabla \cdot b = \partial_r v^r + \frac{v^r}{r} + \partial_z v^z = 0. \end{cases} \quad (1.1)$$

The finite energy smooth solutions of the Navier-Stokes equations satisfy the following energy identity

$$\|v(t)\|_{L^2}^2 + 2 \int_0^t \|\nabla v(s)\|_{L^2}^2 ds = \|v_0\|_{L^2}^2 < +\infty. \quad (1.2)$$

For the axisymmetric velocity  $v$ , we can also compute the vorticity  $w = \nabla \times v = w^r e_r + w^\theta e_\theta + w^z e_z$  as follows

$$w^r = -\partial_z v^\theta, \quad w^\theta = \partial_z v^r - \partial_r v^z, \quad w^z = \left( \partial_r + \frac{1}{r} \right) v^\theta.$$

And  $(w^r, w^\theta, w^z)$  satisfies

$$\begin{cases} \partial_t w^r + (b \cdot \nabla) w^r - \left( \Delta - \frac{1}{r^2} \right) w^r - (w^r \partial_r + w^z \partial_z) v^r = 0, \\ \partial_t w^\theta + (b \cdot \nabla) w^\theta - \left( \Delta - \frac{1}{r^2} \right) w^\theta - 2 \frac{v^\theta}{r} \partial_z v^\theta - w^\theta \frac{v^r}{r} = 0, \\ \partial_t w^z + (b \cdot \nabla) w^z - \Delta w^z - (w^r \partial_r + w^z \partial_z) v^z = 0. \end{cases} \quad (1.3)$$

Let us recall some results on the study of the axisymmetric Navier-Stokes equations. When the swirl  $v^\theta$  is vanishing, Ladyzhenskaya [12] and Ukhovskii-Yudovich [18] independently proved that finite energy weak solutions are regular for all time, see also Leonardi-Malek-Necas-Pokorný [16]. If the swirl  $v^\theta$  is non-trivial, global in-time regularity of solutions is still open. But recently, tremendous efforts and progress have been made on the regularity of solutions to the axisymmetric Navier-Stokes equations, see [1, 6–8] etc. In [3] and [4], Chen-Strain-Yau-Tsai proved that the suitable weak solution is regular if it satisfies  $r|v| \leq C < \infty$ . The same result was obtained by Koch-Nadirashvili-Seregin-Sverak in [10] by using a different method. Lei and Zhang in [13] extend their results under a more general critical assumption on the drift term  $b \in L^\infty([0, T], BMO^{-1})$ .

Also, considering on one velocity component, it has been shown that in [11] the axisymmetric solution is smooth in  $(0, T) \times \mathbb{R}^3$  when  $v^r \in L_t^s L_x^q$  with  $3/q + 2/s \leq 1$ . Neustupa and Pokorný [17] proved the regularity of one component (either  $v^\theta$  or  $v^r$ ) implies regularity of the other components of the solutions. Chae-Lee [2] proved regularity assuming a zero-dimensional integral norm on  $w^\theta$ :  $w^\theta \in L_t^s L_x^q$  with  $3/q + 2/s \leq 2$ . Also regularity results come from the work of Jiu-Xin [8] under the assumption that another zero-dimensional scaled norms  $\int_{\mathbb{R}^3} (R^{-1}|w^\theta|^2 + R^{-3}|v^\theta|^2) dz$  is sufficiently small for  $R > 0$ , small enough. See more refined results in [9] and the work of Zhang-Zhang [20].

Most recently, Chen-Fang-Zhang in [5] proved that if  $rv^\theta$  satisfies  $r|v^\theta| \leq Cr^\alpha$ ,  $\alpha > 0$ , then  $v$  is regular without any other a priori assumptions. Later, Lei-Zhang in [14] improved their result by assuming  $r|v^\theta| \leq C|\ln r|^{-2}$  for small  $r$ . Also Wei in [19] improved the  $\log$  power from  $-2$  to  $-\frac{3}{2}$ .

Our result is a complement of theirs. We prove the regularity of the solution by assuming Hölder continuity on  $rv^r$  and  $rv^z$ . Below is the main theorem.

**Theorem 1.1** *Let  $v$  is an axisymmetric weak solution of the Navier-Stokes equations in  $(0, T) \times \mathbb{R}^3$  with the axisymmetric initial data  $v_0 \in H^2(\mathbb{R}^3)$  and  $\nabla \cdot v_0 = 0$ . If  $rv^r$  or  $rv^z$  satisfies*

$$rv^r \geq -1; \quad \text{or} \quad r|v^r(t, x)| \leq Cr^\alpha, \quad \alpha \in (0, 1]; \quad \text{or} \quad r|v^z(t, x)| \leq Cr^\beta, \quad \beta \in [0, 1];$$

*Then  $v$  is smooth in  $(0, T] \times \mathbb{R}^3$ .*

**Remark 1.1** The case that  $r|v^z| \leq C$ , where  $\beta = 0$  implied the regularity of the solution has already been proved in proposition 4.2 of [15].

**Remark 1.2** Now we can not prove the regularity of the solution under the assumption  $rv^r < -1$ .

## 2 Proof of the Regularity Criteria

First, we state two lemmas which will be very useful in the proof of the theorem.

**Lemma 2.1** *Assume  $v$  is a smooth axisymmetric solution of (1.1) on  $(0, T)$ , then  $v$  can be stated as follow.*

$$v = v^\theta e_\theta + \nabla \times (\psi e_\theta) = -\partial_z \psi e_r + v^\theta e_\theta + \frac{\partial_r(r\psi)}{r} e_z, \quad (2.1)$$

where  $\psi$  is the angular component of the stream function of  $v$ .

Next we give a general Sobolev-Hardy inequality whose proof can be found in Lemma 2.4 of [5]. We omit the detail.

**Lemma 2.2** *Set  $\mathbb{R}^n = \mathbb{R}^k \times \mathbb{R}^{n-k}$  with  $2 \leq k \leq n$ , and write  $x = (x', z) \in \mathbb{R}^k \times \mathbb{R}^{n-k}$ . For  $1 < q < n$ ,  $0 \leq s \leq q$  and  $s < k$ , let  $q_* \in [q, \frac{q(n-s)}{n-q}]$ . Then there exists a positive constant  $C = C(s, q, n, k)$  such that for all  $f \in C_0^\infty(\mathbb{R}^n)$ ,*

$$\int_{\mathbb{R}^n} \frac{|f|^{q_*}}{|x'|^s} dx \leq C \|f\|_q^{\frac{n-s}{q_*} - \frac{n}{q} + 1} \|\nabla f\|_q^{\frac{n}{q} - \frac{n-s}{q_*}}.$$

*In particular, we pick  $n = 3$ ,  $k = 2$ ,  $q = 2$ ,  $q_* \in [2, 2(3-s)]$ , and assume  $0 \leq s < 2$ ,  $r = \sqrt{x_1^2 + x_2^2}$ . Then there exists a positive constant  $C(q_*, s)$  such that for all  $f \in C_0^\infty(\mathbb{R}^n)$ ,*

$$\left\| \frac{f}{r^{\frac{s}{q_*}}} \right\|_{q_*} \leq C \|f\|_2^{\frac{3-s}{q_*} - \frac{1}{2}} \|\nabla f\|_2^{\frac{3}{2} - \frac{3-s}{q_*}}. \quad (2.2)$$

*Proof of Theorem 1.1*

**Case 1:**  $rv^r \geq -1$  or  $r|v^r(t, x)| \leq Cr^\alpha$ ,  $\alpha \in (0, 1]$ .

From the equation of  $v^\theta$  in (1.1), we have

$$\partial_t v^\theta + (b \cdot \nabla) v^\theta - \Delta v^\theta + \frac{rv^r + 1}{r^2} v^\theta = 0. \quad (2.3)$$

1. If  $rv^r \geq -1$ , the coefficient  $\frac{rv^r + 1}{r^2}$  of  $v^\theta$  is nonnegative, which has a lower bound 0.

2. If  $r|v^r(t, x)| \leq Cr^\alpha$ ,  $\alpha \in (0, 1]$ , using a simple computation, we can get that

$$\frac{rv^r + 1}{r^2} \geq \frac{1 - Cr^\alpha}{r^2} \geq c_0,$$

where  $c_0$  is a fixed constant which may be negative. So at last for *Case 1*, we get

$$\frac{rv^r + 1}{r^2} \geq \min\{0, c_0\} \quad (2.4)$$

From (2.3) and (2.4), by using Maximum principle and Sobolev embedding inequality, we obtain

$$\sup_t \|v^\theta\|_{L^\infty} \leq \|v_0^\theta\|_{L^\infty} \leq C \|v_0\|_{H^2} < \infty,$$

which implies the regularity of the solution.

**Case 2:**  $r|v^z| < Cr^\beta$ , where  $\beta = 0$ .

Actually this case has already been proven in proposition 4.2 of [15], but for completion of our paper, we state it here again. From (2.1), we have

$$v^r = -\partial_z \psi, \quad rv^z = \partial_r(r\psi).$$

Then

$$|\psi| = \frac{|\int_0^r s v^z ds|}{r} \leq C,$$

which indicates that  $\psi$  is a bounded function which can be embedded in  $BMO$ . As  $\psi$  is the angular component of the stream function of  $v$ , we can get that  $b \in L^\infty(BMO^{-1})$ . The result in [13] implies the regularity of  $v$ .

**Case 3:**  $r|v^z| < Cr^\beta$ , where  $\beta \in (0, 1]$ .

We will deal with this case by combining a  $L^4$  estimate of  $v^\theta$  and a  $L^2$  estimate of  $w^\theta$ . From the equation of  $v^\theta$  in (1.1), multiplying it with  $(v^\theta)^3$  and integrating it over  $\mathbb{R}^3$ , we can get

$$\begin{aligned} & \frac{1}{4} \frac{d}{dt} \|v^\theta\|_4^4 + \frac{3}{4} \|\nabla(v^\theta)^2\|_2^2 + \int_{\mathbb{R}^3} \frac{(v^\theta)^4}{r^2} dx \\ &= - \int_{\mathbb{R}^3} \frac{v^r}{r} (v^\theta)^4 dx \\ &\triangleq I. \end{aligned} \quad (2.5)$$

From (2.1), we have

$$v^r = -\partial_z \psi, \quad r v^z = \partial_r (r \psi). \quad (2.6)$$

Then

$$|\psi| = \frac{|\int_0^r s v^z ds|}{r} \leq C r^\beta.$$

Using integration by parts and (2.6), we have

$$\begin{aligned} |I| &= \left| -\int_{\mathbb{R}^3} \frac{v^r}{r} (v^\theta)^4 dx \right| \\ &= \left| \int_{\mathbb{R}^3} \frac{\partial_z \psi}{r} (v^\theta)^4 dx \right| \\ &= \left| -2 \int_{\mathbb{R}^3} \frac{\psi}{r} (v^\theta)^2 \partial_z (v^\theta)^2 dx \right| \\ &\leq \frac{1}{4} \|\nabla (v^\theta)^2\|_2^2 + 4 \int_{\mathbb{R}^3} \frac{(v^\theta)^4}{r^{2-\beta}} dx. \end{aligned} \quad (2.7)$$

In (2.2), let  $f = (v^\theta)^2$ ,  $q_* = 2$  and  $s = 2 - \beta$ , then we have

$$\begin{aligned} 4 \int_{\mathbb{R}^3} \frac{(v^\theta)^4}{r^{2-\beta}} dx &\leq C_\beta \|(v^\theta)^2\|_2^\beta \|\nabla (v^\theta)^2\|_2^{2-\beta} \\ &\leq \frac{1}{4} \|\nabla (v^\theta)^2\|_2^2 + C_\beta \|(v^\theta)^2\|_2^2. \end{aligned} \quad (2.8)$$

Combining (2.7) and (2.8), we have

$$|I| \leq \frac{1}{2} \|\nabla (v^\theta)^2\|_2^2 + C_\beta \|(v^\theta)^2\|_2^2. \quad (2.9)$$

At last, combining (2.5) and (2.9), we have

$$\frac{d}{dt} \|v^\theta\|_4^4 + \|\nabla (v^\theta)^2\|_2^2 + \int_{\mathbb{R}^3} \frac{(v^\theta)^4}{r^2} dx \leq C \|v^\theta\|_4^4.$$

Applying Gronwall's inequality, we have

$$\sup_{t \in [0, T)} \|v^\theta\|_4^4 + \int_0^T \left\| \frac{(v^\theta)^2}{r} \right\|_2^2 dt \leq \|v_0^\theta\|_4^4 e^{CT} < \infty. \quad (2.10)$$

Next we perform a  $L^2$  estimate of  $w^\theta$ . From the  $w^\theta$  equation of (1.3), we have

$$\begin{aligned} &\frac{1}{2} \frac{d}{dt} \|w^\theta\|_{L^2}^2 + \|\nabla w^\theta\|_{L^2}^2 + \int_{\mathbb{R}^3} \frac{(w^\theta)^2}{r^2} dx \\ &= \int_{\mathbb{R}^3} \frac{v^r}{r} (w^\theta)^2 dx + \int_{\mathbb{R}^3} w^\theta \frac{\partial_z (v^\theta)^2}{r} dx. \end{aligned} \quad (2.11)$$

We can deal with the first term on the right hand of (2.11) the same as  $I$  in (2.5). Thus we can get

$$\begin{aligned}
 & \frac{d}{dt} \|w^\theta\|_{L^2}^2 + \|\nabla w^\theta\|_{L^2}^2 + \int_{\mathbb{R}^3} \frac{(w^\theta)^2}{r^2} dx \\
 & \leq C \|w^\theta\|_{L^2}^2 + C \int_{\mathbb{R}^3} w^\theta \frac{\partial_z (v^\theta)^2}{r} dx \\
 & = C \|w^\theta\|_{L^2}^2 - C \int_{\mathbb{R}^3} \partial_z w^\theta \frac{(v^\theta)^2}{r} dx \\
 & \leq C \|w^\theta\|_{L^2}^2 + \frac{1}{2} \|\nabla w^\theta\|_{L^2}^2 + C \left\| \frac{(v^\theta)^2}{r} \right\|_{L^2}^2.
 \end{aligned} \tag{2.12}$$

Again, using Gronwall's inequality, we have

$$\sup_{t \in (0, T)} \|w^\theta\|_{L^2}^2 \leq e^{CT} \left( \|w_0^\theta\|_{L^2}^2 + \int_0^T \left\| \frac{(v_\theta)^2}{r} \right\|_{L^2}^2 dt \right) < \infty.$$

This implies  $w^\theta \in L_t^\infty L_x^2$ . From the regularity criteria in [2] (Theorem 1), we proved the regularity of the solution.  $\square$

**Acknowledgements** I want to express my gratitude to the anonymous referees for their ideas and suggestions in the proof of this work.

## References

1. Burke Loftus, J., Zhang, Q.S.: A priori bounds for the vorticity of axially symmetric solutions to the Navier-Stokes equations. *Adv. Differ. Equ.* **15**(5–6), 531–560 (2010)
2. Chae, D., Lee, J.: On the regularity of the axisymmetric solutions of the Navier-Stokes equations. *Math. Z.* **239**(4), 645–671 (2002)
3. Chen, C.-C., Strain, R.M., Yau, H.-T., Tsai, T.-P.: Lower bound on the blow-up rate of the axisymmetric Navier-Stokes equations. *Int. Math. Res. Not.* **2008**(9), rnn016, 31 pp. (2008)
4. Chen, C.-C., Strain, R.M., Tsai, T.-P., Yau, H.-T.: Lower bounds on the blow-up rate of the axisymmetric Navier-Stokes equations. II. *Commun. Partial Differ. Equ.* **34**(1–3), 203–232 (2009)
5. Chen, H., Fang, D., Zhang, T.: Regularity of 3D axisymmetric Navier-Stokes equations. [arXiv:1505.00905](https://arxiv.org/abs/1505.00905)
6. Hou, T.Y., Li, C.: Dynamic stability of the three-dimensional axisymmetric Navier-Stokes equations with swirl. *Commun. Pure Appl. Math.* **61**(5), 661–697 (2008)
7. Hou, T.Y., Lei, Z., Li, C.: Global regularity of the 3D axisymmetric Navier-Stokes equations with anisotropic data. *Commun. Partial Differ. Equ.* **33**(7–9), 1622–1637 (2008)
8. Jiu, Q., Xin, Z.: Some regularity criteria on suitable weak solutions of the 3-D incompressible axisymmetric Navier-Stokes equations. In: *Lectures on Partial Differential Equations*. New Stud. Adv. Math., vol. 2, pp. 119–139. Int. Press, Somerville (2003)
9. Jiu, Q., Xin, Z.: On Liouville type of theorems to the 3-D incompressible axisymmetric Navier-Stokes equations. [arXiv:1501.02412](https://arxiv.org/abs/1501.02412)
10. Koch, G., Nadirashvili, N., Seregin, G.A., Sverak, V.: Liouville theorems for the Navier-Stokes equations and applications. *Acta Math.* **203**(1), 83–105 (2009)
11. Kubica, A., Pokorný, M., Zajaczkowski, W.: Remarks on regularity criteria for axially symmetric weak solutions to the Navier-Stokes equations. *Math. Methods Appl. Sci.* **35**(3), 360–371 (2012)
12. Ladyzenskaja, O.A.: Unique global solvability of the three-dimensional Cauchy problem for the Navier-Stokes equations in the presence of axial symmetry. *Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI)* **7**, 155–177 (1968) (Russian)

13. Lei, Z., Zhang, Q.S.: A Liouville theorem for the axially-symmetric Navier-Stokes equations. *J. Funct. Anal.* **261**(8), 2323–2345 (2011)
14. Lei, Z., Zhang, Q.S.: Criticality of the axially symmetric Navier-Stokes equations. *Pac. J. Math.* (2017, to appear). [arXiv:1505.02628](https://arxiv.org/abs/1505.02628)
15. Lei, Z., Navas, E.A., Zhang, Q.S.: A priori bound on the velocity in axially symmetric Navier-Stokes equations. *Commun. Math. Phys.* **341**(1), 289–307 (2016)
16. Leonardi, S., Málek, J., Nečas, J., Pokorný, M.: On axially symmetric flows in  $\mathbb{R}^3$ . *Z. Anal. Anwend.* **18**(3), 639–649 (1999)
17. Neustupa, J., Pokorný, M.: An interior regularity criterion for an axially symmetric suitable weak solution to the Navier-Stokes equations. *J. Math. Fluid Mech.* **2**(4), 381–399 (2000)
18. Ukhovskii, M.R., Iudovich, V.I.: Axially symmetric flows of ideal and viscous fluids filling the whole space. *Prikl. Mat. Meh.* **32**, 59–69 (1968) (Russian); translated as *J. Appl. Math. Mech.* **32**, 52–61
19. Wei, D.: Regularity criterion to the axially symmetric Navier-Stokes equations. *J. Math. Anal. Appl.* **435**(1), 402–413 (2016)
20. Zhang, P., Zhang, T.: Global axisymmetric solutions to three-dimensional Navier-Stokes system. *Int. Math. Res. Not.* **3**, 610–642 (2014)