Chapter 9 solutions

CH09-E1

1.

$$(2-i)(3+2i) = 6 + 4i - 3i - 2i^2 = 8 + i.$$

$$\frac{2-i}{3+2i} = \frac{(2-i)(3-2i)}{13} = \frac{4-7i}{13}.$$

2.

$$|1 + i\sqrt{3}| = 2, \qquad 1 + i\sqrt{3} = 2e^{i\pi/3}.$$

3.

$$(1 + i\sqrt{3})^6 = 2^6 e^{i6\pi/3} = 64e^{2\pi i} = 64.$$

4.


$$z^4 = -16 = 16e^{i\pi}.$$

Roots:

$$z_k = 2e^{i(\pi + 2\pi k)/4}, \quad k = 0, 1, 2, 3.$$

The arguments are $\pi/4, 3\pi/4, 5\pi/4, 7\pi/4$, equally spaced on the radius-2 circle.

z⁴ = −16: radius 2, angle spacing π/2



Angles counterclockwise from the positive real axis

k	$\theta_k = (\pi + 2\pi k)/4$	z_k
0	π/4	√2 + i√2
1	3π/4	−√2 + i√2
2	5π/4	−√2 − i√2
3	7π/4	√2 − i√2

$|z_k| = 2$; $4\theta_k = \pi + 2\pi k$, hence $z_k^4 = -16$

Lin Cing https://github.com/xinhi6716/computer-science-self-study

CH09-E2

1.

$$\|u\|^2 = |1+i|^2 + |2-i|^2 = 7,$$
$$\|v\|^2 = 7.$$
$$u^H v = (1-i)(2-i) + (2+i)(1+i).$$

The two terms are $1-3i$ and $1+3i$, so

$$u^H v = 2.$$
$$v^H u = 2.$$

2.

$$u^H v = \overline{v^H u} = \overline{2} = 2.$$

3. The inner product is 2, not zero, so the vectors are not orthogonal.

4.

$$q = \frac{1}{\sqrt{7}}(1+i, 2-i)^T,$$
$$q^H q = \frac{1}{7} u^H u = 1.$$

Lin Cing
https://github.com/xinhi6716/computer-science-self-study

CH09-E3

1.

$$S_{12} = i, \qquad S_{21} = -i = \overline{i},$$

Both diagonal entries are real, and the off-diagonal entries are conjugates, proving $S^H = S$.

2.

$$p_S(\lambda) = (2-\lambda)^2 - 1 = (\lambda - 1)(\lambda - 3).$$

Both eigenvalues, 1 and 3, are real.

3. For $\lambda = 3$,

$$-z_1 + iz_2 = 0 \Rightarrow z_1 = iz_2.$$

$$q_1 = \frac{1}{\sqrt{2}}(i, 1)^T.$$

For $\lambda = 1$,

$$z_1 + iz_2 = 0 \Rightarrow z_1 = -iz_2.$$

$$q_2 = \frac{1}{\sqrt{2}}(-i, 1)^T.$$

4.

$$Q = \frac{1}{\sqrt{2}} \begin{bmatrix} i & -i \\ 1 & 1 \end{bmatrix}.$$

$$Q^H Q = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}.$$

Therefore

$$S = Q \begin{bmatrix} 3 & 0 \\ 0 & 1 \end{bmatrix} Q^H.$$

Lin Cing https://github.com/xinhi6716/computer-science-self-study

CH09-E4

1.

$$(A^H)^H = A.$$

For a product,

$$(AB)^H_{ij} = \overline{\sum_k A_{jk}B_{ki}} = \sum_k \overline{B_{ki}A_{jk}} = (B^H A^H)_{ij}.$$

2. If $A^H = -A$ and $Au = \lambda u$, with $u \neq 0$,

$$\lambda u^H u = u^H Au = -u^H A^H u = -(Au)^H u = -\overline{\lambda} u^H u.$$

Thus $\lambda = -\overline{\lambda}$, so its real part is zero.

3.

$$1 = \det(Q^H Q) = |\det Q|^2,$$

Hence the determinant modulus is 1.

4. Hermitian positive definite:

$$z^H Az = \lambda z^H z > 0$$

For a nonzero eigenvector this forces $\lambda > 0$.

Lin Cing
https://github.com/xinhi6716/computer-science-self-study

CH09-E5

1. The given matrix is

$$Q = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix}.$$

$$Q^T Q = \frac{1}{2} \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix} = I.$$

2. The standard basis vectors have inner product zero. Unitarity gives

$$(Qu)^H (Qv) = u^H Q^H Qv = 0.$$

Their norms also remain 1.

3.

$$\det(Q - \lambda I) = \lambda^2 - \sqrt{2}\lambda + 1.$$

$$\lambda = e^{\pm i\pi/4}, \qquad |\lambda| = 1.$$

4. The real matrix has $Q^T = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix} \neq Q$, so it is not Hermitian. Its nonreal eigenvalues give another check. Some matrices, such as the identity, are both unitary and Hermitian; neither property generally implies the other.

CH09-E6

1. $\omega = i$:

$$F_4 = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & i & -1 & -i \\ 1 & -1 & 1 & -1 \\ 1 & -i & -1 & i \end{bmatrix}.$$

2. $c = (1, 2, 0, -1)^T$:

$$y_0 = 2,$$
$$y_1 = 1 + 2i + 0 + i = 1 + 3i,$$
$$y_2 = 1 - 2 + 0 + 1 = 0,$$
$$y_3 = 1 - 2i + 0 - i = 1 - 3i.$$

Therefore

$$y = (2, 1 + 3i, 0, 1 - 3i)^T.$$

3.

$$F_4^H = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & -i & -1 & i \\ 1 & -1 & 1 & -1 \\ 1 & i & -1 & -i \end{bmatrix}.$$
$$F_4^H y = (4, 8, 0, -4)^T,$$

Division by 4 recovers $c = (1, 2, 0, -1)^T$.

4.

$$\|c\|^2 = 1 + 4 + 0 + 1 = 6.$$
$$\|y\|^2 = 4 + 10 + 0 + 10 = 24 = 4\|c\|^2.$$

Lin Cing https://github.com/xinhi6716/computer-science-self-study

CH09-E7


1. The even-indexed coefficients are

$$c' = (c_0, c_2, c_4, c_6)^T,$$

The odd-indexed coefficients are

$$c'' = (c_1, c_3, c_5, c_7)^T.$$

Taking their size-4 DFTs gives $E = F_4 c'$, $O = F_4 c''$.



2. $j = 0, \ldots, 3$:

$$y_j = E_j + \omega_8^j O_j,$$
$$y_{j+4} = E_j - \omega_8^j O_j.$$

The twiddle factor is ω_8^j.

3.

$$T(8) = 2T(4) + 8,$$
$$T(4) = 2T(2) + 4, \qquad T(2) = 2T(1) + 2.$$

Each stage costs linear work and there are $\log_2 n$ stages, so

$$T(n) = \Theta(n \log n).$$

4. Radix-2 requires a power-of-two length. Other lengths can use mixed radices, Bluestein, or zero-padding; padding changes the signal length and frequency grid.

Lin Cing https://github.com/xinhi6716/computer-science-self-study

CH09-E8

1. Let q_j be column j. From $Q^HQ = I$,

$$q_i^H q_j = \delta_{ij},$$

these n columns form an orthonormal basis of $\mathbb{C}^n$. For square Q, $Q^HQ = I$ also gives $Q^{-1} = Q^H$, hence $QQ^H = I$. Writing row i as the column $r_i = (Q_{i1}, \ldots, Q_{in})^T$,

$$r_i^H r_j = \overline{(QQ^H)_{ij}} = \delta_{ij}.$$

the rows also form an orthonormal basis. The two matrix products directly test different vector families.

2. A Hermitian matrix $S = S^H$ has real diagonal entries and hence real trace. Its determinant is the product of real eigenvalues and is real as well.

3.

$$(cF_n)^H(cF_n) = |c|^2 nI.$$

A unitary matrix satisfies

$$|c| = \frac{1}{\sqrt{n}}.$$

> Any global phase $e^{i\phi}$ preserves the norm.

4. Let F_+ and F_- be the unnormalized positive- and negative-sign matrices. For any complex input c,

$$F_-c = \overline{F_+\overline{c}}, \qquad (F_-c)_k = (F_+c)_{(-k) \bmod n}.$$

Simply conjugating the output works only for real input. Both unnormalized matrices satisfy $F_\pm F_\pm^H = nI$; for $Q = F_\pm/\sqrt{n}$, $QQ^H = I$. Parseval's factor depends on scaling, not the sign.

CH09-E9

1. Bit reversal reorders $0, 1, 2, 3$ as $0, 2, 1, 3$, giving
$$(c_0, c_2, c_1, c_3) = (1, 0, 2, -1).$$
2. The first stage is
$$(1, 0) \mapsto (1, 1), \qquad (2, -1) \mapsto (1, 3).$$

The second stage pairs positions $(0, 2)$ and $(1, 3)$, with twiddles 1 and i, respectively:
$$(1, 1) \mapsto (2, 0), \qquad (1, 3) \mapsto (1 + 3i, 1 - 3i).$$

3. Therefore
$$F_4 c = (2, 1 + 3i, 0, 1 - 3i)^T.$$
4. Direct evaluation gives
$$y_1 = 1 + 2i + 0(-1) + (-1)(-i) = 1 + 3i,$$
$$y_3 = 1 - 2i + 0(-1) + (-1)i = 1 - 3i,$$

agreeing with the FFT stages.

CH09-E10

$$A^H = \begin{bmatrix} -i & 0 \\ 1 & 2 \end{bmatrix}, \quad Ax = (i, 0)^T, \quad A^H y = (-i, 1 + 2i)^T.$$

Thus $\langle Ax, y \rangle = (-i, 0)(1, i)^T = -i$, and $\langle x, A^H y \rangle = (1, 0)(-i, 1 + 2i)^T = -i$.

Using A^T instead gives $A^T y = (i, 1 + 2i)^T$, so the right side is i, not $-i$. The difference is the omitted conjugation.

References and attribution

Lin Cing, handwritten linear algebra notes. Gilbert Strang, Introduction to Linear Algebra, 5th ed., Chapter 9.
Original educational synthesis by Lin Cing, licensed under CC BY 4.0.
https://github.com/xinhi6716/computer-science-self-study