Chapter 6 solutions

CH06-E1

1.

$$p_A(\lambda) = (3-\lambda)^2 - 1 = (\lambda - 4)(\lambda - 2).$$

Eigenvalues $4, 2$.

2. For $\lambda = 4$, the equations give $x_1 = x_2$, so

$$E_4 = \text{span}\{(1,1)^T\}.$$

For $\lambda = 2$, they give $x_1 + x_2 = 0$, so

$$E_2 = \text{span}\{(1,-1)^T\}.$$

Both geometric multiplicities are 1.

3.

$$u_0 = (2,0)^T = (1,1)^T + (1,-1)^T.$$
$$u_k = 4^k(1,1)^T + 2^k(1,-1)^T = \begin{bmatrix} 4^k + 2^k \\ 4^k - 2^k \end{bmatrix}.$$

CH06-E2

Lin Cing
https://github.com/xinhi6716/computer-science-self-study

1. Trace 5, det 6:

$$p_A(\lambda) = (\lambda - 3)(\lambda - 2).$$

For eigenvalues 3 and 2, take $x_1 = (2,1)^T$ and $x_2 = (1,1)^T$, respectively.

2.

$$X = \begin{bmatrix} 2 & 1 \\ 1 & 1 \end{bmatrix}, \qquad X^{-1} = \begin{bmatrix} 1 & -1 \\ -1 & 2 \end{bmatrix}.$$
$$A = X \begin{bmatrix} 3 & 0 \\ 0 & 2 \end{bmatrix} X^{-1}.$$

3.

$$A^5 = X \begin{bmatrix} 243 & 0 \\ 0 & 32 \end{bmatrix} X^{-1}.$$

Multiplying gives

$$\begin{bmatrix} 486 & 32 \\ 243 & 32 \end{bmatrix} \begin{bmatrix} 1 & -1 \\ -1 & 2 \end{bmatrix} = \begin{bmatrix} 454 & -422 \\ 211 & -179 \end{bmatrix}.$$

CH06-E3

1. Both triangular diagonal entries are 2.

$$p_A(\lambda) = (\lambda - 2)^2.$$

Algebraic multiplicity 2.

$$(A - 2I)x = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} x = 0$$

Only $x_2 = 0$ is imposed, so

$$E_2 = \operatorname{span}\{(1,0)^T\}, \qquad g_2 = 1.$$

2. Only one eigenvector direction exists, so diagonalization is impossible.

3.

$$A = 2I + N, \qquad N = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \qquad N^2 = 0.$$

Binomial:

$$A^k = 2^k I + k2^{k-1} N = \begin{bmatrix} 2^k & k2^{k-1} \\ 0 & 2^k \end{bmatrix}.$$

Lin Cing
https://github.com/xinhi6716/computer-science-self-study

CH06-E4

1. Matrix

$$\begin{bmatrix} 2 & -1 \\ -1 & 2 \end{bmatrix}$$

has eigenvalues 3 and 1 with orthonormal eigenvectors

$$q_1 = \frac{1}{\sqrt{2}}(1, -1)^T,$$
$$q_2 = \frac{1}{\sqrt{2}}(1, 1)^T.$$

2.

$$u(0) = (1, 3)^T$$
$$q_1^T u(0) = -\sqrt{2}, \qquad q_2^T u(0) = 2\sqrt{2}.$$

3.

$$u(t) = -\sqrt{2}e^{3t}q_1 + 2\sqrt{2}e^{t}q_2.$$

Expanding gives

$$u(t) = \begin{bmatrix} -e^{3t} + 2e^{t} \\ e^{3t} + 2e^{t} \end{bmatrix}.$$

Both eigenvalues are positive. For these initial data, the eigenvalue-3 mode is nonzero and dominates the growth.

Lin Cing https://github.com/xinhi6716/computer-science-self-study

CH06-E5

1.

$$p(\lambda) = \lambda^2 + 1, \qquad \lambda = \pm i.$$

For $\lambda = i$, choose $(1, -i)^T$; for $\lambda = -i$, choose its conjugate $(1, i)^T$.

2. Real solution：

$$u(t) = \begin{bmatrix} c_1 \cos t - c_2 \sin t \\ c_1 \sin t + c_2 \cos t \end{bmatrix}.$$

$u(0) = (c_1, c_2)^T$.

3.

$$\frac{d}{dt}(x^2 + y^2) = 2x(-y) + 2y(x) = 0.$$

Thus the orbit lies on a circle.

4. Matrix

$$\begin{bmatrix} \alpha & -1 \\ 1 & \alpha \end{bmatrix}$$

The eigenvalues are $\alpha \pm i$. For $\alpha < 0$, solutions decay and zero is asymptotically stable. At $\alpha = 0$, bounded rotation gives Lyapunov stability but not asymptotic stability. For $\alpha > 0$, nonzero solutions spiral outward and zero is unstable.

Lin Cing
https://github.com/xinhi6716/computer-science-self-study

CH06-E6

1. Direct transposition gives $S^T = S$, and the energy is
$$2(x_1^2 + x_2^2 + x_3^2 + x_1x_2 + x_2x_3).$$
a symmetric quadratic form.

2. From $Sx = \lambda x$, multiply by x^*.
$$x^*Sx = \lambda x^*x.$$
The left side equals its conjugate and is real. Since $x^*x > 0$, the eigenvalue is real.

3. By part 2 both eigenvalues are real. For $Sx = \lambda x$, $Sy = \mu y$, use the conjugate inner product:
$$\lambda x^*y = (Sx)^*y = x^*Sy = \mu x^*y.$$
Distinct eigenvalues force $x \perp y$ for real eigenvectors, or $x^*y = 0$ for complex choices.

4. There exist an orthonormal eigenbasis Q and a real diagonal Λ such that
$$S = Q\Lambda Q^T.$$
Each vector must belong to the appropriate eigenspace. Orthonormal choices can vary within repeated eigenspaces.

Lin Cing
https://github.com/xinhi6716/computer-science-self-study

CH06-E7

1. Characteristic:
$$p_S(\lambda) = (\lambda - 1)(\lambda - 3),$$
Both eigenvalues, 3 and 1, are positive.

Leading minors:
$$\Delta_1 = 2 > 0, \qquad \Delta_2 = 3 > 0.$$
Energy:
$$x^TSx = x_1^2 + (x_1 - x_2)^2 + x_2^2 > 0$$
for every nonzero x.

2. Spectral coordinates:
$$3y_1^2 + y_2^2 = 1.$$
The semiaxis lengths along y_1 and y_2 are $1/\sqrt{3}$ and 1, respectively.

3. On the unit sphere, the maximum energy is 3 and the minimum is 1, attained at the corresponding unit eigenvectors.

CH06-E8

1. The upper-triangular diagonal entries are all -1.

$$\lambda = -1, -1, -1.$$

Trace -3 , det -1.

2. Let

$$N = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}.$$

$$N^2 = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \qquad N^3 = 0.$$

$$e^{At} = e^{-t} \begin{bmatrix} 1 & t & t^2/2 \\ 0 & 1 & t \\ 0 & 0 & 1 \end{bmatrix}.$$

3. All real parts are $-1 < 0$, so zero is asymptotically stable and $e^{At} \to 0$.

4. In general, asymptotic stability does not imply monotone decay. This matrix does contract in the 2-norm, since the largest eigenvalue of $(A + A^T)/2$ is $-1 + \sqrt{2}/2 < 0$, giving $d\|u\|^2/dt = u^T(A + A^T)u < 0$ for nonzero u; the zero state has derivative zero. A true counterexample is $B = \begin{bmatrix} -1 & 4 \\ 0 & -1 \end{bmatrix}$. Its eigenvalues are both -1, but at $u = (1, 1)^T$, $u^T(B + B^T)u = 4 > 0$. Asymptotic stability ensures eventual decay; a polynomial factor alone does not prove transient growth.

Let $\delta = 1 - \sqrt{2}/2 > 0$. The differential inequality yields $\|e^{At}v\|_2 \le e^{-\delta t}\|v\|_2$ for every v. By the semigroup property, $\|e^{A(s+t)}\|_2 \le e^{-\delta t}\|e^{As}\|_2$ for $s, t \ge 0$; the operator norm therefore decreases monotonically.

CH06-E9

1. A, B invertible:

$$BA = A^{-1}(AB)A.$$

Thus BA is similar to AB, and they share eigenvalues.

2. A diagonalizable A with all eigenvalues zero annihilates every eigenbasis vector.

$$Av_i = 0.$$

Since these vectors span, $A = 0$.

3. A counterexample is

$$N = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$$

It has only eigenvalue zero but is nonzero and defective.

CH06-E10

Since d_5=10 is the full dimension, all later nullities equal 10. Increments for k=1 through 6 are (3,3,2,1,1,0), counting blocks of length at least k. Their adjacent differences are (0,1,1,0,1), so there is one block of each length 2,3,5:

$$J = \mathrm{diag}(J_5(0), J_3(0), J_2(0)).$$

The dimensions add to 10. Rank-nullity gives rank N=10−3=7 and rank N^2=10−6=4. Since d_4=9<10=d_5, $N^4 \neq 0$ and N^5=0, so the least m is 5. The value d_1=3 counts chain endpoints and hence blocks, not maximum chain length.

CH06-E11

Write B=−I+4N, where $N = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$ and $N^2=0$. Then

$$e^{Bt} = e^{-t}(I + 4tN), \qquad u(t) = e^{-t}(1 + 4t, 1)^T/\sqrt{2}.$$

For arbitrary initial data $(a,b)^T$ the solution is $e^{-t}(a+4tb,b)^T$. Both e^{-t} and te^{-t} tend to zero. The evolution operator is also bounded on $t \geq 0$, so the zero solution is asymptotically stable.

Set $h(t) = \|u(t)\|_2^2 = e^{-2t}[(1 + 4t)^2 + 1]/2$. Its derivative is $h'(t) = e^{-2t}(4z - z^2 - 1)$, with $z=1+4t \geq 1$. The only admissible root is $z=2+\sqrt{3}$. The derivative is positive before it and negative after it, so

$$t_* = (1 + \sqrt{3})/4, \quad \max_{t \geq 0} \|u(t)\|_2 = e^{-(1+\sqrt{3})/4}\sqrt{4 + 2\sqrt{3}} \approx 1.379.$$

For C the same initial state gives $e^{-t}u_0$, whose norm decreases from 1. For B, $H = \begin{bmatrix} -1 & 2 \\ 2 & -1 \end{bmatrix}$ satisfies $Hu_0=u_0$, so $h'(0)=2$. But initial state e_1 gives $e^{-t}e_1$ and decays monotonically.

Part 2 fixes one initial direction. The operator norm $\|e^{Bt}\|_2 = \max_{\|v\|_2=1} \|e^{Bt}v\|_2$ optimizes that direction separately at each t. Our curve is only a lower bound on the operator norm.

CH06-E12

$p_A(t) = (t - 2)^2(t + 1)$. Solving $(A - 2I)x = 0$ gives $E_2 = \text{span}\{e_1\}$, so 2 has algebraic multiplicity 2 and geometric multiplicity 1; $E_{-1} = \text{span}\{e_3\}$.

Choose generalized eigenvector $e_2/3$. Then $(A - 2I)(e_2/3) = e_1$, $(A - 2I)e_1 = 0$, and $(A + I)e_3 = 0$. Set

$$P = [e_1, e_2/3, e_3] = \text{diag}(1, 1/3, 1), \quad \det P = 1/3 \neq 0.$$

Checking $AP = PJ$ column by column yields

$$P^{-1}AP = J = \begin{bmatrix} 2 & 1 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & -1 \end{bmatrix}.$$

The superdiagonal 1 follows from $A(e_2/3) = e_1 + 2(e_2/3)$; using e_2 without rescaling would leave a 3 there.

References and attribution

Lin Cing, handwritten linear algebra notes. Gilbert Strang, Introduction to Linear Algebra, 5th ed., Chapter 6.

Original educational synthesis by Lin Cing, licensed under CC BY 4.0.
https://github.com/xinhi6716/computer-science-self-study