Recurrence solutions

DM10-E01

The fixed point solves $a^* = 2a^* - 1$, giving $a^* = 1$. Then $b_n=a_n-1$ has $b_n=2b_{n-1}$,$b_0=2$, hence $a_n=2^{n+1}+1$. Iteration gives $a_n = 2^n a_0 - \sum_{k=0}^{n-1} 2^k = 3 \cdot 2^n - (2^n - 1) = 2^{n+1} + 1$. The initial value is three, and $2a_{n-1}-1=2^{n+1}+1$ verifies substitution.

DM10-E02

The polynomial $r^2-5r+6=(r-2)(r-3)$ gives $a_n=C_1 2^n+C_2 3^n$. Initial equations $C_1+C_2=1$,$2C_1+3C_2=4$ give $C_1=-1$,$C_2=2$, so $a_n=2 \cdot 3^n-2^n$.

DM10-E03

The polynomial is $(r-2)^2$, giving $a_n=(C_1+C_2 n)2^n$. At zero $C_1=2$; at one $2(C_1+C_2)=6$ gives $C_2=1$. Thus $a_n=(n+2)2^n$.

DM10-E04

The homogeneous root is two, giving $C2^n$. Forcing base three does not collide. Trial $A3^n$ gives $A3^n-2A3^{n-1}=A3^{n-1}=3^n$, so $A=3$. The general solution is $a_n=C2^n+3^{n+1}$.

DM10-E05

Near the base $f(2)=2f(1)+2=4$ and $f(4)=2f(2)+4=12$. Conjecture $f(2^m)=(m+1)2^m$. The base $m=0$ gives one. If it holds at m, $f(2^{m+1})=2(m+1)2^m+2^{m+1}=(m+2)2^{m+1}$, proving induction.

For general integers, first prove monotonicity. The base $f(2)>f(1)$ holds. For $n\geq 2$,

$$f(n+1) - f(n) = 2[f(\lfloor (n+1)/2 \rfloor) - f(\lfloor n/2 \rfloor)] + 1.$$

At $n=2k$ the bracket vanishes; at $n=2k+1$ it is $f(k+1)-f(k)$, nonnegative by strong induction. Thus f is nondecreasing. For $m=\lfloor \log_2 n \rfloor$,

$$(m+1)2^m \leq f(n) \leq (m+2)2^{m+1}.$$

For $n\geq 2$ use $m+1\geq \log_2 n$,$2^m\geq n/2$,$m+2\leq 3\log_2 n$,$2^{m+1}<2n$ to obtain

$$\tfrac{1}{2} n \log_2 n \leq f(n) \leq 6n \log_2 n.$$

Hence $\Theta(n \log n)$. Floors change the exact path; this bounding proof does not pretend every n is a power of two.

DM10-E06

With $x_0=0$ all terms stay zero. With $x_0=1$, initial terms are 1,2,6,42 and continue growing without a bounded solution. Nonlinearity rules out the constant-coefficient linear characteristic-polynomial method, without proving that no conceivable closed form exists. Since $x_{n+1} \geq x_n^2$ and $x_1=2$, induction gives $x_n \geq 2^{2^{n-1}}$ for $n \geq 1$, proving unbounded growth.

DM10-E07

The homogeneous solution is $C4^n$. Trial $An+B$ gives $An+B=(4A+3)n+(4B-4A)$. Thus $A=4A+3$ gives $A=-1$, and $B=4B+4$ gives $B=-4/3$. Consequently $b_n=C4^n-n-4/3$. Initial $b_0=1$ gives $C=7/3$:

$$b_n = (7 \cdot 4^n - 3n - 4)/3.$$

Checks give $b_1=7=4 \cdot 1+3$ and $b_2=34=4 \cdot 7+6$.

DM10-E08

The homogeneous polynomial $(r-2)^2$ has root two of multiplicity two. Neither $A2^n$ nor $An2^n$ survives as a particular trial; use $An^2 2^n$. Substitution gives

$$An^2 2^n - 4A(n-1)^2 2^{n-1} + 4A(n-2)^2 2^{n-2} = 2A2^n.$$

Thus $A=3/2$ and $a_n=(C_1+C_2 n+3n^2/2)2^n$. Initial $a_0=1$ gives $C_1=1$; $2(1+C_2+3/2)=5$ gives $C_2=0$. Hence $a_n=(1+3n^2/2)2^n$. At $n=2$ it is $28=4 \cdot 5-4 \cdot 1+12$.

Alternatively, divide by 2^n and put $u_n = a_n/2^n$. Then $u_n - 2u_{n-1} + u_{n-2} = 3$, with $u_0 = 1, u_1 = 5/2$. Set $d_n = u_n - u_{n-1}$. Since $d_1 = 3/2$ and $d_n = d_{n-1} + 3$, we get $d_n = 3n - 3/2$. Summing yields $u_n = 1 + 3n^2/2$, matching coefficient comparison.

n	0	1	2	3	4
a_n	1	5	28	116	400
u_n	1	5/2	7	29/2	25

Checks give $a_3 = 4 \cdot 28 - 4 \cdot 5 + 3 \cdot 8 = 116$ and $a_4 = 4 \cdot 116 - 4 \cdot 28 + 3 \cdot 16 = 400$. The factor n^2 escapes the homogeneous solution space; the constants C_1, C_2 then fit the initial data of the total solution.

DM10-E09

The roots of $(r-1)(r-3)$ give $a_n=C_1+C_2 3^n$. Equations $C_1+C_2=2, C_1+3C_2=7$ give $C_2=5/2, C_1=-1/2$, so $a_n=(5\cdot 3^n-1)/2$.

The OGF equations retain initial corrections:

$$(A-2-7x)-4x(A-2)+3x^2A=0, \qquad A=\frac{2-x}{(1-x)(1-3x)}.$$

Writing $A=\alpha/(1-x)+\beta/(1-3x)$, numerator comparison gives $\alpha+\beta=2, -3\alpha-\beta=-1$, hence $\alpha=-1/2, \beta=5/2$. Extracting coefficients gives the same formula. $A(0)=2$ and $A'(0)=7$ verify the initial numerator.

DM10-E10

Root splitting gives $C_0=1$ and the stated convolution. Summing $n\geq 1$ gives $C-1=xC^2$. The branch with $C(0)=1$ is $C=(1-\sqrt{1-4x})/(2x)$. The generalized binomial expansion of the square root, reindexed by one, gives $C_n=\binom{2n}{n}/(n+1)$. Initial values are 1,1,2,5, and $C_4=\binom{8}{4}/5=14$.

Legal push/pop histories give outputs 123,132,213,231,321. For 231, push 1, push 2, pop 2, push 3, pop 3, pop 1. To output 3 first, both 1 and 2 must already be below it, with 2 above 1. Then 2 must precede 1, ruling out 312. The five outputs match $C_3=5$.

DM10-E11

$P(r)=r^3-2r^2=r^2(r-2)$, so zero has multiplicity two. A complete basis is $\delta_0, \delta_1, 2^n$, where $\delta_j(n)=1$ iff $n=j$. Write $a_n=A\delta_0(n)+B\delta_1(n)+C2^n$. From $a_2=5$, $C=5/4$; from $a_0=3$, $A=7/4$; from $a_1=4$, $B=3/2$. Thus

$$a_n=\tfrac{7}{4}\delta_0(n)+\tfrac{3}{2}\delta_1(n)+\tfrac{5}{4}\,2^n.$$

The terms are 3,4,5,10,20,40; the $n\geq 2$ tail is $5\cdot 2^{n-2}$. Both 0^n and $n0^n$ vanish for $n\geq 1$, regardless of the index-zero convention, so cannot represent arbitrary a_1.

DM10-E12

Internal levels j=0,1,2,3 have 1,2,4,8 nodes. Their total costs are 4,16/3,8,16; sixteen leaves cost one each. Hence T(16)=4+16/3+8+16+16=148/3.

Generally $n=2^k$ and level j costs $n/(k-j)$, $j=0,\ldots,k-1$. Including leaves gives $T(n)=n(1+H_k)$. Integral bounds $\ln(k+1)\le H_k\le 1+\ln k$ give $\Theta(n\log k)=\Theta(n\log\log n)$. At n=1 use the separate base rather than dividing by zero. The ratio $(n/\log n)/n$ tends to zero, so forcing is not $\Theta(n)$; it also has no fixed polynomial gap from critical n. The polynomial-cost theorem does not apply, while layer sums reveal harmonic accumulation.

DM10-E13

Roots of r^2-r+1 are $(1\pm i\sqrt{3})/2=e^{\pm i\pi/3}$. The complex general solution is $Ae^{in\pi/3}+Be^{-in\pi/3}$; the real form is $C\cos(n\pi/3)+D\sin(n\pi/3)$. Initial values give $C=1$ and $1/2+D\sqrt{3}/2=0$, so

$$a_n=\cos(n\pi/3)-\sin(n\pi/3)/\sqrt{3}.$$

The first seven values are 1,0,−1,−1,0,1,1. Both trigonometric terms are unchanged by $n\mapsto n+6$, proving period six. Alternatively $(a_6,a_7)=(1,0)$ returns to the initial pair and recurrence uniqueness proves repetition. A finite observed list alone would not prove it.

Lin Cing
https://github.com/xinhi6716/computer-science-self-study

DM10-E14

Summing from n≥1 gives A−1=2xA+xB,B=xA+xB. Thus B=xA/(1−x), and

$$[(1-2x)(1-x)-x^2]A = 1-x, \quad A = \frac{1-x}{1-3x+x^2}, \quad B = \frac{x}{1-3x+x^2}.$$

For $r_\pm = (3 \pm \sqrt{5})/2$, their sum is three and product one. Numerator checks give

$$B = \frac{1}{\sqrt{5}}\left(\frac{1}{1-r_+x} - \frac{1}{1-r_-x}\right), \quad A = \frac{1}{\sqrt{5}}\left(\frac{r_+-1}{1-r_+x} + \frac{1-r_-}{1-r_-x}\right).$$

Hence

$$b_n = (r_+^n - r_-^n)/\sqrt{5}, \quad a_n = ((r_+-1)r_+^n + (1-r_-)r_-^n)/\sqrt{5}.$$

Direct pairs are (1,0),(2,1),(5,3),(13,8),(34,21). Each component satisfies $u_n=3u_{n-1}-u_{n-2}$ with initial pairs (1,2) and (0,1). In particular $a_4=3\cdot 13-5=34$,$b_4=3\cdot 8-3=21$, agreeing with simultaneous updates.

Generally G−y_0=xMG. The series $H = \sum_{k\ge 0} x^k M^k$ has a finite matrix at every coefficient and satisfies (I−xM)H=H(I−xM)=I by cancellation M^k-M^k at every positive degree. Therefore G=Hy_0=$(I-xM)^{-1}y_0$. Its inverse exists because the constant matrix of I−xM is I, without requiring M invertible. For M=0, G=y_0 still agrees.

References and attribution

Lin Cing, handwritten discrete mathematics notes. Ralph P. Grimaldi, Discrete and Combinatorial Mathematics, 5th ed., Ch. 10.

Original educational synthesis by Lin Cing, licensed under CC BY 4.0.
https://github.com/xinhi6716/computer-science-self-study