Chapter 1 exercises

Use real column vectors. Unless stated otherwise, show the conditions and derivation, rather than only the final answer.

CH01-E1

Let $v = (2, 1)^T$ and $w = (1, 2)^T$. Find real c, d such that

$$cv + dw = (5, 4)^T.$$

State whether the solution is unique and explain why.

CH01-E2

For $u = (3, 4)^T$ and $v = (4, 3)^T$, answer the following.

1. Find the cosine of their angle.
2. Check the Cauchy–Schwarz inequality.
3. Find a nonzero z orthogonal to u, and describe the complete set of vectors orthogonal to u.

Lin Cing
https://github.com/xinhi6716/computer-science-self-study

CH01-E3

Let

$$A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}.$$

Compute $A(2, -1)^T$ by the column and row views. Check that they agree.

CH01-E4

Let $a_1 = (1, 2)^T$ and $a_2 = (2, 4)^T$.

1. Determine whether a_1, a_2 are independent.
2. Determine whether

$$a_1 x_1 + a_2 x_2 = (3, 5)^T$$

has a solution. 3. Replace the right-hand side by $(3, 6)^T$ and find the complete solution set.

CH01-E5

Let

$$A = \begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix}.$$

Chapter 1 exercises (continued)

Check invertibility and derive A^{-1}. Verify the inverse by multiplication rather than quoting an answer.

CH01-E6

Apply the ideas across vector spaces.

1. In $\mathcal{P}_2$, use the basis $B = (1, t, t^2)$ and let $p_1 = 1 + 2t$、$p_2 = 1 - t$、$p_3 = t + t^2$。 Form the coordinate matrix D with these three coordinate columns. Determine independence and express $q(t) = 4 + 3t - t^2$ as a linear combination of p_1, p_2, p_3.
2. In $M_2(\mathbb{R})$, use column-major coordinates and let

$$A_1 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \quad A_2 = \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}, \quad A_3 = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}.$$

Determine whether the matrices are independent. If they are, determine whether they form a basis of $M_2(\mathbb{R})$.

CH01-E7

Let

$$J = \begin{bmatrix} 3 & 1 \\ 0 & 3 \end{bmatrix}, \qquad r(t) = t^3 - 2t + 4.$$

1. Without assuming diagonalizability, prove for every $A \in \mathbb{C}^{n \times n}$ that $Av = \lambda v$ implies $r(A)v = r(\lambda)v$. Prove the reverse spectral inclusion by factoring a polynomial, and apply the result to J.
2. Use $J = 3I + N$, $N^2 = 0$, to compute $r(J)$ completely.
3. If another matrix B has eigenvalue 2 with a one-dimensional eigenspace, does having only one eigenvalue imply $B = 2I$? Give a counterexample or a justification.

Lin Cing https://github.com/xinhi6716/computer-science-self-study

References and attribution

Gilbert Strang. Introduction to Linear Algebra, 5th Edition, 2016.

Original educational synthesis by Lin Cing. Original material is licensed under CC BY 4.0. The referenced textbook is not distributed with these notes.

https://github.com/xinhi6716/computer-science-self-study