Chapter 1 solutions

CH01-E1

Equating components gives

$$2c + d = 5, \qquad c + 2d = 4.$$

Twice the first equation minus the second gives

$$4c + 2d - (c + 2d) = 10 - 4,$$

Thus $3c = 6$ and $c = 2$. Substitution into the first equation gives $d = 1$.

The coefficient matrix

$$\begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix}$$

has determinant $2 \cdot 2 - 1 \cdot 1 = 3 \neq 0$, so the solution is unique.

CH01-E2

The inner product is

$$u^T v = 3 \cdot 4 + 4 \cdot 3 = 24.$$

Both norms are 5, so

$$\cos\theta = \frac{24}{25}.$$

Cauchy–Schwarz states

$$|u^T v| \leq \|u\|_2 \|v\|_2.$$

Here $24 < 25$. The strict inequality agrees with the fact that these vectors are not collinear.

A vector $z = (z_1, z_2)^T$ orthogonal to u must satisfy

$$3z_1 + 4z_2 = 0.$$

Taking $z_1 = 4$ gives $z_2 = -3$, so $z = (4, -3)^T$ is one nonzero answer. All solutions are

$$z = t(4, -3)^T, \qquad t \in \mathbb{R},$$

the line through the origin perpendicular to u. To restrict the set to nonzero vectors, require $t \neq 0$.

CH01-E3

Column view:

$$A(2, -1)^T = 2(1, 3)^T - (2, 4)^T = (0, 2)^T.$$

Lin Cing https://github.com/xinhi6716/computer-science-self-study

Chapter 1 solutions (continued)

Row view:

The first row gives $(1,2)\cdot(2,-1)=2-2=0$. The second gives $(3,4)\cdot(2,-1)=6-4=2$. Hence the output is again $(0,2)^T$.

The two decompositions compute the same vector.

CH01-E4

Since $a_2=2a_1$, the nonzero coefficient vector $(c_1,c_2)=(2,-1)$ satisfies

$$2a_1-a_2=0.$$

The vectors are dependent and span the line

$$\{t(1,2)^T : t\in\mathbb{R}\}.$$

For $(3,5)^T$ to belong to this span, its second component would have to be twice its first. Since $5\neq 2\cdot 3$, there is no solution.

For the target $(3,6)^T$, every solution satisfies

$$x_1+2x_2=3.$$

Set $x_2=t$. Then $x_1=3-2t$, and the complete solution set is

$$\left\{\begin{bmatrix}3\\0\end{bmatrix}+t\begin{bmatrix}-2\\1\end{bmatrix} : t\in\mathbb{R}\right\}.$$

https://github.com/xinhi6716/computer-science-self-study

CH01-E5

The determinant is

$$\det A=2\cdot 2-1\cdot 1=3\neq 0.$$

Over the real field, a square matrix with nonzero determinant is invertible. The 2×2 formula gives

$$A^{-1}=\frac{1}{3}\begin{bmatrix}2&-1\\-1&2\end{bmatrix}.$$

Verification gives

$$\begin{bmatrix}2&1\\1&2\end{bmatrix}\left(\frac{1}{3}\begin{bmatrix}2&-1\\-1&2\end{bmatrix}\right)=\frac{1}{3}\begin{bmatrix}3&0\\0&3\end{bmatrix}=I_2.$$

Matrix multiplication does not commute in general. A matrix and its inverse nevertheless multiply to I_2 in both orders.

CH01-E6

1. The coordinate vectors are

Chapter 1 solutions (continued)

$$[p_1]_B = (1,2,0)^T, \quad [p_2]_B = (1,-1,0)^T, \quad [p_3]_B = (0,1,1)^T.$$

Thus

$$D = \begin{bmatrix} 1 & 1 & 0 \\ 2 & -1 & 1 \\ 0 & 0 & 1 \end{bmatrix}.$$

Expansion along the third row gives $\det D = (+1)\det \begin{bmatrix} 1 & 1 \\ 2 & -1 \end{bmatrix} = -3 \neq 0$，so the coordinate columns and the three polynomials are independent.

The coordinates of q are $(4,3,-1)^T$. Solve

$$c_1 + c_2 = 4, \qquad 2c_1 - c_2 + c_3 = 3, \qquad c_3 = -1.$$

The first equation gives $c_2 = 4 - c_1$. Substitution into the second gives $2c_1 - (4 - c_1) - 1 = 3$, hence $3c_1 = 8$, $c_1 = 8/3$, and $c_2 = 4/3$. Therefore

$$q = \frac{8}{3}p_1 + \frac{4}{3}p_2 - p_3.$$

Expanding verifies the constant term $8/3 + 4/3 = 4$, the coefficient of t, $16/3 - 4/3 - 1 = 3$, and the coefficient -1 of t^2.

2. The column-major coordinate columns are

$$(1,0,0,1)^T, \quad (1,0,1,0)^T, \quad (0,1,1,0)^T.$$

If $c_1A_1 + c_2A_2 + c_3A_3 = 0$, the entries give $c_1 + c_2 = 0$、$c_2 + c_3 = 0$、$c_3 = 0$、$c_1 = 0$, so the matrices are independent. However, $M_2(\mathbb{R})$ has dimension 4. Three vectors cannot span this four-dimensional space, so the independent set is not a basis.

CH01-E7

1. For $Av = \lambda v$, $v \neq 0$, induction gives $A^k v = \lambda^k v$, hence $r(A)v = r(\lambda)v$ for every polynomial. Conversely, let $\mu \in \operatorname{spec}(r(A))$. For nonconstant r, factor $r(t) - \mu = c\prod_j (t - \zeta_j)$ over $\mathbb{C}$. From

$$0 = \det(r(A) - \mu I) = c^n \prod_j \det(A - \zeta_j I)$$

at least one ζ_j is an eigenvalue of A, with $r(\zeta_j) = \mu$. A constant polynomial is handled by $r(A) = cI$. Thus $\operatorname{spec}(r(A)) = r(\operatorname{spec}(A))$, without diagonalization.

Write $J = 3I + N$, where

$$N = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \qquad N^2 = 0.$$

Lin Cing https://github.com/xinhi6716/computer-science-self-study

Chapter 1 solutions (continued)

For every positive integer k, $J^k = 3^k I + k3^{k-1}N$. The polynomial $r(t) = t^3 - 2t + 4$ therefore gives

$$r(J) = r(3)I + r'(3)N.$$

This follows directly from the binomial expansion. If $Ju = 3u$, each term satisfies $J^k u = 3^k u$, so $r(J)u = r(3)u = 25u$.

2.

$$r'(t) = 3t^2 - 2, \qquad r'(3) = 25,$$

Also $r(3) = 27 - 6 + 4 = 25$. Hence

$$r(J) = 25I + 25N = 25 \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}.$$

3. No. The counterexample $\begin{bmatrix} 2 & 1 \\ 0 & 2 \end{bmatrix}$ has only eigenvalue 2, but its eigenvectors lie in the line spanned by e_1, and the matrix is not $2I$. The eigenvalue determines the diagonal of a Jordan block. Without a full independent eigenbasis, the generalized-eigenvector structure can retain nonzero off-diagonal entries.

Lin Cing
https://github.com/xinhi6716/computer-science-self-study

References and attribution

Gilbert Strang. Introduction to Linear Algebra, 5th Edition, 2016.

Original educational synthesis by Lin Cing. Original material is licensed under CC BY 4.0. The referenced textbook is not distributed with these notes.

https://github.com/xinhi6716/computer-science-self-study