Chapter 6 solutions

CH06-E1

1.

$$p_A(\lambda) = (3-\lambda)^2 - 1 = (\lambda - 4)(\lambda - 2).$$

Eigenvalues $4, 2$.

2. For $\lambda = 4$, the equations give $x_1 = x_2$, so

$$E_4 = \text{span}\{(1,1)^T\}.$$

For $\lambda = 2$, they give $x_1 + x_2 = 0$, so

$$E_2 = \text{span}\{(1,-1)^T\}.$$

Both geometric multiplicities are 1.

3.

$$u_0 = (2,0)^T = (1,1)^T + (1,-1)^T.$$

$$u_k = 4^k(1,1)^T + 2^k(1,-1)^T = \begin{bmatrix} 4^k + 2^k \\ 4^k - 2^k \end{bmatrix}.$$

CH06-E2

1. Trace 5, det 6:

$$p_A(\lambda) = (\lambda - 3)(\lambda - 2).$$

For eigenvalues 3 and 2, take $x_1 = (2,1)^T$ and $x_2 = (1,1)^T$, respectively.

2.

$$X = \begin{bmatrix} 2 & 1 \\ 1 & 1 \end{bmatrix}, \qquad X^{-1} = \begin{bmatrix} 1 & -1 \\ -1 & 2 \end{bmatrix}.$$

$$A = X \begin{bmatrix} 3 & 0 \\ 0 & 2 \end{bmatrix} X^{-1}.$$

3.

$$A^5 = X \begin{bmatrix} 243 & 0 \\ 0 & 32 \end{bmatrix} X^{-1}.$$

Multiplying gives

$$\begin{bmatrix} 486 & 32 \\ 243 & 32 \end{bmatrix} \begin{bmatrix} 1 & -1 \\ -1 & 2 \end{bmatrix} = \begin{bmatrix} 454 & -422 \\ 211 & -179 \end{bmatrix}.$$

CH06-E3

1. Both triangular diagonal entries are 2.

$$p_A(\lambda) = (\lambda - 2)^2.$$

Lin Cing https://github.com/xinhi6716/computer-science-self-study

Chapter 6 solutions (continued)

Algebraic multiplicity 2.

$$(A - 2I)x = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} x = 0$$

Only $x_2 = 0$ is imposed, so

$$E_2 = \text{span}\{(1,0)^T\}, \qquad g_2 = 1.$$

2. Only one eigenvector direction exists, so diagonalization is impossible.

3.

$$A = 2I + N, \qquad N = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \qquad N^2 = 0.$$

Binomial:

$$A^k = 2^k I + k2^{k-1} N = \begin{bmatrix} 2^k & k2^{k-1} \\ 0 & 2^k \end{bmatrix}.$$

CH06-E4

1. Matrix

$$\begin{bmatrix} 2 & -1 \\ -1 & 2 \end{bmatrix}$$

has eigenvalues 3 and 1 with orthonormal eigenvectors

$$q_1 = \frac{1}{\sqrt{2}}(1,-1)^T,$$
$$q_2 = \frac{1}{\sqrt{2}}(1,1)^T.$$

2.

$$u(0) = (1,3)^T$$
$$q_1^T u(0) = -\sqrt{2}, \qquad q_2^T u(0) = 2\sqrt{2}.$$

3.

$$u(t) = -\sqrt{2}e^{3t} q_1 + 2\sqrt{2}e^{t} q_2.$$

Expanding gives

$$u(t) = \begin{bmatrix} -e^{3t} + 2e^{t} \\ e^{3t} + 2e^{t} \end{bmatrix}.$$

Both eigenvalues are positive. For these initial data, the eigenvalue-3 mode is nonzero and dominates the growth.

Chapter 6 solutions (continued)

CH06-E5

1.

$$p(\lambda) = \lambda^2 + 1, \qquad \lambda = \pm i.$$

For $\lambda = i$, choose $(1, -i)^T$; for $\lambda = -i$, choose its conjugate $(1, i)^T$.

2. Real solution：

$$u(t) = \begin{bmatrix} c_1 \cos t - c_2 \sin t \\ c_1 \sin t + c_2 \cos t \end{bmatrix}.$$

$u(0) = (c_1, c_2)^T$ 。

3.

$$\frac{d}{dt}(x^2 + y^2) = 2x(-y) + 2y(x) = 0.$$

Thus the orbit lies on a circle.

4. Matrix

$$\begin{bmatrix} \alpha & -1 \\ 1 & \alpha \end{bmatrix}$$

The eigenvalues are $\alpha \pm i$. For $\alpha < 0$, solutions decay and zero is asymptotically stable. At $\alpha = 0$, bounded rotation gives Lyapunov stability but not asymptotic stability. For $\alpha > 0$, nonzero solutions spiral outward and zero is unstable.

CH06-E6

1. Direct transposition gives $S^T = S$, and the energy is

$$2(x_1^2 + x_2^2 + x_3^2 + x_1x_2 + x_2x_3).$$

a symmetric quadratic form.

2. From $Sx = \lambda x$, multiply by x^*.

$$x^*Sx = \lambda x^*x.$$

The left side equals its conjugate and is real. Since $x^*x > 0$, the eigenvalue is real.

3. $Sx = \lambda x$, $Sy = \mu y$:

$$\lambda x^T y = (Sx)^T y = x^T Sy = \mu x^T y.$$

Distinct eigenvalues force $x \perp y$ for real eigenvectors, or $x^*y = 0$ for complex choices.

4. There exist an orthonormal eigenbasis Q and a real diagonal Λ such that

$$S = Q\Lambda Q^T.$$

Chapter 6 solutions (continued)

Each vector must belong to the appropriate eigenspace. Orthonormal choices can vary within repeated eigenspaces.

CH06-E7

1. Characteristic:

$$p_S(\lambda) = (\lambda - 1)(\lambda - 3),$$

Both eigenvalues, 3 and 1, are positive.

Leading minors:

$$\Delta_1 = 2 > 0, \qquad \Delta_2 = 3 > 0.$$

Energy:

$$x^T S x = x_1^2 + (x_1 - x_2)^2 + x_2^2 > 0$$

for every nonzero x.

2. Spectral coordinates:

$$3y_1^2 + y_2^2 = 1.$$

$|y_1|$ axis length $1/\sqrt{3}$, $|y_2|$ axis length 1.

3. On the unit sphere, the maximum energy is 3 and the minimum is 1, attained at the corresponding unit eigenvectors.

CH06-E8

1. The upper-triangular diagonal entries are all -1.

$$\lambda = -1, -1, -1.$$

Trace -3 › det -1 ∘

2. Let

$$N = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}.$$

$$N^2 = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \qquad N^3 = 0.$$

$$e^{At} = e^{-t} \begin{bmatrix} 1 & t & t^2/2 \\ 0 & 1 & t \\ 0 & 0 & 1 \end{bmatrix}.$$

Lin Cing https://github.com/xinhi6716/computer-science-self-study

Chapter 6 solutions (continued)

3. All real parts are $-1 < 0$, so zero is asymptotically stable and $e^{At} \to 0$.
4. In general, asymptotic stability does not imply monotone decay. This matrix does contract in the 2-norm, since the largest eigenvalue of $(A + A^T)/2$ is $-1 + \sqrt{2}/2 < 0$, giving $d\|u\|^2/dt = u^T(A + A^T)u < 0$. A true counterexample is $B = \begin{bmatrix} -1 & 4 \\ 0 & -1 \end{bmatrix}$. Its eigenvalues are both -1, but at $u = (1, 1)^T$, $u^T(B + B^T)u = 4 > 0$. Asymptotic stability ensures eventual decay; a polynomial factor alone does not prove transient growth.

CH06-E9

1. A, B invertible:

$$BA = A^{-1}(AB)A.$$

Thus BA is similar to AB, and they share eigenvalues.

2. A diagonalizable A with all eigenvalues zero annihilates every eigenbasis vector.

$$Av_i = 0.$$

Since these vectors span, $A = 0$.

3. A counterexample is

$$N = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$$

It has only eigenvalue zero but is nonzero and defective.

Lin Cing https://github.com/xinhi6716/computer-science-self-study

Chapter 6 solutions (continued)

Verification code appendix: eigen_dynamics.py

```
"""Check eigen-decompositions, discrete powers, ODE and definiteness examples."""

from math import exp, sqrt


Vec = tuple[float, ...]
Mat = tuple[tuple[float, ...], ...]


def matvec(matrix: Mat, vector: Vec) -> Vec:
    return tuple(
        sum(matrix[i][j] * vector[j] for j in range(len(vector)))
        for i in range(len(matrix))
    )


def matmul(left: Mat, right: Mat) -> Mat:
    return tuple(
        tuple(
            sum(left[i][k] * right[k][j] for k in range(len(right)))
            for j in range(len(right[0]))
        )
        for i in range(len(left))
    )


def residual(matrix: Mat, vector: Vec, value: float) -> float:
    image = matvec(matrix, vector)
    return sqrt(sum((image[i] - value * vector[i]) ** 2 for i in range(len(vector))))


def matrix_power(matrix: Mat, exponent: int) -> Mat:
    result = tuple(
        tuple(1.0 if i == j else 0.0 for j in range(len(matrix)))
        for i in range(len(matrix))
    )
    base = matrix
    while exponent:
        if exponent & 1:
            result = matmul(result, base)
        base = matmul(base, base)
        exponent >>= 1
    return result


a = ((3.0, 1.0), (1.0, 3.0))
x1 = (1 / sqrt(2), 1 / sqrt(2))
x2 = (1 / sqrt(2), -1 / sqrt(2))
assert residual(a, x1, 4.0) < 1e-12
assert residual(a, x2, 2.0) < 1e-12
```

Lin Cing https://github.com/xinhi6716/computer-science-self-study

Chapter 6 solutions (continued)

```
assert abs(x1[0] * x2[0] + x1[1] * x2[1]) < 1e-12

u0 = (2.0, 0.0)
for k in range(8):
    closed = (
        4**k + 2**k,
        4**k - 2**k,
    )
    iterative = matvec(matrix_power(a, k), u0)
    assert closed == iterative

defective = ((2.0, 1.0), (0.0, 2.0))
assert matrix_power(defective, 5) == (
    (32.0, 80.0),
    (0.0, 32.0),
)

t = 0.75
system = ((2.0, -1.0), (-1.0, 2.0))
closed_u = (
    -exp(3.0 * t) + 2.0 * exp(t),
    exp(3.0 * t) + 2.0 * exp(t),
)
steps = 200
dt = t / steps
u = (1.0, 3.0)
for _ in range(steps):
    k1 = matvec(system, u)
    k2 = matvec(system, tuple(u[i] + 0.5 * dt * k1[i] for i in range(2)))
    k3 = matvec(system, tuple(u[i] + 0.5 * dt * k2[i] for i in range(2)))
    k4 = matvec(system, tuple(u[i] + dt * k3[i] for i in range(2)))
    u = tuple(
        u[i] + dt * (k1[i] + 2.0 * k2[i] + 2.0 * k3[i] + k4[i]) / 6.0
        for i in range(2)
    )
assert max(abs(u[i] - closed_u[i]) for i in range(2)) < 1e-8

positive_definite = ((2.0, -1.0), (-1.0, 2.0))
for probe in ((1.0, 0.0), (0.0, 1.0), (1.0, -1.0), (2.0, 1.0)):
    energy = probe[0] * matvec(positive_definite, probe)[0] + probe[1] * matvec(
        positive_definite, probe
    )[1]
    assert energy > 0

print("A^100 =", matrix_power(a, 100))
print("defective A^5 =", matrix_power(defective, 5))
print("u(0.75) =", closed_u)
print("energy samples are positive")
```

Lin Cing https://github.com/xinhi6716/computer-science-self-study

Chapter 6 solutions (continued)

References and attribution

Gilbert Strang. Introduction to Linear Algebra, 5th Edition, 2016.

Original educational synthesis by Lin Cing. Original material is licensed under CC BY 4.0. The referenced textbook is not distributed with these notes.

https://github.com/xinhi6716/computer-science-self-study

Lin Cing
https://github.com/xinhi6716/computer-science-self-study