第 4 章練習解答

CH04-E1

1. 解

$$x_1 + 2x_2 = 0, \qquad x_2 + x_3 = 0.$$

令 $x_2 = t$，則 $x_1 = -2t, x_3 = -t$。所以

$$N(A) = \text{span}\{(-2, 1, -1)^T\}.$$

2. A 的兩 rows 為 $r_1 = (1, 2, 0)^T, r_2 = (0, 1, 1)^T$，形成 row space 的 basis。對 $n = (-2, 1, -1)^T$：

$$r_1^T n = -2 + 2 = 0, \qquad r_2^T n = 1 - 1 = 0.$$

因此 row space 的每一條線性組合都與 n 正交，證明 $C(A^T) \subseteq N(A)^\perp$；dimensions 為 2 與 1，和為 3，故兩者互為 orthogonal complements。

3. A 是 2×3，有兩個獨立 rows，rank 為 2，因此 $C(A)$ 是 $\mathbb{R}^2$ 的 2 維子空間，只能等於 $\mathbb{R}^2$。left nullspace 的 dimension：

$$\dim N(A^T) = m - r = 2 - 2 = 0,$$

所以 $N(A^T) = \{0\}$。

Lin Cing
https://github.com/xinhi6716/computer-science-self-study

CH04-E2

1. $a^T a = 3$、$a^T b = 2 + 1 + 4 = 7$，故

$$p = \frac{7}{3} a = \frac{7}{3}(1, 1, 1)^T.$$

2.

$$e = b - p = \left(2 - \frac{7}{3}, 1 - \frac{7}{3}, 4 - \frac{7}{3}\right)^T = \frac{1}{3}(-1, -4, 5)^T.$$

3. $a^T e = \frac{1}{3}(-1 - 4 + 5) = 0$，所以 residual 垂直於整條 line。

4. line projection matrix是

$$P = \frac{a a^T}{a^T a} = \frac{1}{3} \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix}.$$

它顯然對稱。全 1 矩陣 J_3 的每個乘積項相加為 3，故

$$P^2 = \frac{1}{3^2} \cdot 3\, J_3 = \frac{1}{3} J_3 = P,$$

其中 J_3 是全 1 matrix。

第 4 章練習解答 （續）

CH04-E3

1.

$$A = \begin{bmatrix} 1 & 0 \\ 1 & 1 \\ 1 & 2 \end{bmatrix}, \quad x = \begin{bmatrix} c \\ m \end{bmatrix}, \quad b = \begin{bmatrix} 1 \\ 2 \\ 2 \end{bmatrix}.$$

2.

$$A^T A = \begin{bmatrix} 3 & 3 \\ 3 & 5 \end{bmatrix}, \qquad A^T b = \begin{bmatrix} 5 \\ 6 \end{bmatrix}.$$

方程是

$$3c + 3m = 5, \qquad 3c + 5m = 6.$$

兩式相減得 $2m = 1$，所以 $m = 1/2$；代回第一式得 $c = 7/6$。

3.

$$p = Ax = \left(\frac{7}{6}, \frac{7}{6} + \frac{3}{6}, \frac{7}{6} + \frac{6}{6}\right)^T = \left(\frac{7}{6}, \frac{5}{3}, \frac{13}{6}\right)^T.$$

$$e = b - p = \left(-\frac{1}{6}, \frac{1}{3}, -\frac{1}{6}\right)^T.$$

因 e 之和為 0，constant column 與它正交。再者

$$t \cdot e = 0 + \frac{1}{3} + \left(-\frac{1}{3}\right) = 0.$$

故 $A^T e = 0$。

4. $\|e\|^2 = 2(1/36) + 4/36 = 6/36 = 1/6$。這是所有 c, m 中最小的 residual 平方和。

CH04-E4

1. 因 $A^T A$ 對稱，其 inverse 也對稱：

$$P^T = A((A^T A)^{-1})^T A^T = A(A^T A)^{-1} A^T = P.$$

又

$$P^2 = A(A^T A)^{-1} A^T A(A^T A)^{-1} A^T = P.$$

2. $I - P$ 對稱。並且

$$(I - P)^2 = I - 2P + P^2 = I - P.$$

所以是 orthogonal projection。交叉項

$$P(I - P) = P - P^2 = 0.$$

其 range 與 P 的 range 正交，因 $P^T(I - P) = P(I - P) = 0$。

Lin Cing https://github.com/xinhi6716/computer-science-self-study

第 4 章練習解答 （續）

3. 若 $A^TAz=0$，左乘 z^T 得

$$0=z^TA^TAz=\|Az\|^2.$$

所以 $Az=0$。independent columns 使 $z=0$，故 $N(A^TA)=\{0\}$，A^TA 可逆。

4. dependent columns 給非零 z 使 $Az=0$。于是

$$A^TAz=0,$$

所以 A^TA 有非零 nullspace，不可逆。

CH04-E5

1. a_1 的長度是 $\sqrt{2}$，故

$$q_1=\left(\frac{1}{\sqrt{2}},\frac{1}{\sqrt{2}},0\right)^T.$$

又 $q_1^Ta_2=1/\sqrt{2}$，

$$z_2=a_2-\frac{1}{2}(1,1,0)^T=\left(\frac{1}{2},-\frac{1}{2},1\right)^T.$$

$$\|z_2\|=\sqrt{\frac{1}{4}+\frac{1}{4}+1}=\sqrt{\frac{3}{2}},$$

$$q_2=\frac{1}{\sqrt{6}}(1,-1,2)^T.$$

2. Gram-Schmidt 的 coefficients：

$$a_1=\|a_1\|q_1=\sqrt{2}\,q_1,$$

$$a_2=\frac{1}{\sqrt{2}}q_1+\sqrt{\frac{3}{2}}q_2.$$

所以

$$R=\begin{bmatrix}\sqrt{2} & 1/\sqrt{2}\\ 0 & \sqrt{3/2}\end{bmatrix}.$$

3. $q_1^Tq_1=q_2^Tq_2=1$，且

$$q_1^Tq_2=\frac{1+(-1)+0}{\sqrt{12}}=0,$$

所以 $Q^TQ=I_2$。按 columns 展開

$$\sqrt{2}q_1=a_1,$$

$$\frac{1}{\sqrt{2}}q_1+\sqrt{\frac{3}{2}}q_2=\left(\frac{1}{2},\frac{1}{2},0\right)^T+\left(\frac{1}{2},-\frac{1}{2},1\right)^T=a_2,$$

故 $QR=A$。

Lin Cing https://github.com/xinhi6716/computer-science-self-study

第 4 章練習解答 （續）

CH04-E6

1.

$$Q^T b = \left(\frac{1+2}{\sqrt{2}}, \frac{1-2}{\sqrt{6}}\right)^T = \left(\frac{3}{\sqrt{2}}, -\frac{1}{\sqrt{6}}\right)^T.$$

解

$$\sqrt{2}\,x_1 + \frac{1}{\sqrt{2}}x_2 = \frac{3}{\sqrt{2}}, \qquad \sqrt{\frac{3}{2}}x_2 = -\frac{1}{\sqrt{6}}.$$

第二式得 $x_2 = -1/3$，代回第一式得 $x_1 = 5/3$。因此最小平方解是

$$x = \begin{bmatrix} 5/3 \\ -1/3 \end{bmatrix}.$$

2.

$$p = Ax = \left(\frac{4}{3}, \frac{5}{3}, -\frac{1}{3}\right)^T,$$

$$e = b - p = \frac{1}{3}(-1, 1, 1)^T, \qquad \|e\|^2 = \frac{1}{3}.$$

3.

$$a_1^T e = -\frac{1}{3} + \frac{1}{3} = 0,$$

$$a_2^T e = -\frac{1}{3} + \frac{1}{3} = 0.$$

此 residual 非零，不能把「最小平方」誤解成「精確解」。逐欄正交正是 normal equation $A^T e = 0$ 的座標形式；full column rank 使這組參數唯一。

CH04-E7

1. 第二 column 是第一 column 的倍數，所以 columns dependent，rank 為 1。
2. 取 $a = (1, 2)^T$。則

$$a^T a = 5, \qquad a^T b = 1 + 6 = 7,$$

$$p = \frac{7}{5}a = \left(\frac{7}{5}, \frac{14}{5}\right)^T.$$

3.

$$A^T A = \begin{bmatrix} 5 & 5 \\ 5 & 5 \end{bmatrix}, \qquad A^T b = \begin{bmatrix} 7 \\ 7 \end{bmatrix}.$$

所有 minimizers 為

第 4 章練習解答 （續）

$$x^\star = \begin{bmatrix} 7/5 \\ 0 \end{bmatrix} + t \begin{bmatrix} -1 \\ 1 \end{bmatrix}, \quad t \in \mathbb{R}.$$

4. 若 $Ax_1 = Ax_2 = p$，差 $d = x_1 - x_2$ 在 $N(A)$ 中。A 有兩個相同 columns，nullspace 為 $\text{span}\{(-1,1)\}$，所以 $x^\star$ 有非唯一方向；但 A 作用後兩者都映射到同一個 column-space point p。投影到 $C(A)$ 仍是唯一的最近點。

CH04-E8

1. 直接相乘：

$$P^2 = \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} = P.$$

2. 但

$$P^T = \begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix} \neq P.$$

3. $P(x,y)^T = (x+y, 0)^T$，所以

$$R(P) = \text{span}\{(1,0)^T\}, \qquad N(P) = \text{span}\{(-1,1)^T\}.$$

4. $R(P)$ 與 $N(P)$ 並非正交：$(1,0)^T \cdot (-1,1)^T = -1 \neq 0$。因此 $P^2 = P$ 只說明它是斜投影；正交投影還需要 $P^T = P$。

Lin Cing
https://github.com/xinhi6716/computer-science-self-study

第 4 章練習解答 （續）

原創驗算程式附錄: projection_and_qr.py

```
"""Check the line projection and Gram-Schmidt example in chapter four."""

from math import sqrt


def dot(left, right):
    return sum(x * y for x, y in zip(left, right))


def scale(value, vector):
    return [value * x for x in vector]


def add(left, right):
    return [x + y for x, y in zip(left, right)]


def norm(vector):
    return sqrt(dot(vector, vector))


a = [1.0, 1.0, 1.0]
b = [2.0, 1.0, 4.0]
p = scale(dot(a, b) / dot(a, a), a)
e = add(b, scale(-1.0, p))
assert abs(dot(a, e)) < 1e-12
assert abs(norm(e) - sqrt(42) / 3) < 1e-12

a1 = [1.0, 1.0, 0.0]
```

Lin Cing
https://github.com/xinhi6716/computer-science-self-study

第 4 章練習解答 （續）

```
a2 = [1.0, 0.0, 1.0]
q1 = scale(1.0 / norm(a1), a1)
z2 = add(a2, scale(-dot(q1, a2), q1))
q2 = scale(1.0 / norm(z2), z2)
assert abs(dot(q1, q1) - 1.0) < 1e-12
assert abs(dot(q2, q2) - 1.0) < 1e-12
assert abs(dot(q1, q2)) < 1e-12

r11 = sqrt(2.0)
r12 = 1.0 / sqrt(2.0)
r22 = sqrt(1.5)
reconstructed_a1 = scale(r11, q1)
reconstructed_a2 = add(scale(r12, q1), scale(r22, q2))
assert all(abs(x - y) < 1e-12 for x, y in zip(reconstructed_a1, a1))
assert all(abs(x - y) < 1e-12 for x, y in zip(reconstructed_a2, a2))

target = a2
qtb = [dot(q1, target), dot(q2, target)]
x2 = qtb[1] / r22
x1 = (qtb[0] - r12 * x2) / r11
fit = add(scale(x1, a1), scale(x2, a2))
assert abs(x1) < 1e-12 and abs(x2 - 1.0) < 1e-12
assert norm(add(target, scale(-1.0, fit))) < 1e-12

print("line projection residual norm =", norm(e))
print("q1 =", q1)
print("q2 =", q2)
```

Lin Cing
https://github.com/xinhi6716/computer-science-self-study

參考資料與署名

Gilbert Strang. Introduction to Linear Algebra, 5th Edition, 2016.

Lin Cing 原創教學整理。原創內容以 CC BY 4.0 授權；參考書籍不隨教材散布。
https://github.com/xinhi6716/computer-science-self-study