Chapter 7 solutions

CH07-E1

1.

$A^TA = \begin{bmatrix} 3 & 0 \\ 0 & 2 \end{bmatrix}$.

Eigenvalues $3, 2$, eigenvectors $v_1 = (1,0)^T$, $v_2 = (0,1)^T$.

2.

$\sigma_1 = \sqrt{3}, \qquad \sigma_2 = \sqrt{2}.\ u_1 = \frac{1}{\sqrt{3}}(1,1,1)^T, \qquad u_2 = \frac{1}{\sqrt{2}}(1,-1,0)^T.$

3.

$A = \begin{bmatrix} 1/\sqrt{3} & 1/\sqrt{2} \\ 1/\sqrt{3} & -1/\sqrt{2} \\ 1/\sqrt{3} & 0 \end{bmatrix} \begin{bmatrix} \sqrt{3} & 0 \\ 0 & \sqrt{2} \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$.

4. Complete the basis with

$u_3 = (1,1,-2)^T/\sqrt{6}$.

Full column rank gives $N(A) = \{0\}$, while $\dim N(A^T) = 3 - 2 = 1$. The vector u_3 is a unit left-nullspace basis vector: $u_1^T u_3 = u_2^T u_3 = 0$. Thus $U = [u_1\ u_2\ u_3]$ has $U^TU = I_3$, and multiplying the full $U\Sigma V^T$ reconstructs both columns of A. The full rectangular diagonal factor is

$\begin{bmatrix} \sqrt{3} & 0 \\ 0 & \sqrt{2} \\ 0 & 0 \end{bmatrix}$.

Lin Cing
https://github.com/xinhi6716/computer-science-self-study

CH07-E2

1.

$m = 3, n = 2, r = 2$. $\dim C(A^T) = 2, \quad \dim N(A) = 0, \dim C(A) = 2, \quad \dim N(A^T) = 1.$

2. Bases:
   - $C(A^T)$:e_1, e_2;
   - $N(A)$ is the zero subspace, whose basis is the empty set, not a set containing zero.
   - $C(A)$:u_1, u_2;
   - $N(A^T)$:u_3.

3.

$Av_1 = \sqrt{3}u_1, \qquad Av_2 = \sqrt{2}u_2.$

4. The three left vectors form an orthonormal basis of $\mathbb{R}^3$, giving the column-space and left-nullspace sum. The row space already equals $\mathbb{R}^2$, with trivial nullspace.

CH07-E3

1. Columns dependent, rank 1:

$u_1 = \frac{1}{\sqrt{5}}(1,2)^T, \qquad v_1 = \frac{1}{\sqrt{5}}(1,2)^T$. $A = 5u_1v_1^T, \qquad \sigma_1 = 5.$

2.

$5u_1v_1^T = A.$

3.

$\|A\|_2 = 5, \qquad \|A\|_F = 5$. $A^+ = \frac{1}{5}v_1u_1^T = \frac{1}{25}\begin{bmatrix}1 & 2\\ 2 & 4\end{bmatrix}.$

4.

$x^+ = A^+b = \frac{7}{25}(1,2)^T.$

Normal equation:

$x_1 + 2x_2 = \frac{7}{5}.$

All minimizers are

$x = x^+ + t(-2,1)^T, \quad t \in \mathbb{R}.$

The added vector lies in the nullspace and is orthogonal to x^+. Pythagoras proves that x^+ uniquely minimizes the norm.

Lin Cing https://github.com/xinhi6716/computer-science-self-study

CH07-E4

1. Every finite real rectangular matrix has an SVD. Here $|\det A| = 20$ equals the product of its two singular values, so neither is zero and A is invertible.
2. For a full-rank square matrix,

$\kappa_2(A) = \frac{\sigma_1}{\sigma_2} \geq 1.$

3. A positive singular value becoming zero makes the matrix rank deficient and singular. The ordinary inverse condition number becomes infinite. Changes in the other singular values depend on the perturbation.

CH07-E5

1. Three singular values are positive, so the rank is 3.
2. Spectral errors:

$k = 1 : 4, \qquad k = 2 : 1, \qquad k = 3 : 0.$

Frobenius errors:

$k = 1 : \sqrt{17}, \qquad k = 2 : 1, \qquad k = 3 : 0.$

3. Retained energy:

$\frac{\sigma_1^2 + \cdots + \sigma_k^2}{\sigma_1^2 + \cdots + \sigma_r^2}.$

4. The spectral error is σ_{k+1}. Since $\sigma_2 = 4$, rank 1 suffices, while rank 0 has error 9. Thus the minimum is $k = 1$.

Lin Cing https://github.com/xinhi6716/computer-science-self-study

CH07-E6

1. Each row sums to zero, so the samples are centered.
2. $X_0X_0^T = \begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix}$. $u_1 = (1,1)^T/\sqrt{2}, \qquad u_2 = (1,-1)^T/\sqrt{2}.$

Scores:

$s_1 = (-\sqrt{2}, 1/\sqrt{2}, 1/\sqrt{2}), s_2 = (0, -1/\sqrt{2}, 1/\sqrt{2}).$


3. $\widehat{C} = \frac{1}{2}\begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix}$.


The first and second component sample variances are $3/2$ and $1/2$, respectively. The first explains $3/4$ of the total variance.

4. Scaling row 2 by 100 makes its variance dominate, moving the first direction toward the second coordinate axis. Standardization avoids domination by units but replaces covariance geometry by correlation geometry. That choice depends on the application.

Projections and translation

5. The projections are (−1,−1), (1/2,1/2), (1/2,1/2). The residuals are (0,0), (−1/2,1/2), (1/2,−1/2). Their squared norms total 1, exactly the squared discarded singular value.





6. Centering subtracts the added mean μ, leaving the same matrix. The principal direction is unchanged; the best affine line becomes μ+span(u_1). All projections translate by μ, so the residuals and minimum error remain unchanged.

Lin Cing https://github.com/xinhi6716/computer-science-self-study

CH07-E7

1.

$A = \sqrt{10}\, u_1 v_1^T, \quad u_1 = (1,2)^T/\sqrt{5}, \quad v_1 = (1,1)^T/\sqrt{2}.$

2.

$A^+ = \frac{1}{\sqrt{10}} v_1 u_1^T = \frac{1}{10}\begin{bmatrix} 1 & 2 \\ 1 & 2 \end{bmatrix}. \; x^+ = A^+ b = \frac{7}{10}(1,1)^T. \; AA^+ b = \frac{7}{5}(1,2)^T.$

3. Residual:

$e = \left(-\frac{2}{5}, \frac{1}{5}\right)^T. \; e^T(1,2) = 0.$

4. All minimizers:

$x = x^+ + t(-1,1)^T, \quad t \in \mathbb{R}.$

The minimum-norm solution lies in row space, perpendicular to nullspace. Adding a nonzero nullspace vector increases its squared norm.

CH07-E8

1. Insert $V^T V = I$ into the SVD.

$A = U\Sigma V^T = (UV^T)(V\Sigma V^T) = QS.$

2. A product of orthogonal square matrices is orthogonal.

$Q^T Q = I. \; S^T = V\Sigma^T V^T = S.$

For any x,

$x^T S x = \sum_i \sigma_i (V^T x)_i^2 \geq 0.$

3. Invertibility makes every singular value positive, proving positive definiteness and invertibility of S.

4.

$\det A = \det Q \det S.$

Since $\det S > 0$ in the invertible case, $\operatorname{sgn}(\det A) = \operatorname{sgn}(\det Q)$. A negative determinant of Q records orientation reversal.

Lin Cing https://github.com/xinhi6716/computer-science-self-study

CH07-E9

1.

$AA^+ = U_r U_r^T, \qquad A^+A = V_r V_r^T.$

Both matrices are symmetric, and

$(U_r U_r^T)^2 = U_r U_r^T,$

The same calculation applies to $V_r V_r^T$. Hence both are orthogonal projectors.

2.

$AA^+A = U_r U_r^T A = A.\ A^+AA^+ = V_r V_r^T A^+ = A^+.$

3. If the cutoff splits a tie, different retained directions within the tied subspace give the same optimal error. The complete tied subspace remains fixed, but individual axes and the retained part of it need not be unique.

4. Flipping only u_1 changes the first term from

$\sigma_1 u_1 v_1^T$

to its negative. Flipping v_1 as well supplies a second minus sign and preserves the factorization.

Lin Cing
https://github.com/xinhi6716/computer-science-self-study

CH07-E10

1. Eckart–Young gives $\sigma_i(A) \le \|A - C\|_2$ for every C of rank less than i. Choose $C = B_{i-1}$. Then $\sigma_i(A) \le \|A - B\|_2 + \|B - B_{i-1}\|_2 = \|A - B\|_2 + \sigma_i(B)$. Exchanging A, B and combining the inequalities proves the bound.
2. The first two singular values exceed 10^{-10}, so every compatible B has rank at least 2. The rank-2 model $B_2 = \mathrm{diag}(1, 10^{-8}, 0)$ has distance 10^{-12}; the rank-3 model $B_3 = A$ has distance zero. Both are compatible, so the third direction is unresolved. A truncated pseudoinverse retaining the second direction still amplifies it by 10^8; retention also requires an application error budget.
3. $\sigma_1 \ge \|Re_2\|_2 = \sqrt{10^{12} + 1} > 10^6$, and $\sigma_1\sigma_2 = |\det R| = 1$, so $\sigma_2 < 10^{-6}$. Without computing an SVD, an explicit rank-1 model is

$$B = \begin{bmatrix} 1 & 10^6 \\ 10^{-6} & 1 \end{bmatrix} = \begin{bmatrix} 10^6 \\ 1 \end{bmatrix} \begin{bmatrix} 10^{-6} & 1 \end{bmatrix}.$$

Only the lower-left entry changes, so $\|R - B\|_2 = 10^{-6}$. Since R is already upper triangular, ordinary QR permits $Q = I$. Its diagonal entries are both 1 despite its proximity to deficient rank. QR diagonals cannot replace singular values.

4. If $\sigma_i(A) > t + \delta$, then $\sigma_i(B) \ge \sigma_i(A) - \delta > t$. If $\sigma_i(A) < t - \delta$, then $\sigma_i(B) \le \sigma_i(A) + \delta < t$. Within the band, perturbations may cross the threshold; equality depends on the strict or nonstrict convention. If the threshold itself changes with matrix scale, include that change as well.

CH07-E11

Successive coordinates are (1,0), (4/5,−3/5), (12/5,−6/5), (12/5,6/5). The final multiplication gives ((36+24)/25, (48−18)/25)=(12/5,6/5). Directly, $A = \frac{1}{5}\begin{bmatrix} 12 & -1 \\ 6 & 12 \end{bmatrix}$, whose first column gives the same answer. Norms are 1,1,6/$\sqrt{}$5,6/$\sqrt{}$5: both orthogonal factors preserve length, while Σ changes it.

For arbitrary unit x, put w=V^Tx. Then $w_1^2+w_2^2=1$ and $\|Ax\|^2 = 9w_1^2 + 4w_2^2 = 4 + 5w_1^2$. Hence $2 \le \|Ax\| \le 3$. Equality at 3 requires $w_2=0$ and $w_1=\pm1$, equivalent to $x=\pm v_1$; equality at 2 requires $w_1=0$ and $w_2=\pm1$, equivalent to $x=\pm v_2$. Here both w components are nonzero, so the output length lies strictly between the two singular values. Singular values are the lengths attained along right singular directions, not a list of all possible output lengths.

CH07-E12

1. Unit-input error versus a squared sum

> $A_1 = \mathrm{diag}(8,0,0)$, $E = \mathrm{diag}(0,3,2)$. For every unit x,
>
> $\|Ex\|^2 = 9x_2^2 + 4x_3^2 \le 9(x_1^2 + x_2^2 + x_3^2) = 9.$
>
> Input e₂ attains equality, so $\|E\|_2 = 3$. Meanwhile $\|E\|_F^2 = 0 + 9 + 4 = 13$. Input e₂+e₃ has length $\sqrt{2}$, not one. Normalization gives output length $\sqrt{13/2} < 3$, consistent with the spectral bound.

CH07-E12 (continued)

2. A strict gap does not force spectral uniqueness

> $\|A - B_c\|_2 = \max\{|8-c|, 3, 2\}, \qquad \|A - B_c\|_F^2 = (8-c)^2 + 13.$
>
> The global spectral lower bound is three. This family attains it exactly when $|8 - c| \le 3$, or $5 \le c \le 11$. Every such c is nonzero, so B has rank one. The global Frobenius bound $\sqrt{13}$ is attained within this family only at c=8. We have classified only diagonal candidates Bc, not excluded other spectral minimizers.

Lin Cing
https://github.com/xinhi6716/computer-science-self-study

CH07-E12 (continued)

3. A kernel witness for a nondiagonal candidate

> Take $z = (1,-1,0)^T/\sqrt{2}$. Direct multiplication gives Bz=0 and $Az = (8,-3,0)^T/\sqrt{2}$, hence
>
> $\|(A-B)z\|^2 = \|Az\|^2 = 73/2 \ge 9.$
>
> Since z is unit, $\|A - B\|_2 \ge \sqrt{73/2} > 3$: this B is not optimal. For the general bound, the kernel witness must be in a specific subspace. If $\mathrm{rank}\, B \le 1$, then $\dim \ker B \ge 2$, so its intersection with $\mathrm{span}\{e_1, e_2\}$ is nonzero. For a unit z in the intersection, $Bz = 0$ and $\|Az\|^2 = 64z_1^2 + 9z_2^2 \ge 9$; hence $\|A - B\|_2 \ge 3$.

CH07-E12 (continued)

4. Global uniqueness in Frobenius norm

Let P project onto C(B), with rank at most one. Set $\alpha_i = \|Pe_i\|^2$. Then $0 \le \alpha_i \le 1$ and their sum is at most one. Thus

$\mathrm{tr}(PAA^T) = 64\alpha_1 + 9\alpha_2 + 4\alpha_3 \le 64\alpha_1 + 9(1-\alpha_1) \le 64.$

Attaining squared Frobenius error 13 requires trace 64. This forces α_1=1 and then α_2=α_3=0. Therefore P projects onto span(e_1). Columnwise Pythagoras,

$\|A-B\|_F^2 = \|(I-P)A\|_F^2 + \|PA-B\|_F^2,$

also forces its second term to zero. Hence B=PA=diag(8,0,0), uniquely among all allowed matrices, not merely among the diagonal family.

CH07-E12 (continued)

5. A cutoff through a repeated singular value

Let $w_\theta^\perp = (-\sin\theta, \cos\theta, 0)^T$. In the orthonormal basis $w_\theta, w_\theta^\perp, e_3$, D acts by 3,3,1 and Cθ by 3,0,0. The error therefore acts by 0,3,1:

$\|D-C_\theta\|_2 = 3, \qquad \|D-C_\theta\|_F = \sqrt{3^2+1^2} = \sqrt{10}.$

Both attain the lower bounds. Angles zero and π/2 retain different axes and produce different matrices. Angles θ and θ+π give the same projector, so not every angle produces a distinct matrix. The complete singular subspace for value three is always span(e_1,e_2); its selected one-dimensional part changes.

Lin Cing https://github.com/xinhi6716/computer-science-self-study

CH07-E13

A is 1×2 with rank 1, hence full row rank; B is 2×1 with rank 1, hence full column rank. Since $AB = [1]$, $(AB)^+ = [1]$.

The full-row-rank formula gives $A^+ = A^T(AA^T)^{-1} = [1\ 1]^T/2$; the full-column-rank formula gives $B^+ = (B^TB)^{-1}B^T = [1\ 0]$. Thus $B^+A^+ = [1/2] \neq (AB)^+$. All dimensions match, but maximum rank of each factor does not ensure the reverse-order law.

References and attribution

Lin Cing, handwritten linear algebra notes. Gilbert Strang, Introduction to Linear Algebra, 5th ed., Chapter 7.

Original educational synthesis by Lin Cing, licensed under CC BY 4.0.
https://github.com/xinhi6716/computer-science-self-study

Lin Cing
https://github.com/xinhi6716/computer-science-self-study