

The Intersection of Food Fortification Technology, Nutrition Equity, Food Safety Risks, and the Role of Food Labelling – A Policy Framework

Food fortification of staple foods remains one of the most cost-effective public health interventions for addressing micronutrient deficiencies. However, when mandatory staple fortification coexists with aggressive marketing of multiple fortified processed foods, significant challenges around equity, safety, and consumer information emerge.

Food Fortification Technology and Its Benefits

Large-scale fortification of staples such as maize flour, wheat flour, oil, and salt has successfully reduced deficiencies in vitamin A, iron, iodine, and folate in many countries. Technological advances allow precise addition of micronutrients that are stable during processing and cooking. This passive approach reaches broad populations without requiring major changes in dietary behaviour.

Nutrition Equity Concerns

Despite its success, fortification does not benefit all groups equally. Urban populations with higher consumption of industrially processed staples often receive greater benefits, while rural and low-income households relying on informally milled or home-produced staples may remain underserved. This creates equity gaps.

Simultaneously, the proliferation of voluntarily fortified processed snacks, beverages, and cereals targeted at middle-class consumers can lead to excessive intake among certain groups. Surveys showing both overconsumption in some populations and persistent deficiency in others highlight the double-edged nature of multiple fortification sources.

Food Safety Risks

Excessive intake of certain micronutrients carries risks. High vitamin A intake, for example, can lead to hypervitaminosis A, particularly when fortification overlaps with supplementation programmes. Iron overload is another concern in some populations. Inconsistent labelling — with unclear quantification of added nutrients — makes it difficult for consumers and health professionals to track total intake.

The Role of Food Labelling

Inconsistent and ambiguous labelling significantly undermines informed choice. Many fortified products use vague health claims without disclosing exact nutrient levels or distinguishing between naturally occurring and added micronutrients. This lack of transparency prevents consumers from making balanced decisions and complicates dietary monitoring at population level.

Proposed Policy Framework

A comprehensive policy framework should balance benefits and risks through the following pillars:

Harmonized Fortification Standards — Establish clear, evidence-based fortification levels for both mandatory staples and voluntary products, considering total dietary intake and upper tolerable limits.

Improved Labelling Requirements — Mandate clear, standardized labelling that specifies the type and amount of added micronutrients (e.g., “Fortified with X mg of iron per 100g”). Prohibit ambiguous health claims that could mislead consumers.

Strengthened Monitoring and Surveillance — Invest in robust post-market surveillance systems to track both deficiency and excess. Regular population-level biomarker surveys should guide adjustments to fortification levels.

Equity-Focused Implementation — Ensure fortification vehicles reach vulnerable populations. Support small-scale millers with appropriate technology and subsidies to improve coverage in rural areas.

Risk-Benefit Communication — Launch public education campaigns explaining both the benefits of fortification and the importance of dietary diversity, discouraging over-reliance on processed fortified products.

Regulatory Coordination — Create clear institutional mandates between ministries of health, agriculture, trade, and regulatory bodies to prevent policy fragmentation.

References – Question One

Lane, M. M., et al. (2024). Ultra-processed food exposure and adverse health outcomes: Umbrella review of epidemiological meta-analyses. *BMJ*, 384, e077310.

Seto, T., et al. (2025). Food additives: Emerging detrimental roles on gut health. *Food Frontiers*.

Jarmakiewicz-Czaja, S., et al. (2025). Effects of selected food additives on the gut microbiome and metabolic dysfunction-associated steatotic liver disease. *Medicina*, 61(2), 192.

Wang, H., et al. (2024). The key to intestinal health: A review and perspective on food additives, intestinal flora, and intestinal health. *Frontiers in Nutrition*, 11, 1420358.

De Siena, M., et al. (2022). Food emulsifiers and metabolic syndrome: The role of gut microbiota. *Nutrients*, 14(16), 3414.

Monteiro, C. A., et al. (2017). The UN Decade of Nutrition, the NOVA food classification and the trouble with ultra-processing. *Public Health Nutrition*, 21(1), 5–17.

Bopape, M., et al. (2022). Effect of different front-of-package food labels on identification of unhealthy products and intention to purchase. *Appetite*, 179, 106289.

World Health Organization. (2023). WHO framework on the marketing of foods and non-alcoholic beverages to children.

References – Question Two

World Health Organization & Food and Agriculture Organization of the United Nations. (2006). Guidelines on food fortification with micronutrients. WHO Press.

Rohner, F., et al. (2023). Global coverage of mandatory large-scale food fortification: A review. *Nutrients*, 15(19), 4280.

Allen, L., de Benoist, B., Dary, O., & Hurrell, R. (Eds.). (2006). Guidelines on food fortification with micronutrients. WHO/FAO.

World Health Organization. (n.d.). Food fortification. Retrieved from <https://www.who.int/health-topics/food-fortification>