

Data Science with Actuarial Applications**Week 7**

Xintong Li

Department of Statistics and Actuarial Science

Last Week

- ▶ Y is a key ratio: claim frequency or claim severity

Last Week

- ▶ Y is a key ratio: claim frequency or claim severity
- ▶ X is a vector of rating factors, modeled as categorical variables

Last Week

- ▶ Y is a key ratio: claim frequency or claim severity
- ▶ X is a vector of rating factors, modeled as categorical variables
- ▶ Exponential Dispersion Models (EDM)

Last Week

- ▶ Y is a key ratio: claim frequency or claim severity
- ▶ X is a vector of rating factors, modeled as categorical variables
- ▶ Exponential Dispersion Models (EDM)
- ▶ Multiplicative models (logarithmic link function)

Last Week

- ▶ Y is a key ratio: claim frequency or claim severity
- ▶ X is a vector of rating factors, modeled as categorical variables
- ▶ Exponential Dispersion Models (EDM)
- ▶ Multiplicative models (logarithmic link function)
- ▶ Claim frequency: Poisson regression

Last Week

- ▶ Y is a key ratio: claim frequency or claim severity
- ▶ X is a vector of rating factors, modeled as categorical variables
- ▶ Exponential Dispersion Models (EDM)
- ▶ Multiplicative models (logarithmic link function)
- ▶ Claim frequency: Poisson regression
- ▶ Claim severity: Gamma regression

Today

- ▶ Estimating the parameters of a GLM from:

Today

- ▶ Estimating the parameters of a GLM from:
- ▶ Writing the log-likelihood function

Today

- ▶ Estimating the parameters of a GLM from:
- ▶ Writing the log-likelihood function
- ▶ Taking the derivative of the log-likelihood function

Today

- ▶ Estimating the parameters of a GLM from:
- ▶ Writing the log-likelihood function
- ▶ Taking the derivative of the log-likelihood function
- ▶ Solving for the maximum likelihood estimators (MLE)

2.6 Parameter Estimation (MLE)



Taking derivative using the chain rule



Evaluating each partial derivative



Remark: saturated model



Corresponding GLM overfits. However, it is useful in the definition of deviance.

Multiplicative Poisson frequency model



Multiplicative gamma severity model



Summary of Theoretical Results

- ▶ **Goal:** estimate the parameters of a GLM using MLE

Summary of Theoretical Results

- ▶ **Goal:** estimate the parameters of a GLM using MLE
- ▶ The $(r + 1)$ equations:

$$\sum_{i=1}^n w_i \frac{y_i - \mu_i}{v(\mu_i) g'(\mu_i)} x_{ij} = 0.$$

Summary of Theoretical Results

- ▶ **Goal:** estimate the parameters of a GLM using MLE
- ▶ The $(r + 1)$ equations:

$$\sum_{i=1}^n w_i \frac{y_i - \mu_i}{v(\mu_i) g'(\mu_i)} x_{ij} = 0.$$

- ▶ For multiplicative Poisson frequency model:

$$\sum_{i=1}^n w_i (y_i - \mu_i) x_{ij} = 0.$$

Summary of Theoretical Results

- **Goal:** estimate the parameters of a GLM using MLE
- The $(r + 1)$ equations:

$$\sum_{i=1}^n w_i \frac{y_i - \mu_i}{v(\mu_i) g'(\mu_i)} x_{ij} = 0.$$

- For multiplicative Poisson frequency model:

$$\sum_{i=1}^n w_i (y_i - \mu_i) x_{ij} = 0.$$

- For multiplicative gamma severity model:

$$\sum_{i=1}^n w_i \frac{y_i - \mu_i}{\mu_i} x_{ij} = 0.$$

Example: Moped dataset

- **Goal:** Use everything we learned to build a GLM for the moped dataset.

Example: Moped dataset

- ▶ **Goal:** Use everything we learned to build a GLM for the moped dataset.
- ▶ **Tasks:**
 1. Load the data

Example: Moped dataset

- ▶ **Goal:** Use everything we learned to build a GLM for the moped dataset.
- ▶ **Tasks:**
 1. Load the data
 2. Explore the data

Example: Moped dataset

- ▶ **Goal:** Use everything we learned to build a GLM for the moped dataset.
- ▶ **Tasks:**
 1. Load the data
 2. Explore the data
 3. Set the base tariff cells

Example: Moped dataset

- ▶ **Goal:** Use everything we learned to build a GLM for the moped dataset.
- ▶ **Tasks:**
 1. Load the data
 2. Explore the data
 3. Set the base tariff cells
 4. Fit the frequency model

Example: Moped dataset

- ▶ **Goal:** Use everything we learned to build a GLM for the moped dataset.
- ▶ **Tasks:**
 1. Load the data
 2. Explore the data
 3. Set the base tariff cells
 4. Fit the frequency model
 5. Fit the severity model

Recap from Last Time

- ▶ Estimating the parameters of a GLM using MLE

Recap from Last Time

- ▶ Estimating the parameters of a GLM using MLE
- ▶ Example in R: Moped dataset

Today

- ▶ Hypothesis testing for GLMs

Today

- ▶ Hypothesis testing for GLMs
- ▶ Deviance statistic: likelihood ratio test

Today

- ▶ Hypothesis testing for GLMs
- ▶ Deviance statistic: likelihood ratio test
- ▶ Pearson's chi-squared test

Today

- ▶ Hypothesis testing for GLMs
- ▶ Deviance statistic: likelihood ratio test
- ▶ Pearson's chi-squared test
- ▶ Estimating the dispersion parameter ϕ

Today

- ▶ Hypothesis testing for GLMs
- ▶ Deviance statistic: likelihood ratio test
- ▶ Pearson's chi-squared test
- ▶ Estimating the dispersion parameter ϕ
- ▶ Testing hierarchical models

Today

- ▶ Hypothesis testing for GLMs
- ▶ Deviance statistic: likelihood ratio test
- ▶ Pearson's chi-squared test
- ▶ Estimating the dispersion parameter ϕ
- ▶ Testing hierarchical models
- ▶ Confidence intervals based on Fisher's information

2.7 GLM Model Building



Deviance Statistic from Likelihood Ratio Test



Deviance Statistic from Likelihood Ratio Test

Null hypothesis:

H_0 : fitted GLM model

Deviance Statistic from Likelihood Ratio Test

Null hypothesis:

H_0 : fitted GLM model

Test Statistic:

$$D^* = 2\{\mathfrak{l}(y_1, \dots, y_n; \theta, \phi) - \mathfrak{l}(\hat{\mu}_1, \dots, \hat{\mu}_n; \theta, \phi)\}.$$

Deviance Statistic from Likelihood Ratio Test

Null hypothesis:

$$H_0 : \text{fitted GLM model}$$

Test Statistic:

$$D^* = 2\{l(y_1, \dots, y_n; \theta, \phi) - l(\hat{\mu}_1, \dots, \hat{\mu}_n; \theta, \phi)\}.$$

Under general conditions, the deviance statistic D^* is approximately chi-squared distributed with $n - p$ degrees of freedom, where

- ▶ n is the number of observations,
- ▶ and $p = r + 1$ is the number of parameters in the model.

Deviance Statistic from Likelihood Ratio Test

Null hypothesis:

$$H_0 : \text{fitted GLM model}$$

Test Statistic:

$$D^* = 2\{l(y_1, \dots, y_n; \theta, \phi) - l(\hat{\mu}_1, \dots, \hat{\mu}_n; \theta, \phi)\}.$$

Under general conditions, the deviance statistic D^* is approximately chi-squared distributed with $n - p$ degrees of freedom, where

- ▶ n is the number of observations,
- ▶ and $p = r + 1$ is the number of parameters in the model.

Compare the deviance statistic with $c_{1-\alpha}$,

- ▶ $c_{1-\alpha}$ is the $1 - \alpha$ quantile of the chi-squared distribution with $n - p$ degrees of freedom.
- ▶ Reject if $D^* > c_{1-\alpha}$, accept otherwise.

Deviance Statistic from Likelihood Ratio Test

Null hypothesis:

$$H_0 : \text{fitted GLM model}$$

Test Statistic:

$$D^* = 2\{l(y_1, \dots, y_n; \theta, \phi) - l(\hat{\mu}_1, \dots, \hat{\mu}_n; \theta, \phi)\}.$$

Under general conditions, the deviance statistic D^* is approximately chi-squared distributed with $n - p$ degrees of freedom, where

- ▶ n is the number of observations,
- ▶ and $p = r + 1$ is the number of parameters in the model.

Compare the deviance statistic with $c_{1-\alpha}$,

- ▶ $c_{1-\alpha}$ is the $1 - \alpha$ quantile of the chi-squared distribution with $n - p$ degrees of freedom.
- ▶ Reject if $D^* > c_{1-\alpha}$, accept otherwise.

Using **p-values**, reject if **p-value** $< \alpha$, accept otherwise.

Pearson's Goodness-of-Fit Test

Null hypothesis: H_0 : fitted GLM model

$$\chi^2 = \sum_{i=1}^n \frac{(y_i - \hat{\mu}_i)^2}{\text{Var}(Y_i)} = \frac{1}{\phi} \sum_{i=1}^n w_i \frac{(y_i - \hat{\mu}_i)^2}{v(\hat{\mu}_i)}.$$

χ^2 is approximately chi-squared distributed with $n - (r + 1)$ degrees of freedom.

Pearson's Goodness-of-Fit Test

Null hypothesis: H_0 : fitted GLM model

$$X^2 = \sum_{i=1}^n \frac{(y_i - \hat{\mu}_i)^2}{\text{Var}(Y_i)} = \frac{1}{\phi} \sum_{i=1}^n w_i \frac{(y_i - \hat{\mu}_i)^2}{v(\hat{\mu}_i)}.$$

X^2 is approximately chi-squared distributed with $n - (r + 1)$ degrees of freedom.

Compare the Pearson test statistic with $c_{1-\alpha}$,

- ▶ $c_{1-\alpha}$ is the $1 - \alpha$ quantile of the chi-squared distribution with $n - p$ degrees of freedom.
- ▶ Reject if $X^2 > c_{1-\alpha}$, accept otherwise.

Pearson's Goodness-of-Fit Test

Null hypothesis: H_0 : fitted GLM model

$$X^2 = \sum_{i=1}^n \frac{(y_i - \hat{\mu}_i)^2}{\text{Var}(Y_i)} = \frac{1}{\phi} \sum_{i=1}^n w_i \frac{(y_i - \hat{\mu}_i)^2}{v(\hat{\mu}_i)}.$$

X^2 is approximately chi-squared distributed with $n - (r + 1)$ degrees of freedom.

Compare the Pearson test statistic with $c_{1-\alpha}$,

- ▶ $c_{1-\alpha}$ is the $1 - \alpha$ quantile of the chi-squared distribution with $n - p$ degrees of freedom.
- ▶ Reject if $X^2 > c_{1-\alpha}$, accept otherwise.

Using p-values, **reject if p-value $< \alpha$, accept otherwise.**

Estimation of ϕ

- **Method 1:** Matching moments with the Pearson statistic

$$\hat{\phi} = \frac{1}{n - (r + 1)} \sum_{i=1}^n w_i \frac{(y_i - \hat{\mu}_i)^2}{v(\hat{\mu}_i)}.$$

where we set $X^2 = n - (r + 1)$ by matching the expected value of X^2

$$\mathbb{E}[X^2] = n - (r + 1).$$

Estimation of ϕ

- **Method 1:** Matching moments with the Pearson statistic

$$\hat{\phi} = \frac{1}{n - (r + 1)} \sum_{i=1}^n w_i \frac{(y_i - \hat{\mu}_i)^2}{v(\hat{\mu}_i)}.$$

where we set $X^2 = n - (r + 1)$ by matching the expected value of X^2

$$\mathbb{E}[X^2] = n - (r + 1).$$

- **Method 2:** Matching moments with the deviance statistic

$$\hat{\phi}_D = \frac{D}{n - (r + 1)}.$$

where we set $D^* = n - (r + 1)$ by matching the expected value of D^*

Testing Hierarchical Models

Consider two models $M_s \subset M_t$:

H_0 : data comes from M_s

H_α : data comes from the more complicated M_t

Testing Hierarchical Models

Consider two models $M_s \subset M_t$:

H_0 : data comes from M_s

H_α : data comes from the more complicated M_t

Test statistic:

$$D^*(\mathbf{y}, \hat{\mu}^{(s)}) - D^*(\mathbf{y}, \hat{\mu}^{(t)}).$$

Testing Hierarchical Models

Consider two models $M_s \subset M_t$:

H_0 : data comes from M_s

H_α : data comes from the more complicated M_t

Test statistic:

$$D^*(\mathbf{y}, \hat{\mu}^{(s)}) - D^*(\mathbf{y}, \hat{\mu}^{(t)}).$$

- M_s and M_t are required to be from the same EDM family.

Testing Hierarchical Models

Consider two models $M_s \subset M_t$:

H_0 : data comes from M_s

H_α : data comes from the more complicated M_t

Test statistic:

$$D^*(\mathbf{y}, \hat{\mu}^{(s)}) - D^*(\mathbf{y}, \hat{\mu}^{(t)}).$$

- ▶ M_s and M_t are required to be from the same EDM family.
- ▶ $D^*(\mathbf{y}, \hat{\mu}^{(s)}) - D^*(\mathbf{y}, \hat{\mu}^{(t)})$ is approximately chi-squared distributed with $p_t - p_s$ degrees of freedom.

Testing Hierarchical Models

Consider two models $M_s \subset M_t$:

H_0 : data comes from M_s

H_α : data comes from the more complicated M_t

Test statistic:

$$D^*(\mathbf{y}, \hat{\mu}^{(s)}) - D^*(\mathbf{y}, \hat{\mu}^{(t)}).$$

- ▶ M_s and M_t are required to be from the same EDM family.
- ▶ $D^*(\mathbf{y}, \hat{\mu}^{(s)}) - D^*(\mathbf{y}, \hat{\mu}^{(t)})$ is approximately chi-squared distributed with $p_t - p_s$ degrees of freedom.
- ▶ Using p-values, reject if p-value $< \alpha$, accept otherwise.

Confidence intervals based on Fisher's information

To get the confidence intervals, we need the Fisher's information matrix:

$$[I]_{jk} = -\mathbb{E} \left[\frac{\partial^2 \ell}{\partial \beta_j \partial \beta_k} \right], \quad j, k = 0, 1, \dots, r.$$

Confidence intervals based on Fisher's information

To get the confidence intervals, we need the Fisher's information matrix:

$$[I]_{jk} = -\mathbb{E} \left[\frac{\partial^2 \ell}{\partial \beta_j \partial \beta_k} \right], \quad j, k = 0, 1, \dots, r.$$

It can be shown that $I = X^T D X$, where D is a diagonal matrix with diagonal elements

$$d_i = \frac{w_i}{\phi(\hat{\mu}_i)(g'(\hat{\mu}_i)^2)}.$$

Confidence intervals based on Fisher's information

To get the confidence intervals, we need the Fisher's information matrix:

$$[I]_{jk} = -\mathbb{E} \left[\frac{\partial^2 \ell}{\partial \beta_j \partial \beta_k} \right], \quad j, k = 0, 1, \dots, r.$$

It can be shown that $I = X^T D X$, where D is a diagonal matrix with diagonal elements

$$d_i = \frac{w_i}{\phi(\hat{\mu}_i)(g'(\hat{\mu}_i))^2}.$$

Asymptotically, we have $\hat{\beta} \sim N(\beta, I^{-1})$. Thus we can find the [confidence intervals for \$\beta_j\$](#) :

$$\hat{\beta}_j \pm z_{1-\alpha/2} \sqrt{[I^{-1}]_{jj}}.$$

where $z_{1-\alpha/2}$ is the $(1 - \alpha/2)$ -quantile of the standard normal distribution.

Confidence intervals based on Fisher's information

In practice, we also want to find the confidence intervals for μ_i .

Confidence intervals based on Fisher's information

In practice, we also want to find the confidence intervals for μ_i .

We do this by finding the confidence intervals for η_i :

$$\hat{\eta}_i \pm z_{1-\alpha/2} \sqrt{[I_{\eta}^{-1}]_{jj}}.$$

where $I_{\eta}^{-1} = (x_{i0}, x_{i1}, \dots, x_{ir}) I^{-1} (x_{i0}, x_{i1}, \dots, x_{ir})^T$ is the variance of $\hat{\eta}_i$.

Confidence intervals based on Fisher's information

In practice, we also want to find the confidence intervals for μ_i .

We do this by finding the confidence intervals for η_i :

$$\hat{\eta}_i \pm z_{1-\alpha/2} \sqrt{[I_{\eta}^{-1}]_{jj}}.$$

where $I_{\eta}^{-1} = (x_{i0}, x_{i1}, \dots, x_{ir}) I^{-1} (x_{i0}, x_{i1}, \dots, x_{ir})^T$ is the variance of $\hat{\eta}_i$.

Then we can find the confidence intervals for μ_i :

$$(e^{\hat{\eta}_i - z_{1-\alpha/2} \sqrt{[I_{\eta}^{-1}]_{jj}}}, e^{\hat{\eta}_i + z_{1-\alpha/2} \sqrt{[I_{\eta}^{-1}]_{jj}}}).$$

Example: Moped dataset

- **Goal:** Test our built GLM for the moped dataset.

Example: Moped dataset

► **Goal:** Test our built GLM for the moped dataset.

► **Tasks:**

1. Test the frequency model using deviance test using its p-value.
2. Test the severity model using deviance test using its p-value.
3. Test the frequency model using Pearson's chi-squared test using its p-value.
4. Test the severity model using Pearson's chi-squared test using its p-value.
5. Estimate ϕ for the severity model.
6. Combine zone 5, 6, 7 into zone 4. Fit a new frequency model with the new zone 4.
7. Test the new frequency model against the previous one.
8. Repeat step 6 and 7 for the severity model.
9. Find the 95% confidence intervals for the parameters and the relativities of the frequency model.
10. Repeat step 9 for the severity model.