

# A Survey in Theta Graph

Xinyu Luo (xl3147)

## I. History

The story of spanners is actually started by a bunch of people who wanted to simplify the complete graph. In 1986, Chew tried to simplify a graph to a planar graph, and he succeeded with a transmission distance between each pair of vertices at most  $\sqrt{10}$  times of the euclidean distance based on Delaunay Triangulation, and an algorithm with time complexity  $O(n \log n)$  to build this graph [5]. Well, in 1988, J. Keil mentioned an approach to simplify the complete graph which is better than Delaunay Approximating, this approach is based on the theta graph [3]. In 1991, J. Ruppert and R. Seidel came with another good bound for approximating complete euclidean graphs, which is also based on theta graph but with a completely different proving method, and the graph building algorithm was extended to d-dimensional [1]. And, in 1992, J. Keil came back with a summary of approximating the complete Euclidean graph without realizing there was a better bound for theta graph that came out one year ago [2]. Then, the spanners topic was staying in silence for 15 years.

With the flourishing of the mobile industry, the requirement for networks became higher, so people cared about this topic again. In 2007, a textbook by G. Narasimhan et. called “Geometric Spanner Networks” came out, then wiki named the definition in this book as the official definition for spanners [10]. Then, in these years many researchers came with tighter upper bounds for spanners and clearer proof about the previous results. The new era for this topic was coming [6, 7, 8, 9].

## II. Different proofs

The different classes of spanners can be divided by the polygon approach and theta graph. The first one is mainly the Chew algorithm which is already shown in course notes. For the second class, there are some different proofs J. Keil (1988), J. Ruppert and R. Seidel (1991), clearer proof for J. Keil is from G. Narasimhan (2007), clearer proof for J. Ruppert and R. Seidel is from a lot of people my reference is from A. Verdone (2015) [1, 2, 3, 5, 10].

### A. J. Keil (1988) and G. Narasimhan (2007) proof for spanning ratio $\frac{1}{\cos \theta - \sin \theta}$

They got this result for  $k > 8$ . This constraint is because when  $k > 8$  it can make sure that the new distance is smaller than the old distance to the target proved by Trig. Thus, there will be no cycle on the path.

The original version is J. Keil version, this proof is 3 pages long, the clearer proof is given by G. Narasimhan and it is one page.

- B. J. Ruppert and R. Seidel (1991) and A. Verdonchot (2015) proof for spanning ratio

$$\frac{1}{1-2\sin\frac{\theta}{2}}$$

The spanning ratio is improved from  $\frac{1}{\cos\theta - \sin\theta}$  to  $\frac{1}{1-2\sin\frac{\theta}{2}}$ , based on another proof based on triangle inequality. And also, they improved  $k > 8$  to  $k > 6$ . They found this is because Keil's constructing theta graph algorithm required  $k > 6$ . They said Keil said this, but I didn't find this was told in Keil's paper. I will write some of my ideas about why it is  $k > 6$  at the end of the proof.

### J. Ruppert and R. Seidel (1991):

Construct a path from p to q: greedy algorithm (PS: In the original paper, it said if there is a direct edge between  $p_i$  and  $q$  connect it, otherwise, connect the vertex in the cone which  $p_i$  and  $q$  belongs to has the closest orthogonal projection onto the axis of that cone. I think it is the greedy algorithm.) [1]

**Def.** Let the path between p and q is  $p_0 p_1 \dots p_i q$ , the  $i$ th p is  $p_i$ ,  $e_i$  is the edge from  $p_{i-1}$  to  $p_i$ . Let  $l_i$  be the euclidean distance between  $p_i$  and  $q$  ( $\text{dist}(p_i, q)$ ),  $l_{i-1}$  be the  $\text{dist}(p_{i-1}, q)$ , there are  $m$  edges on the path.

We get the formula  $|e_i| + l_i \leq l_{i-1} + 2|e_i|\sin\frac{\theta}{2}$ , then by telescoping sum, we get routing ratio  $\frac{1}{1-2\sin\frac{\theta}{2}}$ .

Prove formula 1:

If  $p_i = q$ , trivially this hold.

Otherwise, there are three cases.

**Def.** Let  $\alpha$  be the angle between ray  $p_{i-1}p_i$  and the cone axis. Let  $\beta$  be the angle between  $p_{i-1}q$  and the cone axis.  $r$  is the symmetry vertex of  $p_i$  from the cone axis.

Case 1:  $\beta < \alpha$  (Figure 1.)

First,  $p_{i-1}$  is not directly connected to  $q$  which means  $q$  has to be at the right side of  $p_i$ .

Then,

$$|p_{i-1}r| \leq |p_{i-1}w| + |wr| \text{ and } |p_iq| \leq |p_iw| + |wq|$$

$$\Rightarrow |p_{i-1}r| + |p_iq| \leq |p_{i-1}q| + |p_i r|$$

$$\Rightarrow |e_i| + l_i \leq l_{i-1} + 2|e_i|\sin\alpha \leq l_{i-1} + 2|e_i|\sin\frac{\theta}{2}$$




Figure 1. Case 1

Case 2:  $\beta \geq \alpha$ ,  $p_i$  and  $q$  are on the same side (Figure 2.)

First, make a circle  $C_0$  with the center  $p_{i-1}$  and radius  $|p_{i-1}p_i|$ , the intersection point with ray  $p_{i-1}q$  is  $r$ , then  $s$  is an image cross cone axis of  $r$ . Make a circle  $C$  with the center  $q$  and radius  $|p_iq|$ .

If we get  $|p_{i-1}p_i| + |p_iq| \leq |p_{i-1}q| + |rs|$ , then  $|e_i| + l_i \leq l_{i-1} + 2|e_i|\sin\beta \leq l_{i-1} + 2|e_i|\sin\frac{\theta}{2}$

We have  $|p_{i-1}p_i| = |p_{i-1}q| - |rq|$ , we need to prove  $|p_iq| \leq |rq| + |rs|$ .

For  $|qs| \leq |rq| + |rs|$ , only need to prove  $|p_iq| \leq |qs|$ .

We notice that  $p_{i-1}$  and  $q$  are the center for the two circle, so the intersect points of  $C$  and  $C_0$   $p_i$  and  $w$  should be on the different side of ray  $p_{i-1}q$ , and one of the intersection is  $p_i$ , thus,  $s$  must outside the circle  $C$ . We get  $|p_iq| \leq |qs|$ .

Case 3:  $\beta \geq \alpha$ ,  $p_i$  and  $q$  are on the different side (Figure 2.)

The proof is the same as case 2.




Figure 2. Case 2 and Case 3 (figure from [1])

About why  $k > 6$ ?

To be honest, the author thought out this spanning ratio based on  $k > 6$ . We already know  $p_i$  is on the left side of  $q$  and we have  $\frac{\angle p_{i-1}p_iq}{p_{i-1}q} = \frac{\angle p_{i-1}q}{p_iq}$ , thus,

$\angle p_{i-1}p_iq > \angle p_ip_{i-1}q$  when  $k > 6$ , so  $|p_iq| < |p_{i-1}q|$ . There will not be a cycle. Also, the spanning ratio becomes infinity when  $k = 6$  and negative when  $k < 6$ .

**Clearer proof in some recent paper (2015) which is very different from 1991 paper:**

This proof comes from a proof for Yao’s graph. [4]

**Def.** A path from  $a$  to  $b$ ,  $c$  is the only intermediate point. Rotate  $c$  towards  $ab$ , using  $a$  as the center of the circle and  $|ac|$  as the radius for  $\theta$  degrees, then get  $c'$ , as figure 3 showed on the next page. We call the arc from  $c$  to  $c'$  is  $arc\ C$ .

Proof for spanning ratio  $\frac{1}{1-2\sin\frac{\theta}{2}}$ :

If  $cc'$  and  $ab$  are intersected,

we get  $|ac'| \leq |as| + |sc'|$  and  $|bc| \leq |bs| + |sc|$ .

$$\Rightarrow |ac| + |bc| \leq |as| + |sc'| + |bs| + |sc| = |ab| + |cc'| = |ab| + 2\sin\frac{\theta}{2}|ac|,$$

Then, it is the same as the formula 1, we get  $|e_i| + l_i \leq l_{i-1} + 2|e_i|\sin\frac{\theta}{2}$ .




Figure 3. Rotating proof (figure from [4])

We need to prove  $cc'$  and  $ab$  is always intersected. (This proof in the paper is not very clear, the proof below is rewritten by myself).

There will be 2 cases:  $|ac| \leq |ab|$  and  $|ac| > |ab|$ .

Case 1:  $|ac| \leq |ab|$ .

We know that  $\text{arc } C$  will intersect with  $ab$ . And the  $\text{arc } C$ 's chord  $cc'$  must intersect with  $ab$  for the rotating angle  $\theta$  less than  $\pi$ .

Case 2:  $|ac| > |ab|$ , this time *arc*  $C$  will not intersect with  $ab$ .

We select  $c$  instead of  $b$  is because  $c$  is on the left side of  $b$ . Thus, Case 2 could happen if and only if the angle  $\sigma$  between  $ac$  and cone axis is larger than the angle  $\gamma$  between  $ab$  and the cone axis, as shown in the figure 4.




Figure 4.

Thus, if we make a mirror point  $c''$  of cone axis for  $c$ .  $cc''$  is must on the left of  $b$ , and  $\angle c''ac = 2\sigma < \theta$ , thus,  $c'$  is on the left of  $b$ , and the different side of  $ab$ . Also,  $cc'$  will on the right side of  $a$  when  $\theta < \pi$ . Thus,  $cc'$  has to intersect with  $ab$ .

#### C. $\Theta_4 - 4$ , $\Theta_5 - 5$ , $\Theta_6 - 6$

We have already got all the spanning ratio for  $k > 6$ . And for  $\Theta_2$  and  $\Theta_3$ , they are proved to be not a spanner by the translate prove for  $Y_{ao} - 2$  and  $Y_{ao} - 3$ ,  $Y_2$  and  $Y_3$  are not spanners is proved by Nawar M. El Molla in his Ph.D. thesis in 2009. For  $\Theta_4\Theta_5\Theta_6$ , they are all spanners, but they are proved different ways.

##### ➤ $\Theta_4$ is a spanner:

Proved by L. Barba, P. Bose, et al on paper "On the stretch factor of the Theta-4 graph" in 2012. The approximating spanning ratio for  $\Theta_4$  is very loose, it is around 237. [7]

##### ➤ $\Theta_5$ is a spanner - this is a hard topic, it is the last one solved in $\Theta_4\Theta_5\Theta_6$ :

First mentioned by J. Ruppert and R. Seidel (1991), the same persons who get spanning ratio  $\frac{1}{1-2\sin \frac{\pi}{k}}$ . They said in the conclusion of their paper: "In the planar case, some improvement can be made on the constants. In particular, when  $k$  is odd, there is an asymmetry between the cones at point  $p$  and  $q$  that we can take advantage of by growing paths from both ends. Interestingly, this asymmetry allows us to prove a bound near 10 on the path lengths even for the case  $k = 5$ . For even  $k$ , similar improvements can be made by altering the apertures of some cones at each point." Unfortunately, no details are given.

Then, P. Bose, P. Morin, et al, proved this in 2015. Actually, this paper “The theta-5 graph is a spanner” actually showed in a conference in 2013 and another conference in 2015.[9]

➤  **$\Theta_6$  is a spanner:**

Proved by N. Bonichon, et al in the paper “Connections between theta-graphs, Delaunay triangulations, and orthogonal surfaces”. He found the connection between theta-graphs and DT. The DT part was mentioned in Chew’s paper in 1986.[8]

PS: It is very interesting that when  $Y_k$  graph doesn’t fail as a spanner,  $\Theta_k$  never fail as well, it is still an opening question to prove when  $Y_k$  graph is a spanner and  $\Theta_k$  graph is a spanner as well. [4]

### **III. $O(n \log n)$ time complexity algorithm for building theta graph in 2-dimensional (plane swipe algo)**

The  $O(n \log n)$  algorithm in 2-dimensional space is first mentioned by J. Keil (1988), using the 2-3 tree. And this algorithm is transferred to fit high dimensional space by J. Ruppert and R. Seidel (1991), using the range tree, time complexity  $O(n(\log n)^{d-1})$ . In the book of G. Narasimhan (2007), they said using skip lists makes the algorithm more clear for the high dimensional space. [3, 10]

This algorithm is very abstract in the original paper, and I added a lot of details to make this algorithm seem more reasonable.

**To explain this easier, we assume that the cone we are looking at the cone axis is the x-coordinate (we can do it by rotation for other cones).** Some def. will be shown in Figure 5.

**Def.** The point set on this plane is  $A = \{a_1 \dots a_n\}$ , we sort  $A$  by x-coordinate (it is the same as sorting by the project of the cone axis).

**Def.** For a cone original from the point  $a$ , the upper bound ray for this cone is  $h_1$ , the lower bound ray is  $h_2$ .  $D_1$  is the orthogonal vector of  $h_1$ ,  $D_2$  is the orthogonal vector of  $h_2$ . The point project on  $D_1$  is  $D_1(a_i)$ , the point project on  $D_2$  is  $D_2(a_i)$ .




Figure 5. Project on axis

**Def.**  $P$  a point set equal to  $A$  but sorted by  $D_1(a_i)$ ,  $Q$  is another point set equal to  $A$  but sorted by  $D_2(a_i)$ . For an example of sorting,  $D_1(a_1) < D_1(a_3) < D_1(a_2)$ , then we got the order for  $P$ .

**Def.** BST  $T$ , the leaf nodes of  $T$  represent points in  $Q$ , the rank of the leaf nodes is the same rank as  $Q$ . All the nodes of  $T$  augmenting two values, assume a node  $u$  and  $u$  has both subtrees: 1) value  $z_1$ , info for searching operation:  $u$ 's left subtree's rightmost leaf 2) value  $z_2$ : the point has the lowest rank in  $A$  of  $u$ 's subtree. The example is in the figure below.




Figure 6. The structure of  $T$

1. Initially, we have an empty BST  $T$ .
2. Iterate on  $P$  from  $p_n$  to  $p_1$ : (for example, we are dealing with  $p_i$ )
  - 1) Insert  $p_i$  in  $T$  (update  $T$  at the same time)
  - 2) Search  $p_i$  on  $T$ . For this searching path from the root, if we need to turn left on one node  $v$ , we record  $v$ 's right child to a new set  $M$  (clear  $M$  each iteration). This step will be shown in figure 7 ( $M$  is the nodes in red circle). Then, we notice that we can only collect  $O(\log n)$  nodes in  $M$ , and these nodes in  $M$  represent all nodes on the upper side of  $p_i$ 's  $h_2$ . (Because you can only turn left at node  $v$  only when you are sure all the points on the right subtree of  $v$  is projected on the positive part of  $D_2$ , which mean upper than  $p_i$ 's  $h_2$ ).




Figure 7. Searching process

- 3) If  $M$  is not empty, select the minimum  $z_2$  (for example  $a_i$ ) in all nodes in  $M$ . This step helps as to select the point with the minimum x-coordinate in all points which are upper than  $p_i$ 's  $h_2$ . (It is all points because non-leaf nodes can represent the minimum  $z_2$  of its subtrees.)
- 4) Connect  $p_i$  with  $a_i$ . Continue the next iteration.

Extra proof for why this is correct:

We use the reverse order in  $P$ , this makes sure that when we are going to connect the edge of  $p_i$  in the cone, we make sure that  $p_{i+1} \dots p_n$  has to be below  $p_i$ 's  $h_1$ , otherwise  $p_{i+1} \dots p_n$  won't be larger than  $p_i$  on  $D_1$ . When we insert  $p_i$  to  $T$ , we get all nodes on the upper side of  $p_i$ 's  $h_2$ , and the nodes in the  $T$  has to be  $p_{i+1} \dots p_n$ . Thus, we get all points in between  $p_i$ 's  $h_1$  and  $h_2$  and stored in  $M$ . After that, we select the point which has minimum x-coordinate, which means the closest point of  $p_i$  in that cone.

Time complexity:  $O(n \log n)$ , for sorting, and  $n$  time BST operations, each cone.



Space:  $O(n)$  for the BST.

**IV. My experiment towards tighter upper bound (a failure for now, will try something more in the future)**

I was doing the experiment for finding counterexamples for the routing ratio and the spanning ratio in the table below. I did find the routing ratio for  $\Theta_5$  fail some cases in  $\Theta_{4k+5}$ . But there is a bound about  $k \geq 1$ , so it is a mistake, to be honest. Then, nothing is different from the table.[6]

**Table 1. Different Spanning Ratio & Routing Ratio**

	Current Spanning	Current Routing	Previous Spanning & Routing
$\theta_{(4k+2)}$ -graph	$1 + 2 \sin\left(\frac{\theta}{2}\right)$	$\frac{1}{1 - 2 \sin\left(\frac{\theta}{2}\right)}$	$\frac{1}{1 - 2 \sin\left(\frac{\theta}{2}\right)}$
$\theta_{(4k+3)}$ -graph	$\frac{\cos\left(\frac{\theta}{4}\right)}{\cos\left(\frac{\theta}{2}\right) - \sin\left(\frac{3\theta}{4}\right)}$	$1 + \frac{2 \sin\left(\frac{\theta}{2}\right) \cos\left(\frac{\theta}{4}\right)}{\cos\left(\frac{\theta}{2}\right) - \sin\left(\frac{\theta}{2}\right)}$	$\frac{1}{1 - 2 \sin\left(\frac{\theta}{2}\right)}$
$\theta_{(4k+4)}$ -graph	$1 + \frac{2 \sin\left(\frac{\theta}{2}\right)}{\cos\left(\frac{\theta}{2}\right) - \sin\left(\frac{\theta}{2}\right)}$	$1 + \frac{2 \sin\left(\frac{\theta}{2}\right)}{\cos\left(\frac{\theta}{2}\right) - \sin\left(\frac{\theta}{2}\right)}$	$\frac{1}{1 - 2 \sin\left(\frac{\theta}{2}\right)}$
$\theta_{(4k+5)}$ -graph	$\frac{\cos\left(\frac{\theta}{4}\right)}{\cos\left(\frac{\theta}{2}\right) - \sin\left(\frac{3\theta}{4}\right)}$	$1 + \frac{2 \sin\left(\frac{\theta}{2}\right) \cos\left(\frac{\theta}{4}\right)}{\cos\left(\frac{\theta}{2}\right) - \sin\left(\frac{\theta}{2}\right)}$	$\frac{1}{1 - 2 \sin\left(\frac{\theta}{2}\right)}$

## **Reference**

- [1] Ruppert J, Seidel R. Approximating the d-dimensional complete Euclidean graph. In Proceedings of the 3rd Canadian Conference on Computational Geometry (CCCG 1991) 1991 (pp. 207-210).
- [2] Keil JM, Gutwin CA. Classes of graphs which approximate the complete Euclidean graph. Discrete & Computational Geometry. 1992 Jan 1;7(1):13-28.
- [3] Keil JM. Approximating the complete Euclidean graph. In Scandinavian Workshop on Algorithm Theory 1988 Jul 5 (pp. 208-213). Springer, Berlin, Heidelberg.
- [4] Verdonschot S. Flips and Spanners. arXiv preprint arXiv:1509.02563. 2015 Sep 8.
- [5] Chew, L. Paul. "There Is a Planar Graph Almost as Good as the Complete Graph." (1986).
- [6] Bose P, De Carufel JL, Morin P, van Renssen A, Verdonschot S. Towards tight bounds on theta-graphs: More is not always better. Theoretical Computer Science. 2016 Feb 22;616:70-93.
- [7] Barba L, Bose P, De Carufel JL, Van Renssen A, Verdonschot S. On the stretch factor of the Theta-4 graph. In Workshop on Algorithms and Data Structures 2013 Aug 12 (pp. 109-120). Springer, Berlin, Heidelberg.
- [8] Bonichon N, Gavoille C, Hanusse N, Ilcinkas D. Connections between theta-graphs, Delaunay triangulations, and orthogonal surfaces. In International Workshop on Graph-Theoretic Concepts in Computer Science 2010 Jun 28 (pp. 266-278). Springer, Berlin, Heidelberg.
- [9] Bose P, Morin P, Van Renssen A, Verdonschot S. The  $\theta_5$ -graph is a spanner. Computational Geometry. 2015 Feb 1;48(2):108-19.
- [10] Narasimhan G, Smid M. Geometric spanner networks. Cambridge University Press; 2007 Jan 8.